

**SPECIFICATION****TITLE OF INVENTION**

NITROGEN-CONTAINING AROMATIC DERIVATIVES

Related Applications

5

[0001] This application claims the benefit of U.S. Provisional Application No.60/464690, filed April 22, 2003.

BACKGROUND OF THE INVENTION

10

Field of the Invention

15

[0002] The present invention relates to novel compounds effective for prevention and treatment of various diseases associated with abnormal angiogenesis, and to the medical compositions such as angiogenesis inhibitors and antitumor agents containing the novel compounds.

Related Background of the Invention

20

[0003] Angiogenesis is an essential biological phenomenon for fetal vascular formation and morphological and functional development of organs. New blood vessels are assembled through several processes including endothelial cell migration, proliferation and tube formation, and the participation of mast cells, lymphocytes, interstitial cells and the

25

like has been shown to be important in this process
(non-patent literature 1).

5 [0004] A multiple in vivo angiogenesis-stimulating
factors have been identified, particularly Vascular
Endothelial Growth Factor (hereinafter abbreviated as
"VEGF") and Fibroblast Growth Factor (hereinafter
abbreviated as "FGF") are reported to enhance
angiogenesis (non-patent literature 2 and 3).

10

[0005] Although physiological angiogenesis occurs at
the time of healing of wound or in a female estrous
cycle in adult individuals, it is known that
pathological increase in angiogenesis in adult
15 individuals is involved in onset or progression of
various disease. Specific diseases associated with
abnormal angiogenesis include cancer, rheumatoid
arthritis, atherosclerosis, diabetic retinopathy,
angioma, psoriasis, and the like (non-patent literature
20 4). In particular, a literature has indicated
angiogenesis dependency for solid tumor growth, and
angiogenesis inhibitors are therefore promising as new
therapeutic agents for intractable solid tumors (non-
patent literature 5).

25

[0006] Patent literature 1 and 2 are provided as prior

arts with regard to 6-membered nitrogen-containing aromatic derivatives bonded with substituted indole.

5 [0007] Although patent literature 1 describes indole derivatives which suppress VEGF-stimulated angiogenesis based on a selective tyrosine kinase inhibition, the pharmacological test results on their inhibition action are not disclosed. Although patent literature 2 describes pyridine derivatives bonded with indole ring
10 via an oxygen atom at the 4-position, neither the compound according to the present invention nor their inhibiting actions on FGF-stimulated angiogenesis are disclosed.

15 [0008] [patent literature 1] WO 02/16348
[patent literature 2] WO 02/32872
[non-patent literature 1] J. Biol. Chem., 267,
10931, 1992.
[non-patent literature 2] Endocrinology, 133,
20 848, 1993.
[non-patent literature 3] Biochem. Biophys. Res. Commun., 147, 876, 1987.
[non-patent literature 4] N. Engl. J. Med., 333,
1757, 1995.
25 [non-patent literature 5] J. Natl. Cancer Inst.,
82, 4, 1990.


SUMMARY OF THE INVENTION

[0009] It is an object of the present invention to investigate and discover angiogenesis-inhibiting compounds which: (1) exhibit antitumor activity by strongly suppressing both of angiogenesis included by VEGF and FGF which are major *in vivo* angiogenesis factors, (2) are highly useful as drug materials in terms of their properties, biokinetics and safety, and (3) are useful for amelioration, prevention and treatment of various diseases associated with abnormal increase in angiogenesis.

[0010] As a result of much diligent research in light of the circumstances described above, the present inventors have succeeded in synthesizing novel pyridine derivatives and pyrimidine derivatives represented by the following general formula (I) , salts thereof, or hydrates of the foregoing. At the same time, the inventors have completed the present invention upon discovering that these compounds, the salts thereof, or the hydrates of the foregoing exhibit an excellent angiogenesis-inhibiting effect.

[0011] Specifically, the present invention provides the followings:


<1> a compound represented by the general formula:



wherein X_1 represents a nitrogen atom or a group represented by the formula $-CR_{10}=$, X_2 represents a
 5 nitrogen atom or a group represented by the formula $-CR_{11}=$, and X_1 and X_2 do not represent a nitrogen atom at the same time;


Y represents an oxygen atom, a sulfur atom, a sulfinyl group, a sulfonyl group, or a group represented by the
 10 formula $-NR_Y-$ (wherein R_Y represents a hydrogen atom or a C_{1-6} alkyl group);

R_1 represents an optionally substituted C_{1-6} alkoxy group, an optionally substituted C_{6-10} aryloxy group, a group represented by the formula $-NR_{12a}R_{12b}$, a group
 15 represented by the formula:




(wherein Y_{A1} and Y_{A2} each independently represent a group represented by the formula $-A_{10}-A_{11}-A_{12}$ (wherein
 20 A_{10} represents a single bond or an optionally substituted C_{1-6} alkylene; A_{11} represents a single bond, an oxygen atom, a carbonyl group or a sulfonyl group;

and A₁₂ represents a hydrogen atom, a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₂₋₆ alkynyl group, a C₃₋₈ cycloalkyl group, a C₆₋₁₀ aryl group, a 5- to 10-membered heteroaryl group, a group represented by the formula -NR_{A10}R_{A11}, a group represented by the formula -OR_{A12} (wherein R_{A10}, R_{A11} and R_{A12} each independently represent a hydrogen atom, a C₁₋₆ alkyl group or C₃₋₈ cycloalkyl group) or a group represented by the formula:




(wherein e represents 1 or 2; Z represents an oxygen atom, a group represented by the formula -CR_{X7}R_{X8}- or a group represented by the formula -NR_{X9}-; R_{X7}, R_{X8} and R_{X9} each independently represent a hydrogen atom, a hydroxyl group or a C₁₋₆ alkyl group)); and Y_{A3} represents a hydrogen atom or an optionally substituted C₁₋₆ alkyl group) or a group represented by the formula:




(wherein T1 represents an optionally substituted 5- to 10- membered aromatic heterocycle which may have X in the ring or an optionally substituted 3- to 10-membered heterocycle which may have X in the ring); R₃, R₄, R₅, R₆, R₇, R₈, R₁₀ and R₁₁ each independently represent a hydrogen atom, a halogen atom, a cyano

group, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₂₋₆ alkynyl group, an optionally substituted C₃₋₈ cycloalkyl group, a group represented by the formula -CO-R₁₃, a group represented by the formula -NR₁₄-CO-R₁₃, a group represented by the formula -SO₂-R₁₅, a group represented by the formula -NR₁₄-SO₂-R₁₅, or a group represented by the formula -NR_{16a}R_{16b}; R₉ represents a group represented by the formula -NR_{16a}R_{16b} or a group represented by the formula:



(wherein T2 represents an optionally substituted 5- to 10- membered aromatic heterocycle or an optionally substituted 3- to 10- membered heterocycle); R_{12a} and R_{12b} each independently represent a hydrogen atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₃₋₆ alkenyl group, an optionally substituted C₃₋₆ alkynyl group, an optionally substituted C₃₋₈ cycloalkyl group, an optionally substituted 3- to 10- membered heterocyclic group, or an optionally substituted C₁₋₆ alkoxy group; R₁₃ represents a hydrogen atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₂₋₆ alkynyl group, an optionally substituted C₃₋₈ cycloalkyl

group, an optionally substituted C₆₋₁₀ aryl group, an optionally substituted 5- to 10- membered heteroaryl group, an optionally substituted 3- to 10- membered heterocyclic group, an optionally substituted C₁₋₆ alkoxy group, an optionally substituted C₆₋₁₀ aryloxy group, a group represented by the formula -NR_{12a}R_{12b}, or a group represented by the formula:




(wherein T2 represents an optionally substituted 5- to 10- membered aromatic heterocycle or an optionally substituted 3- to 10- membered heterocycle);

R₂ and R₁₄ each independently represent a hydrogen atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₂₋₆ alkynyl group, an optionally substituted C₃₋₈ cycloalkyl group, or a group represented by the formula -CO-R₁₃;


R₁₅ represents an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₂₋₆ alkynyl group, an optionally substituted C₃₋₈ cycloalkyl group, an optionally substituted C₆₋₁₀ aryl group, an optionally substituted 5- to 10- membered heteroaryl group, or an optionally substituted 3- to 10- membered heterocyclic group;

R_{16a} and R_{16b} each independently represent a hydrogen

atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₃₋₆ alkenyl group, an optionally substituted C₃₋₆ alkynyl group, an optionally substituted C₃₋₈ cycloalkyl group, an optionally substituted C₆₋₁₀ aryl group, an optionally substituted 5- to 10- membered heteroaryl group, an optionally substituted 3- to 10- membered heterocyclic group, or an optionally substituted C₁₋₆ alkoxy group; and X represents an oxygen atom, a sulfur atom, a carbonyl group, a sulfonyl group, a group represented by the formula -CR_{X1}R_{X2}-, or a group represented by the formula -NR_{X3}- (wherein R_{X1}, R_{X2} and R_{X3} each independently represent a hydrogen atom or a group represented by the formula -A₁-A₂-A₃ (wherein A₁ and A₂ each independently represent a single bond, an optionally substituted C₁₋₆ alkylene group or a carbonyl group; and A₃ represents a hydrogen atom, a C₃₋₈ cycloalkyl group, a group represented by the formula -NR_{A1}R_{A2}, or the formula -OR_{A3} (wherein, R_{A1}, R_{A2} and R_{A3} each independently represent a hydrogen atom or a C₁₋₆ alkyl group), or an optionally substituted group represented by the formula:



(wherein a represents 1 or 2))),
a salt thereof, or a hydrate of the foregoing;
<2> a compound represented by the general formula:



wherein X_1 , X_2 , Y , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 and R_9 represent the same definitions as X_1 , X_2 , Y , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 and R_9 in <1>, respectively,

5 a salt thereof, or a hydrate of the foregoing;

<3> a compound according to <1> or <2>, a salt of the compound, or a hydrate of the foregoing, wherein Y represents an oxygen atom, a group represented by the formula $-NH-$, or a group represented by the formula $-$

10 $N(CH_3)-$;

<4> a compound according to <1> or <2>, a salt of the compound, or a hydrate of the foregoing, wherein Y represents an oxygen atom;

<5> a compound according to any of <1> to <4>, a salt of the compound, or a hydrate of the foregoing, wherein one of X_1 and X_2 represents a group represented by the formula $-CH=$ and the other represents a nitrogen atom;

<6> a compound according to any of <1> to <4>, a salt of the compound, or a hydrate of the foregoing, wherein both X_1 and X_2 represent a group represented by the formula $-CH=$;


20 <7> a compound according to any of <1> to <6>, a salt

of the compound, or a hydrate of the foregoing, wherein R_3 , R_4 , R_5 , R_6 and R_8 each represent a hydrogen atom, and R_7 represents a hydrogen atom, a halogen atom or an optionally substituted C_{1-6} alkyl group;

5 <8> a compound according to any of <1> to <7>, a salt of the compound, or a hydrate of the foregoing, wherein R_9 represents a group represented by the formula $-NHR_{17}$ (wherein R_{17} represents an optionally substituted C_{1-6} alkyl group, a C_{3-6} alkynyl group, a C_{3-8} cycloalkyl group, an optionally substituted C_{6-10} aryl group or an optionally substituted 5- to 10- membered heteroaryl group);

10 <9> a compound according to any of <1> to <7>, a salt of the compound, or a hydrate of the foregoing, wherein R_9 represents a group represented by the formula $-NR_{18a}R_{18b}$ (wherein R_{18a} and R_{18b} each independently represent a C_{1-6} alkyl group);

15 <10> a compound according to any of <1> to <7>, a salt of the compound, or a hydrate of the foregoing, wherein R_9 represents a group represented by the formula:



(wherein b_1 represents 1 or 2; X represents the same definition as X in <1>);

20 <11> a compound according to any of <1> to <7>, a salt of the compound, or a hydrate of the foregoing, wherein

R₉ represents a group represented by the formula -NHR₁₉ (wherein R₁₉ represents a C₁₋₆ alkyl group, a C₃₋₆ alkynyl group, a C₃₋₈ cycloalkyl group or a C₆₋₁₀ aryl group);

5 <12> a compound according to any of <1> to <11>, a salt of the compound, or a hydrate of the foregoing, wherein R₃, R₄, R₅, R₆, R₇ and R₈ each represent a hydrogen atom;

<13> a compound according to any of <1> to <12>, a salt of the compound, or a hydrate of the foregoing, wherein

10 R₂ represents a hydrogen atom;

<14> a compound according to any of <1> to <13>, a salt of the compound, or a hydrate of the foregoing, wherein


R₉ represents a group represented by the formula -NHR₂₀ (wherein R₂₀ represents a methyl group, an ethyl group

15 or a cyclopropyl group);

<15> a compound according to any of <1> to <13>, a salt of the compound, or a hydrate of the foregoing, wherein

R₉ represents a group represented by the formula -NH(CH₃);


20 <16> a compound according to any of <1> to <15>, a salt of the compound, or a hydrate of the foregoing, wherein R₁ represents a further optionally substituted group represented by the formula:



25 (wherein b₂ represents 0, 1 or 2; and X represents the

same definition as X in <1>;


<17> a compound according to any of <1> to <16>, a salt of the compound, or a hydrate of the foregoing, wherein R_1 represents a group represented by the formula:



(wherein X represents the same definition as X in <1>;

<18> a compound according to <17>, a salt of the compound, or a hydrate of the foregoing, wherein X in the formula (IV) represents an oxygen atom;


10 <19> a compound according to <17>, a salt of the compound, or a hydrate of the foregoing, wherein X in the formula (IV) represents a group represented by the formula:




15 (wherein R_{X4} represents a hydrogen atom or a group represented by the formula $-A_4-A_5-A_6$ (wherein A_4 and A_5 each independently represent a single bond, an optionally substituted C_{1-6} alkylene or a carbonyl group; and A_6 represents a hydrogen atom, a C_{3-8} cycloalkyl group or a group represented by the formula $-NR_{A4}R_{A5}$ or the formula $-OR_{A6}$ (wherein R_{A4} , R_{A5} and R_{A6} each independently represent a hydrogen atom or a C_{1-6} alkyl group))));

20 <20> a compound according to <17>, a salt of the

compound, or a hydrate of the foregoing, wherein X in the formula (IV) represents a group represented by the formula:




5 (wherein R_{X5} and R_{X6} each independently represent a hydrogen atom or a group represented by the formula - $A_7-A_8-A_9$ (wherein A_7 and A_8 each independently represent a single bond, an optionally substituted C_{1-6} alkylene group or a carbonyl group; and A_9 represents a hydrogen atom,
10 a C_{3-8} cycloalkyl group, a group represented by the formula $-NR_{A7}R_{A8}$, or the formula $-OR_{A9}$ (wherein R_{A7} , R_{A8} , and R_{A9} each independently represent a hydrogen atom or a C_{1-6} alkyl group), or a group represented by the formula:



15 (wherein c_1 represents 0, 1 or 2));
<21> a compound according to <20>, a salt of the compound, or a hydrate of the foregoing, wherein one of R_{X5} and R_{X6} in the formula (VI) represents a hydroxyl group and the other represents a hydrogen atom or a C_{1-6} alkyl group;
20


<22> a compound according to <20>, a salt of the compound, or a hydrate of the foregoing, wherein one of R_{X5} or R_{X6} in the formula (VI) represents a hydrogen

atom and the other represents a group represented by the formula:




(wherein c_2 represents 1 or 2);

- 5 <23> a compound according to any of <1> to <16>, a salt of the compound, or a hydrate of the foregoing, wherein R_1 represents a group represented by the formula:




- (wherein R_{X51} and R_{X61} each independently represent a hydrogen atom or a group represented by the formula -
 10 $A_{71}-A_{81}-A_{91}$ (wherein A_{71} and A_{81} each independently represent a single bond, an optionally substituted C_{1-6} alkylene group or a carbonyl group; and A_{91} represents a hydrogen atom, a C_{3-8} cycloalkyl group, a group
 15 represented by the formula $-NR_{A71}R_{A81}$, or the formula $-OR_{A91}$ (wherein R_{A71} , R_{A81} , and R_{A91} each independently represent a hydrogen atom or a C_{1-6} alkyl group), or a group represented by the formula:




- 20 (wherein c_{11} represents 0, 1 or 2));
- <24> A compound according to any of <1> to <15>, a salt of the compound, or a hydrate of the foregoing, wherein

R₁ represents a group represented by the formula:




(wherein Y_{A1} and Y_{A2} each independently represent a group represented by the formula -A₁₀-A₁₁-A₁₂ (wherein A₁₀ represents a single bond or an optionally substituted C₁₋₆ alkylene group; A₁₁ represents a single bond, an oxygen atom, a carbonyl group, or a sulfonyl group; and A₁₂ represents a hydrogen atom, a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₂₋₆ alkynyl group, a C₃₋₈ cycloalkyl group, a C₆₋₁₀ aryl group, a 5- to 10-membered heteroaryl group, a group represented by the formula -NR_{A10}R_{A11}, or the formula -OR_{A12} (wherein, R_{A10}, R_{A11} and R_{A12} each independently represent a hydrogen atom, a C₁₋₆ alkyl group or a C₃₋₈ cycloalkyl group), or a group represented by the formula:



(wherein e represents 1 or 2; and Z represents an oxygen atom or a group represented by the formula -CR_{X7}R_{X8}- or the formula -NR_{X9}- (wherein R_{X7}, R_{X8} and R_{X9} each independently represent a hydrogen atom, a hydroxyl group or a C₁₋₆ alkyl group)); and Y_{A3} represents a hydrogen atom or an optionally substituted C₁₋₆ alkyl group);


<25> a compound according to <24>, a salt of the

compound, or a hydrate of the foregoing, wherein one of Y_{A1} and Y_{A2} in the formula (VIII) represents a hydrogen atom and the other represents a group represented by the formula $-(CH_2)_2-A_{13}-A_{14}$ (wherein A_{13} represents a single bond, a carbonyl group or a sulfonyl group; and A_{14} represents a C_{1-6} alkyl group, a group represented by the formula $-NR_{A13}R_{A14}$ (wherein R_{A13} and R_{A14} each independently represent a hydrogen atom, a C_{1-6} alkyl group or a C_{3-8} cycloalkyl group), or a group represented by the formula:




(wherein e and Z represent the same definitions as e and Z in <24>, respectively)))); and Y_{A3} in the formula (VIII) represents a hydrogen atom;

<26> a compound according to any of <1> to <15>, a salt of the compound, or a hydrate of the foregoing, wherein R_1 represents a group represented by the formulas:




(each of the foregoing members being optionally substituted with a group selected from Substituent Group Alpha,

wherein Substituent Group Alpha is a group consisting of a halogen atom, a hydroxyl group, a thiol group, a nitro group, a cyano group, a carboxyl group, an amino group, a C₁₋₆ alkyl group, a C₃₋₈ cycloalkyl group, and a group represented by the formulas:



(wherein R_{N1} and R_{N2} each independently represent a hydrogen atom or a C₁₋₆ alkyl group));


<27> a compound according to any of <1> to <15>, a salt of the compound, or a hydrate of the foregoing, wherein R₁ represents a group represented by the formulas:



<28> a compound according to any of <1> to <15>, a salt

of the compound, or a hydrate of the foregoing, wherein


R₁ represents a group represented by the formulas:



<29> a compound according to any of <1> to <15>, a salt

5 of the compound, or a hydrate of the foregoing, wherein


R₁ represents a group represented by the formulas:




<30> a compound according to <1> or <2>, a salt of the

compound, or a hydrate of the foregoing, wherein the

10 compound is represented by the general formula:




(wherein R_1 represents a group represented by the formulas:



5 (each of the foregoing members being optionally substituted with a group selected from Substituent Group Beta,

wherein Substituent Group Beta is a group consisting of a hydroxyl group, a C_{1-6} alkyl group, a C_{3-8} cycloalkyl group, and a group represented by the formulas:

10



(wherein R_{N1} and R_{N2} each independently represent a hydrogen atom or a C_{1-6} alkyl group)); and R_9 represents a group represented by the formula $-NHR_{20}$ (wherein R_{20}

represents a methyl group, an ethyl group or a cyclopropyl group));

<31> a compound according to <1>, a salt of the compound, or a hydrate of the foregoing, wherein the compound is a compound selected from a group consisting of

(1) N1-ethyl-5-(2-((methoxylamino)carbonyl)amino-4-pyrimidyl)oxy-1H-indolecarboxamide,

(2) 5-(6-(3-(3-diethylaminopropylamino)ureido)pyrimidin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,

(3) 5-(6-(((4-hydroxypiperidin-1-yl)carbonyl)amino)-pyrimidin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,

(4) 5-(6-((4-pyrrolidin-1-yl)piperidin-1-yl)carbonylamino)pyrimidin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,

(5) 5-(2-(3-((1R)-1-carbamoyl-2-phenylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,

(6) 5-(2-(3-((1S)-1-carbamoyl-2-phenylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,

(7) 5-(2-(3-(2-oxo-2-(pyrrolidin-1-yl)ethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,

- (8) 5-(2-(3-(2-(4-hydroxy-4-methylpiperidin-1-yl)-2-oxoethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- (9) 5-(2-(3-((1S)-1-carbamoylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- (10) 5-(2-(3-((1S)-1-carbamoyl-3-methylbutyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- (11) 5-(2-(3-carbamoylmethylureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- (12) 5-(2-(3-cyclopropylcarbamoylmethylureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- (13) 5-(2-(3-((1S)-1-carbamoyl-2-hydroxyethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- (14) 5-(2-(3-((1R)-1-carbamoyl-2-hydroxyethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- (15) (2S)-2-(3-(4-(1-methylcarbamoyl-1H-indol-5-yloxy)pyridin-2-yl)ureido)-1,5-pentanedicarboxylic acid diamide,
- (16) (2S)-2-(3-(4-(1-methylcarbamoyl-1H-indol-5-yloxy)pyridin-2-yl)ureido)succinamide,
- (17) 5-(2-(3-((1S)-1-cyclopropylcarbamoyl-2-hydroxyethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,

- (18) 5-(2-(3-((1S)-1-hydroxymethyl-2-oxo-2-pyrrolidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- 5 (19) 5-(2-(3-((1R)-1-hydroxymethyl-2-oxo-2-pyrrolidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- (20) 5-(2-(3-((1S)-1-hydroxymethyl-2-oxo-2-piperidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- 10 (21) 5-(2-(3-((1R)-1-hydroxymethyl-2-oxo-2-piperidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- (22) 5-(2-(3-((1S)-1-hydroxymethyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- 15 (23) 5-(2-(3-((1S)-1-hydroxymethyl-2-(morpholin-4-yl)-2-oxoethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- (24) 5-(2-(3-(2-cyclopropylcarbamoyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- 20 (25) 5-(2-(3-(3-oxo-3-(pyrrolidin-1-yl)propyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- (26) 5-(2-(3-(3-(4-hydroxy-4-methylpiperidin-1-yl)-3-oxopropyl)ureido)pyridin-4-yloxy)-1H-indole-1-
- 25

carboxylic acid methylamide,

- (27) N1-ethyl-5-(2-(((2-ethoxyethyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- 5 (28) N1-methyl-5-(2-((4-(2-hydroxy-2-methylpropionyl)piperazin-1-yl)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (29) N1-methyl-5-(2-((3-(diethylamino)propylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- 10 (30) N1-methyl-5-(2-(((3-(4-hydroxypiperidino)propyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (31) N1-methyl-5-(2-(((3-(4-methylpiperazin-1-yl)propyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- 15 (32) 5-(2-(3-(4-oxo-4-(pyrrolidin-1-yl)butyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- 20 (33) 5-(2-(3-(3-(cyclopropylcarbamoyl)propyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- (34) 5-(2-(3-(4-(4-hydroxy-4-methylpiperidin-1-yl)-4-oxobutyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- 25 (35) 5-(2-(3-(3-

- (diethylcarbamoyl)propyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- (36) 5-(2-(3-(3-(methylcarbamoyl)propyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- 5 (37) N1-methyl-5-(2-(pyrrolidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (38) N1-methyl-5-(2-(piperidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (39) N1-methyl-5-(2-((4-hydroxypiperidino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- 10 (40) N1-methyl-5-(2-(4-oxopiperidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (41) 5-(2-(((4-hydroxy-4-methylpiperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- 15 (42) N1-methyl-5-(2-((4-(1-hydroxy-1-methylethyl)piperidino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- 20 (43) 5-(2-(((4-(3-methylcarbamoylpropyl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- (44) 5-(2-(((4-(3-carbamoylpropyl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,
- 25 (45) 5-(2-((4-(pyrrolidin-1-yl)carbonyl)piperidin-1-

yl)carbonylamino)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,

(46) N1-methyl-5-(2-(((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-

5 indolecarboxamide,

(47) N1-methyl-5-(2-(((4-(piperidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-

indolecarboxamide,

(48) N1-methyl-5-(2-((4-ethylpiperazin-1-yl)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,

10

(49) N1-methyl-5-(2-((4-(2-hydroxyethyl)piperazin-1-yl)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,

(50) N1-methyl-5-(2-((3-methylsulfonylpropylamino)carbonyl)amino-4-pyridyl)oxy-

15

1H-1-indolecarboxamide,

(51) N1-methyl-5-(2-((4-(2-dimethylaminoacetyl)piperazin-1-yl)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,

(52) N1-methyl-5-(2-((4-cyclohexylpiperazin-1-yl)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,

20

(53) N4-(4-(1-(methylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-4-morpholinecarboxamide,

(54) N1-methyl-5-(2-((1,1-dioxothiomorpholin-4-ylcarbonyl)amino)pyridin-4-yloxy)-1H-1-

25

indolecarboxamide,

(55) 5-(2-(3-((1R)-1-hydroxymethyl-2-oxo-2-pyrrolidin-

- 1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid ethylamide,
- (56) 5-(2-(3-((1S)-1-hydroxymethyl-2-oxo-2-pyrrolidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid ethylamide,
- 5 (57) 5-(2-(3-((1R)-1-hydroxymethyl-2-oxo-2-piperidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid ethylamide,
- (58) 5-(2-(3-((1S)-1-hydroxymethyl-2-oxo-2-piperidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid ethylamide,
- 10 (59) 5-(2-(3-(2-(4-hydroxy-4-methylpiperidin-1-yl)-2-oxoethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid ethylamide,
- 15 (60) N1-ethyl-5-(2-(((1-methyl-4-piperidyl)methyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (61) N1-ethyl-5-(2-(((2-diethylamino)ethyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- 20 (62) N1-ethyl-5-(2-(((2-(morpholin-4-yl)ethyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (63) N1-ethyl-5-(2-(((2-(4-hydroxypiperidino)ethyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- 25

- (64) N1-methyl-5-(2-((2-(4-hydroxypiperidino)ethyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- 5 (65) N1-ethyl-5-(2-((3-(diethylamino)propylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (66) N1-ethyl-5-(2-((3-(morpholin-4-yl)propyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- 10 (67) N1-ethyl-5-(2-((3-(4-methylpiperazin-1-yl)propyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (68) N1-cyclopropyl-5-(2-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide,
- 15 (69) 5-(2-(3-((1R)-1-hydroxymethyl-2-oxo-2-pyrrolidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid cyclopropylamide,
- (70) 5-(2-(3-((1S)-1-hydroxymethyl-2-oxo-2-pyrrolidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid cyclopropylamide,
- 20 (71) 5-(2-(3-(2-oxo-2-(pyrrolidin-1-yl)ethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid cyclopropylamide,
- 25 (72) 5-(2-(3-(3-oxo-3-(pyrrolidin-1-yl)propyl)ureido)pyridin-4-yloxy)-1H-indole-1-

- carboxylic acid cyclopropylamide,
- (73) 5-(2-(3-((1R)-1-hydroxymethyl-2-oxo-2-piperidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid cyclopropylamide,
- 5 (74) 5-(2-(3-((1S)-1-hydroxymethyl-2-oxo-2-piperidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid cyclopropylamide,
- (75) N1-phenyl-5-(2-((3-(diethylamino)propyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- 10 (76) N1-phenyl-5-(2-((3-(4-methylpiperazin-1-yl)propyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (77) N1-ethyl-5-(2-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide,
- 15 (78) 5-(2-((4-hydroxy-4-methylpiperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-indole-1-carboxylic acid ethylamide,
- 20 (79) N1-ethyl-5-(2-((4-hydroxypiperidin-1-yl)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (80) N1-ethyl-5-(2-(piperidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (81) N1-ethyl-5-(2-((pyrrolidin-1-ylcarbonyl)amino)-4-pyridyl)oxy)-1H-1-indolecarboxamide,
- 25 (82) N4-(4-((1-(ethylamino)carbonyl-1H-5-indolyl)oxy)-

- 2-pyridyl)-4-morpholinecarboxamide,
- (83) N1-ethyl-5-(2-((1,1-dioxothiomorpholin-4-ylcarbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide,
- 5 (84) N1-ethyl-5-(2-((methoxylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (85) N1-cyclopropyl-5-(2-((4-hydroxypiperidino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- 10 (86) N1-cyclopropyl-5-(2-((4-hydroxy-4-methylpiperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide,
- (87) N4-(4-(1-(cyclopropylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-4-morpholinecarboxamide,
- 15 (88) N1-cyclopropyl-5-(2-((pyrrolidin-1-ylcarbonyl)amino)-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (89) N1-cyclopropyl-5-(2-(piperidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (90) N4-(4-(1-(cyclopentylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-4-morpholinecarboxamide,
- 20 (91) 5-(2-((4-hydroxypiperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-indole-1-carboxylic acid cyclopentylamide,
- (92) N1-cyclopentyl-5-(2-((4-(pyrrolidin-1-yl)piperidin-1-ylcarbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide,
- 25

- (93) N1-(3-methylbutyl)-5-(2-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide,
- 5 (94) N1-(3-methylbutyl)-5-(2-((4-hydroxypiperidino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (95) N4-(4-(1-((3-methylbutyl)amino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-4-morpholinecarboxamide,
- 10 (96) N1-(1-ethylpropyl)-5-(2-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide,
- (97) N1-(1-ethylpropyl)-5-(2-((4-hydroxypiperidino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- 15 (98) N4-(4-(1-((1-ethylpropyl)amino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-4-morpholinecarboxamide,
- (99) N4-(4-(1-((1-pentyl)amino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-4-morpholinecarboxamide,
- (100) N1-(1-pentyl)-5-(2-((4-hydroxypiperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide,
- 20 (101) N1-(1-pentyl)-5-(2-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide,
- 25 (102) N1-methyl-3-chloro-5-(2-((3-(diethylamino)propyl)amino)carbonyl)amino-4-

pyridyl) oxy-1H-1-indolecarboxamide,

(103) N1-methyl-3-chloro-5-(2-((4-(pyrrolidin-1-yl)piperidino) carbonyl) amino-4-pyridyl) oxy-1H-1-indolecarboxamide,

5 (104) N1-methyl-3-chloro-5-(2-((4-hydroxypiperidino) carbonyl) amino-4-pyridyl) oxy-1H-1-indolecarboxamide,

(105) N1-methyl-3-chloro-5-(2-((3-(4-hydroxypiperidino) propyl) amino) carbonyl) amino-4-

10 pyridyl) oxy-1H-1-indolecarboxamide,

(106) N1-methyl-3-chloro-5-(2-((4-(2-hydroxyethyl) piperazin-1-yl) carbonyl) amino-4-pyridyl) oxy-1H-1-indolecarboxamide,

(107) N4-(4-(3-chloro-1-(methylanino) carbonyl-1H-5-indolyl) oxy-2-pyridyl)-4-morpholinecarboxamide,

15 (108) N1-methyl-3-chloro-5-(2-((4-(ethylpiperazin-1-yl) carbonyl) amino-4-pyridyl) oxy-1H-1-indolecarboxamide,

(109) N1-ethyl-3-chloro-5-(2-((4-hydroxypiperidino) carbonyl) amino-4-pyridyl) oxy-1H-1-

20 indolecarboxamide,

(110) N1-ethyl-3-chloro-5-(2-((3-(4-hydroxypiperidino) propyl) amino) carbonyl) amino-4-pyridyl) oxy-1H-1-indolecarboxamide,

(111) N1-ethyl-3-chloro-5-(2-((3-(diethylamino) propyl) amino) carbonyl) amino-4-pyridyl) oxy-1H-1-indolecarboxamide,

25

- (112) N1,3-dimethyl-5-(2-((4-hydroxypiperidino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (113) N1,3-dimethyl-5-(2-((4-(pyrrolidin-1-yl)piperidino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (114) N1-cyclopropyl-5-(2-((4-hydroxypiperidino)carbonyl)amino-4-pyridyl)oxy-3-methyl-1H-1-indolecarboxamide,
- (115) N1-cyclopropyl-5-(2-((4-(2-hydroxyethyl)piperazin-1-yl)carbonyl)amino-4-pyridyl)oxy-3-methyl-1H-1-indolecarboxamide,
- (116) N1-methyl-5-(2-((methylanino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (117) N1-methyl-5-(2-((diethylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (118) N1-(2-propynyl)-5-(2-((pyrrolidin-1-yl)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide;
- (119) N1-methyl-5-(2-(azetidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (120) N1-ethyl-5-(2-(azetidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (121) N1-cyclopropyl-5-(2-(azetidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (122) N1-methyl-5-(2-(((4-(morpholin-4-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-

indolecarboxamide,

(123) N1-methyl-5-(2-(((4-(azetidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide,

5 (124) N1-methyl-5-(2-(((4-(diethylamino)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide,

(125) N1-methyl-5-(2-(((4-(4-hydroxypiperidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide, and

(126) N1-propyl-5-(2-(pyrrolidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide;

<32> a compound according to <1>, a salt of the compound, or a hydrate of the foregoing, wherein the compound is a compound selected from a group consisting of

(1) 5-(2-(3-(2-oxo-2-(pyrrolidin-1-yl)ethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,

20 (2) 5-(2-(3-carbamoylmethylureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,

(3) 5-(2-(3-((1S)-1-hydroxymethyl-2-oxo-2-pyrrolidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,

25 (4) N1-methyl-5-(2-((4-(2-hydroxy-2-methylpropionyl)piperazin-1-yl)carbonyl)amino-4-

pyridyl)oxy-1H-1-indolecarboxamide,

(5) 5-(2-(3-(4-oxo-4-(pyrrolidin-1-yl)butyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,

5 (6) 5-(2-(3-(3-(cyclopropylcarbamoyl)propyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,

(7) 5-(2-(3-(4-(4-hydroxy-4-methylpiperidin-1-yl)-4-oxobutyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,

(8) 5-(2-(3-(3-(methylcarbamoyl)propyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,

(9) N1-methyl-5-(2-(pyrrolidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,

15 (10) N1-methyl-5-(2-((4-hydroxypiperidino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,

(11) N1-methyl-5-(2-(4-oxopiperidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,

20 (12) 5-(2-(((4-hydroxy-4-methylpiperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide,

(13) 5-(2-(((4-(3-methylcarbamoylpropyl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-indole-1-

25 carboxylic acid methylamide,

(14) 5-(2-(((4-(3-carbamoylpropyl)piperidin-1-

- yl)carbonyl)amino)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methanamide,
- (15) N1-methyl-5-(2-(((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide,
- (16) N1-methyl-5-(2-(((4-(piperidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide,
- (17) N1-methyl-5-(2-((3-methylsulfonylpropylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (18) N4-(4-(1-(methylanino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-4-morpholinecarboxamide,
- (19) N1-cyclopropyl-5-(2-(((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide,
- (20) 5-(2-(((4-hydroxy-4-methylpiperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-indole-1-carboxylic acid ethanamide,
- (21) N1-ethyl-5-(2-((4-hydroxypiperidin-1-yl)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,
- (22) N1-ethyl-5-((2-((pyrrolidin-1-ylcarbonyl)amino)-4-pyridyl)oxy)-1H-1-indolecarboxamide,
- (23) N4-(4-((1-(ethylanino)carbonyl-1H-5-indolyl)oxy)-2-pyridyl)-4-morpholinecarboxamide,
- (24) N1-cyclopropyl-5-(2-((pyrrolidin-1-

- ylcarbonyl) amino) -4-pyridyl) oxy-1H-1-indolecarboxamide,
 (25) N1-methyl-3-chloro-5-(2-((4-hydroxypiperidino) carbonyl) amino-4-pyridyl) oxy-1H-1-indolecarboxamide,
- 5 (26) N1-methyl-5-(2-((methylamino) carbonyl) amino-4-pyridyl) oxy-1H-1-indolecarboxamide,
- (27) N1-methyl-5-(2-((diethylamino) carbonyl) amino-4-pyridyl) oxy-1H-1-indolecarboxamide,
- (28) N1-(2-propynyl)-5-(2-((pyrrolidin-1-yl) carbonyl) amino-4-pyridyl) oxy-1H-1-indolecarboxamide,
- 10 (29) N1-methyl-5-(2-(azetidin-1-ylcarbonyl) amino-4-pyridyl) oxy-1H-1-indolecarboxamide,
- (30) N1-ethyl-5-(2-(azetidin-1-ylcarbonyl) amino-4-pyridyl) oxy-1H-1-indolecarboxamide,
- 15 (31) N1-cyclopropyl-5-(2-(azetidin-1-ylcarbonyl) amino-4-pyridyl) oxy-1H-1-indolecarboxamide,
- (32) N1-methyl-5-(2-(((4-(morpholin-4-yl)piperidin-1-yl) carbonyl) amino) pyridin-4-yloxy)-1H-1-indolecarboxamide,
- 20 (33) N1-methyl-5-(2-(((4-(azetidin-1-yl)piperidin-1-yl) carbonyl) amino) pyridin-4-yloxy)-1H-1-indolecarboxamide,
- (34) N1-methyl-5-(2-(((4-(diethylamino)piperidin-1-yl) carbonyl) amino) pyridin-4-yloxy)-1H-1-indolecarboxamide,
- 25 (35) N1-methyl-5-(2-(((4-(4-hydroxypiperidin-1-

yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide, and

(36) N1-propyl-5-(2-(pyrrolidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide;

5 <33> a compound according to <1>, a salt of the compound, or a hydrate of the foregoing, wherein the compound is a compound selected from a group consisting of

(1) 5-(2-(((4-hydroxy-4-methylpiperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methyllamide,

(2) N1-methyl-5-(2-((4-hydroxypiperidino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide,

15 (3) N1-methyl-5-(2-(((4-(piperidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide,

(4) N1-methyl-5-(2-(((4-(piperidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-

20 indolecarboxamide, and

(5) N4-(4-(1-(methyllamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-4-morpholinecarboxamide;

<34> a pharmaceutical composition comprising a compound according to any of <1> to <33> and a pharmaceutical
25 adjuvant;

<35> a prophylactic or therapeutic agent for a disease

for which angiogenesis inhibition is effective,
comprising as an active ingredient, a compound
according to any of <1> to <33>, a salt thereof, or a
hydrate of the foregoing;

5 <36> an angiogenesis inhibitor comprising as an active
ingredient, a compound according to any of <1> to <33>,
a salt thereof, or a hydrate of the foregoing;

 <37> an antitumor agent comprising as an active
ingredient, a compound according to any of <1> to <33>,
10 a salt thereof, or a hydrate of the foregoing;

 <38> an antitumor agent according to <37>, wherein the
tumor is a pancreatic cancer, a gastric cancer, a colon
cancer, a breast cancer, a prostate cancer, a lung
cancer, a renal cancer, a brain tumor, a blood cancer,
15 or an ovarian cancer;

 <39> a therapeutic agent for hemangioma comprising as
an active ingredient, a compound according to any of
<1> to <33>, a salt thereof, or a hydrate of the
foregoing;

20 <40> a cancer metastasis inhibitor comprising as an
active ingredient, a compound according to any of <1>
to <33>, a salt thereof, or a hydrate of the foregoing;

 <41> a therapeutic agent for retinal neovascularization
or diabetic retinopathy comprising as an active
25 ingredient, a compound according to any of <1> to <33>,
a salt thereof, or a hydrate of the foregoing;

<42> a therapeutic agent for an inflammatory disease comprising as an active ingredient, a compound according to any of <1> to <33>, a salt thereof, or a hydrate of the foregoing;

5 <43> a therapeutic agent for an inflammatory disease according to <42>, wherein the inflammatory disease is deformatant arthritis, rheumatoid arthritis, psoriasis or delayed hypersensitivity reaction;

10 <44> a therapeutic agent for atherosclerosis comprising as an active ingredient, a compound according to any of <1> to <33>, a salt thereof, or a hydrate of the foregoing;

15 <45> an angiogenesis inhibition-based antitumor agent comprising as an active ingredient, a compound according to any of <1> to <33>, a salt thereof, or a hydrate of the foregoing;

20 <46> a prophylactic or therapeutic method for a disease for which angiogenesis inhibition is effective, comprising administering to a patient, a pharmacologically effective dose of a compound according to any of <1> to <33>, a salt thereof, or a hydrate of the foregoing; and

25 <47> use of a compound according to any of <1> to <33>, a salt thereof, or a hydrate of the foregoing for the manufacture of a prophylactic or therapeutic agent for a disease for which angiogenesis inhibition is

effective.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

5 [0012] The meanings of the terms, symbols or the like used in the specification are described and the present invention is described in detail below.

10 [0013] It should be noted that, although the structural formula of a compound may indicate a certain isomer for convenience's sake in this specification, the present invention include all geometrical isomers generated in the structures of compounds, isomers such as optical isomers based on asymmetric carbon atom, stereoisomers and tautomers, and a mixture of isomers, which are not
15 limited to the descriptions of formulas for convenience's sake, either of isomers or mixtures may be included. Therefore, although optically active compounds and racemic compounds may be existent when they have asymmetric carbon atoms in a molecule, they
20 are not particularly limited in the present invention and any cases are included. In addition, although a variety of crystal morphism are existent, these are not limited similarly. Specifically, any of a single crystal form or mixtures may be included, in addition,
25 anhydrides, hydrates or solvates may be included.

[0014] In addition, compounds according to the present invention also include compounds which still indicate a desired activity after they are subjected to metabolism such as oxidation, reduction, hydrolysis and conjugation in an organism, and the present invention also includes compounds which produce the compounds according to the present invention after they are subjected to metabolism such as oxidation, reduction and hydrolysis.

[0015] The term "C₁₋₆ alkyl group" as described in the specification represents a linear or branched alkyl group of 1 to 6 carbon atoms, which is a monovalent group derived by removing a hydrogen atom from an aliphatic hydrocarbon of 1 to 6 carbon atoms. As specific examples there may be mentioned methyl group, ethyl group, n-propyl group, i-propyl group, n-butyl group, i-butyl group, sec-butyl group, t-butyl group, n-pentyl group, i-pentyl group, sec-pentyl group, neopentyl group, 1-methylbutyl group, 2-methylbutyl group, 1,1-dimethylpropyl group, 1,2-dimethylpropyl group, n-hexyl group, i-hexyl group, 1-methylpentyl group, 2-methylpentyl group, 3-methylpentyl group, 1,1-dimethylbutyl group, 1,2-dimethylbutyl group, 2,2-dimethylbutyl group, 1,3-dimethylbutyl group, 2,3-dimethylbutyl group, 3,3-dimethylbutyl group, 1-

ethylbutyl group, 2-ethylbutyl group, 1,1,2-trimethylpropyl group, 1,2,2-trimethylpropyl group, 1-ethyl-1-methylpropyl group, 1-ethyl-2-methylpropyl group or the like, and preferably methyl group, ethyl group, n-propyl group, i-propyl group, n-butyl group, i-butyl group, sec-butyl group and t-butyl group.

[0016] The term "C₂₋₆ alkenyl group" as described in the specification represents a linear or branched alkenyl group of 2 to 6 carbon atoms which may contain 1 to 2 double bonds. As specific examples there may be mentioned ethenyl group, 1-propenyl group, 2-propenyl group, 1-butenyl group, 2-butenyl group, 3-butenyl group, 2-methyl-1-propenyl group, pentenyl group, hexenyl group, hexadienyl group or the like, and preferably ethenyl group, 1-propenyl group, 2-propenyl group, 1-butenyl group, 2-butenyl group, 3-butenyl group and 2-methyl-1-propenyl group.

[0017] The term "C₃₋₆ alkenyl group" as described in the specification represents a linear or branched alkenyl of 3 to 6 carbon atoms which may contain 1 to 2 double bonds. As specific examples there may be mentioned 1-propenyl group, 2-propenyl group, 1-butenyl group, 2-butenyl group, 3-butenyl group, 2-methyl-1-propenyl group, pentenyl group, hexenyl group, hexadienyl group

or the like, and preferably 1-propenyl group, 2-propenyl group, 1-butenyl group, 2-butenyl group, 3-butenyl group and 2-methyl-1-propenyl group.

5 [0018] The term "C₂₋₆ alkynyl group" as described in the specification represents a linear or branched alkynyl group of 2 to 6 carbon atoms which may contain 1 to 2 triple bonds. As specific examples there may be mentioned ethynyl group, 1-propynyl group, 2-propynyl
10 group, 1-butyne group, 2-butyne group, 3-butyne group, pentynyl group, hexynyl group, hexadiynyl group or the like, and preferably ethynyl group, 1-propynyl group, 2-propynyl group, 1-butyne group, 2-butyne group and 3-butyne group.

15 [0019] The term "C₃₋₆ alkynyl group" as described in the specification represents a linear or branched alkynyl group of 3 to 6 carbon atoms which may contain 1 to 2 triple bonds. As specific examples there may be
20 mentioned 1-propynyl group, 2-propynyl group, 1-butyne group, 2-butyne group, 3-butyne group, pentynyl group, hexynyl group, hexadiynyl group or the like, and preferably 1-propynyl group, 2-propynyl group, 1-butyne group, 2-butyne group and 3-butyne group.

25 [0020] The term "C₃₋₈ cycloalkyl group" as described in

the specification represents a cyclic aliphatic hydrocarbon group of 3 to 8 carbon atoms, and as specific examples there may be mentioned cyclopropyl group, cyclobutyl group, cyclopentyl group, cyclohexyl group, cycloheptyl group, cyclooctyl group or the like, and preferably cyclopropyl group, cyclobutyl group and cyclopentyl group.

[0021] The term "C₁₋₆ alkylene group" as described in the specification represents a divalent group derived by further removing a hydrogen atom from the aforementioned definition of "C₁₋₆ alkyl group." As specific examples there may be mentioned methylene group, ethylene group, methylethylene group, propylene group, ethylethylene group, 1,1-dimethylethylene group, 1,2-dimethylethylene group, tetramethylene group, pentamethylene group, hexamethylene group or the like, and preferably methylene group and ethylene group.

[0022] The term "C₁₋₆ alkoxy group" as described in the specification represents an oxy group bonded with the aforementioned definition of "C₁₋₆ alkyl group." As specific examples there may be mentioned methoxy group, ethoxy group, n-propoxy group, i-propoxy group, n-butoxy group, i-butoxy group, sec-butoxy group, t-butoxy group, n-pentyloxy group, i-pentyloxy group,

sec-pentyloxy group, neopentyloxy group, 1-methylbutoxy group, 2-methylbutoxy group, 1,1-dimethylpropoxy group, 1,2-dimethylpropoxy group, n-hexyloxy group, i-hexyloxy group, 1-methylpentyloxy group, 2-methylpentyloxy group, 3-methylpentyloxy group, 1,1-dimethylbutoxy group, 1,2-dimethylbutoxy group, 2,2-dimethylbutoxy group, 1,3-dimethylbutoxy group, 2,3-dimethylbutoxy group, 3,3-dimethylbutoxy group, 1-ethylbutoxy group, 2-ethylbutoxy group, 1,1,2-trimethylpropoxy group, 1,2,2-trimethylpropoxy group, 1-ethyl-1-methylpropoxy group, 1-ethyl-2-methylpropoxy group or the like, and preferably methoxy group, ethoxy group, n-propoxy group, i-propoxy group, n-butoxy group, i-butoxy group, sec-butoxy group, and t-butoxy group.

[0023] The term "C₁₋₆ alkylthio group" as described in the specification represents a thio group bonded with the aforementioned definition of "C₁₋₆ alkyl group." As specific examples there may be mentioned methylthio group, ethylthio group, n-propylthio group, i-propylthio group, n-butylthio group, i-butylthio group, sec-butylthio group, t-butylthio group, n-pentylthio group, i-pentylthio group, sec-pentylthio group, neopentylthio group, 1-methylbutylthio group, 2-methylbutylthio group, 1,1-dimethylpropylthio group, 1,2-dimethylpropylthio group, n-hexylthio group, i-

hexylthio group, 1-methylpentylthio group, 2-methylpentylthio group, 3-methylpentylthio group, 1,1-dimethylbutylthio group, 1,2-dimethylbutylthio group, 2,2-dimethylbutylthio group, 1,3-dimethylbutylthio group, 2,3-dimethylbutylthio group, 3,3-dimethylbutylthio group, 1-ethylbutylthio group, 2-ethylbutylthio group, 1,1,2-trimethylpropylthio group, 1,2,2-trimethylpropylthio group, 1-ethyl-1-methylpropylthio group, 1-ethyl-2-methylpropylthio group or the like, and preferably methylthio group, ethylthio group, n-propylthio group, i-propylthio group, n-butylthio group, i-butylthio group, sec-butylthio group and t-butylthio group.

[0024] The term "C₆₋₁₀ aryl group" as described in the specification represents an aromatic hydrocarbon ring group of 6 to 10 carbon atoms. As specific examples there may be mentioned phenyl group, 1-naphthyl group, 2-naphthyl group, indenyl group, azulenyl group, heptalenyl group or the like, and preferably phenyl group, 1-naphthyl group and 2-naphthyl group.

[0025] The term "C₆₋₁₀ aryloxy group" as described in the specification represents an oxy group bonded with the aforementioned definition of "C₆₋₁₀ aryl group." As specific examples there may be mentioned phenoxy group,

1-naphthyloxy group, 2-naphthyloxy group, indenyl group, azulenyloxy group, heptalenyloxy group or the like, and preferably phenoxy group, 1-naphthyloxy group and 2-naphthyloxy group.

5

[0026] The term "halogen atom" as described in the specification represents fluorine atom, chlorine atom, bromine atom or iodine atom, and preferably fluorine atom, chlorine atom and bromine atom.

10

[0027] The term "heteroatom" as described in the specification represents nitrogen atom, sulfur atom, or oxygen atom.

15

[0028] The term "5- to 10- membered aromatic heterocycle" as described in the specification represents an aromatic ring in which the number of atoms forming the ring is 5 to 10, and 1 to a plurality of heteroatoms are contained in the atoms forming the ring. Specific examples are pyrrole ring, pyridine ring, pyridazine ring, pyrimidine ring, pyrazine ring, pyrazole ring, imidazole ring, triazole ring, tetrazole ring, indole ring, isoindole ring, indazole ring, quinoline ring, isoquinoline ring, cinnoline ring, quinazoline ring, quinoxaline ring, naphthyridine ring, phthalazine ring, carbazole ring, purine ring, furan

20

25

ring, thiophene ring, benzimidazole ring,
imidazopyridine ring, imidazotriazine ring,
pyrrolopyridine ring, pyrrolopyrimidine ring,
pyridopyrimidine ring, oxazole ring, isoxazole ring,
5 thiazole ring, isothiazole ring, phenoxazine ring,
phenothiazine ring, furopyrrole ring, imidazothiazole
ring, benzoxazole ring, benzthiazole ring,
pyrazoloxazole ring, pyridoxazine ring, benzofuran ring,
benzothiophene ring or the like, and preferably furan
10 ring, thiophen ring, and thiazole ring.

[0029] The term "5- to 10- membered heteroaryl group"
as described in the specification represents a
monovalent group derived by removing a hydrogen atom
15 from the aforementioned definition of "5- to 10-
membered aromatic heterocycle."

[0030] The term "3- to 10- membered heterocycle" as
described in the specification represents,
20 (1) a monocyclic or bicyclic non-aromatic ring
(2) having 3 to 10 atoms in the ring,
(3) containing 1 to 2 hetero atoms among the atoms of
the ring,
(4) optionally including 1 to 2 double bonds in the
25 ring, and
(5) optionally including 1 to 3 carbonyl groups or 1 to

3 sulfonyl groups in the ring.

Specific examples are aziridine ring, azetidine ring, pyrrolidine ring, piperidine ring, 4-oxopiperidine ring, homopiperidine ring, piperazine ring, homopiperazine ring, morpholine ring, thiomorpholine ring, 1,1-dioxothiomorpholine ring, pyridone ring, phthalimide ring, succinimide ring or the like, and preferably azetidine ring, pyrrolidine ring, piperidine ring, piperazine ring, morpholine ring and thiomorpholine ring.

[0031] The term "3- to 10- membered heterocyclic group" as described in the specification represents a monovalent group derived by removing a hydrogen atom from the aforementioned definition of "3- to 10-membered heterocycle."

[0032] The term "optionally substituted" as described in the specification is equivalent in the meaning as in "which may have 1 or a plurality of substituents by arbitrarily combining them at substitutable positions". As specific examples of such substituents there may be mentioned the following:

- (1) a halogen atom,
- (2) a hydroxyl group,
- (3) a thiol group,

- (4) a nitro group,
(5) a cyano group,
(6) an azido group,
(7) a formyl group,
5 (8) a carboxyl group,
(9) an amino group,
(10) an oxo group or
(11) a group represented by the formula $-T^1-T^2-T^3$,
wherein T^1 represents a single bond or a C_{1-6} alkylene
10 group; T^2 represents a single bond, a C_{1-6} alkylene
group, an oxygen atom, an sulfur atom, a sulfinyl group,
a sulfonyl group, a carbonyl group, or a group
represented by the formula $-O-CO-$, the formula $-CO-O-$,
the formula $-NR^{T1}-$, the formula $-CO-NR^{T1}-$, the formula $-$
15 $NR^{T1}-CO-$, the formula $-SO_2-NR^{T1}-$, or the formula $-NR^{T1}-$
 SO_2- ; T^3 represents a hydrogen atom, a C_{1-6} alkyl group,
a C_{2-6} alkenyl group, a C_{2-6} alkynyl group, a C_{3-8}
cycloalkyl group, a C_{6-10} aryl group, a 5- to 10-
membered heteroaryl group, a 3- to 10- membered
20 heterocyclic group or a group represented by the
formula $-N(R^{T2})(R^{T3})$; R^{T1} , R^{T2} , or R^{T3} each independently
represent a hydrogen atom or a C_{1-6} alkyl group; wherein
a C_{1-6} alkyl group, a C_{2-6} alkenyl group, a C_{2-6} alkynyl
group, a C_{3-8} cycloalkyl group, a C_{6-10} aryl group, a 5-
25 to 10- membered heteroaryl group and a 3- to 10-
membered heterocyclic group in T^3 may each

independently have 1 to 3 groups selected from a group
of the below-mentioned Substituent Group;

<Substituent Group>

a halogen atom, a hydroxyl group, a thiol group, a
5 nitro group, a cyano group, a C₁₋₆ alkyl group, a C₃₋₈
cycloalkyl group, a C₂₋₆ alkenyl group, a C₂₋₆ alkynyl
group, a C₆₋₁₀ aryl group, a 5- to 10- membered
heteroaryl group, a 3- to 10- membered heterocyclic
group, a C₁₋₆ alkoxy group and a C₁₋₆ alkylthio group.

10

[0033] The term "leaving group" as described in the
specification may be any group commonly known as a
leaving group in organic synthesis, with no special
restrictions, and as specific examples there may be
15 mentioned a halogen atom such as a chlorine atom, a
bromine atom, an iodine atom; a nitro group; an
alkylthio group such as a methylthio group, an
ethylthio group and a propylthio group; an arylthio
group such as a phenylthio group, a tolylthio group
20 and a 2-pyridylthio group; an alkylsulfonyloxy group
such as a methanesulfonyloxy group, a
trifluoromethanesulfonyloxy group, an ethanesulfonyloxy
group, a propanesulfonyloxy; an arylsulfonyloxy group
such as a benzenesulfonyloxy group, a p-
25 toluenesulfonyloxy group; an alkanoyloxy group such as
an acetoxy group and a trifluoroacetoxy group; an

alkoxy group such as a methoxy group, an ethoxy group
and a propoxy group; an alkylamino group such as a
methylamino group, an ethylamino group, a propylamino
group and a butylamino group; a dialkylamino group such
5 as a dimethylamino group, a diethylamino group, a
dipropylamino group, a methylethylamino group, an
ethylpropylamino group and a methylpropylamino group; a
substituted phosphoryloxy group such as
diphenoxyphosphoryloxy group or the like, and
10 preferably a halogen atom such as a chlorine atom, a
bromine atom and an iodine atom, a
trifluoromethanesulfonyl group or the like.

[0034] As a "salt" described in the specification,
15 there may be mentioned, for example, a salt with
inorganic acid, a salt with organic acid, a salt with
inorganic base, a salt with organic base, a salt with
acidic or basic amino acid or the like, preferably a
pharmacologically acceptable salt. A salt is formed in
20 an appropriate ratio of 0.1 to 5 molecules of acid or
base to one molecule of the compound.

[0035] As preferable examples of a salt with inorganic
acid, there may be mentioned, for example, a salt with
25 hydrochloric acid, hydrobromic acid, sulfuric acid,
nitric acid, phosphoric acid, or the like, and as

preferable examples of a salt with organic acid, there may be mentioned, for example, a salt with acetic acid, succinic acid, fumaric acid, maleic acid, tartaric acid, citric acid, lactic acid, stearic acid, benzoic acid, methanesulfonic acid, p-toluenesulfonic acid or the like.

[0036] As preferable examples of a salt with inorganic base, there may be mentioned, for example, an alkali metal salt such as a sodium salt and a potassium salt, an alkali earth metal salt such as a calcium salt and a magnesium salt, an aluminum salt, an ammonium salt or the like. As preferable examples of a salt with organic base, there may be mentioned, for example, a salt with diethylamine, diethanolamine, meglumine, N,N'-dibenzylethylenediamine or the like.

[0037] As preferable examples of a salt with acidic amino acid, there may be mentioned, for example, a salt with aspartic acid, glutamic acid or the like, and as preferable examples of a salt with basic amino acid, there may be mentioned, for example, a salt with arginine, lysine, ornithine or the like.

[0038] As a "adjuvant" described in the specification, there may be mentioned, for example, a excipient, a

binder, a disintegrator, a lubricant, a coloring agent, a corrective coating, a stabilizer, a emulsifier, a absorbefacient, a surfactant, a pH adjustor, a preservative, an antioxidant or the like.


5

[0039] Production methods for the compounds of the invention will now be described. Various methods may be considered for production of compounds of the invention represented by the general formulas (I) and
 10 (II) with synthesis carried out by ordinary organic synthesis means, and the following are representative examples of methods for their production.

[0040] [General synthesis method]

15 [Production method 1]

A typical production method of the compound represented by the formula (Ia)



wherein, R_{101} , R_{102} may represent the same definitions as
 20 the formula R_{12a} , R_{12b} (R_{12a} and R_{12b} represent the same definitions as the aforementioned definition), respectively; or R_{101} and R_{102} form a ring, and the formula $-NR_{101}R_{102}$ may represent the same definition as


the formula



(wherein T1 represents the same definition as the
aforementioned definition); other symbols represent the
same definitions as the aforementioned definitions.

[0041] [Production method 1-A]

A typical production method of the compound (1e), which
is the compounds represented by the formula (1a),
wherein Y represents an oxygen atom, a sulfur atom or a
group represented by the formula $-NR_Y-$ (R_Y represents a
hydrogen atom or a C_{1-6} alkyl group)



wherein, Y_1 represents an oxygen atom, a sulfur atom or
a group represented by the formula $-NR_{Y1}-$ (R_{Y1}
represents a hydrogen atom or a C_{1-6} alkyl group); L_1
represents a leaving group; other symbols represent the
same definitions as the aforementioned definition.

<Step 1A-1>

[0042] This is a step for obtaining a compound (1c) by condensing pyrimidine or a pyridine derivative (1a) having a leaving group (L_1) at the 4-position with an indole derivative (1b). As a reaction solvent, N-methylpyrrolidone, N,N-dimethylformamide, dimethyl sulfoxide, 2-ethoxyethanol, chlorobenzene or the like can be used. A base or an acid may be added thereto, specifically, an organic base such as diisopropylethylamine, an inorganic base such as potassium carbonate, cesium carbonate and sodium hydride and an acid such as pyridine hydrochloride and hydrochloric acid can be used. The reaction can be performed at a temperature ranging from room temperature to reflux temperature for a reaction time ranging from 10 minutes to 30 hours. In addition, a compound where a halogen atom which is not as a leaving group is bonded on pyrimidine or pyridine ring may be used as a starting material, and the halogen atom can be reduced by the catalytic reduction method or the like after this step.

<Step 1A-2>

[0043] This is a step for obtaining a compound (1d) by carboxamidating the 1-position of indole in compound

(1c). As a reagent, a carbamate derivative, an isocyanate derivative, a halogenated carbamoyl derivative or the like can be used. As a reaction solvent, chloroform, toluene, N-methylpyrrolidone, N,N-dimethylformamide, dimethyl sulfoxide, chlorobenzene can be used. A base may be added thereto, specifically, an organic base such as pyridine, triethylamine and diisopropylethylamine, an inorganic base such as potassium carbonate, cesium carbonate and sodium hydride can be used, for example. The reaction can be performed for a time of 10 minutes to 30 hours at a temperature of 0 °C to reflux temperature.

<Step 1A-3>

[0044] This is a step for converting a compound (1d) into a urea derivative (1e). Carbamate ester derivative is prepared by using phenyl chlorocarbonate or the like as a reagent, for example. After this intermediate is isolated, or not isolated, the intermediate is allowed to react with an amine, thereby a urea derivative can be obtained. Alternatively, by reacting a carbamate derivative or an isocyanate derivative as a reagent, a corresponding urea derivative can be converted into. As a reaction solvent, chloroform, toluene, N-methylpyrrolidone, N,N-dimethylformamide, dimethyl sulfoxide, chlorobenzene or


the like can be used. A base may be added thereto, specifically, an organic base such as pyridine, triethylamine, and diisopropylethylamine, an inorganic base such as potassium carbonate, cesium carbonate and sodium hydride can be used, for example. The reaction can be performed for a time of 10 minutes to 30 hours at a temperature of 0 °C to reflux temperature.

[0045] It should be noted that a substituent conversion in R_2 , R_{101} , R_{102} can be also performed by suitably using an oxidation reaction, a reduction reaction, a reductive amination reaction, an ester formation reaction, an amide formation reaction, a protecting group introduction reaction, a deprotection reaction, a hydrolysis reaction or the like which are generally used before and/or after each process. Specifically, for example, in the case that R_2 is a hydrogen atom in the compounds (1a), (1c) and (1d), the following methods come under the above-mentioned substituent conversions; that is, a method for converting R_2 into a C_{1-6} alkyl group by performing a reductive amination reaction with aldehyde or ketone, a method in which, after a corresponding urea derivative is obtained as in <Step 1A-3> from the compound (1c) and an amine having ketone or aldehyde, an amine side chain is introduced into R_{101} , R_{102} by further performing a reductive

amination reaction with an amine, or the like. In these cases, sodium cyanoborohydride, sodium trimethoxyborohydride, sodium triacetoxyborohydride or the like can be used as a reducing agent, and methanol, tetrahydrofuran, dichloromethane, dichloroethane or the like can be used as a reaction solvent. In addition, a method that a benzotriazole derivative is prepared and the derivative is reduced by sodium borohydride as reported in Tetrahedron 47, 2683 (1991), or the like is useful. Alternatively, a corresponding urea is formed as in <Step 1A-3> from the compound (1c) and an amine having an ester. After the ester is hydrolyzed by bases such as lithium hydroxide, sodium hydroxide or potassium hydroxide in aqueous ethanol, an amide derivative can be also obtained by using a condensing agent. In this case, N,N-dimethylformamide, tetrahydrofuran or the like can be used as a reaction solvent, and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride, (1H-1,2,3-benzotriazole-1-yloxy) (tri(dimethylamino))phosphonium hexafluorophosphate can be used as a condensing agent. The reaction can be performed for a time of 10 minutes to 30 hours at a temperature of 0 °C to reflux temperature.

[Production method 1-B]

[0046] A production method of the compound (1g), which is the compound represented by the formula (Ia), wherein Y is a sulfinyl group or a sulfonyl group




5 wherein, Y₂ represents a sulfinyl group or a sulfonyl group; other symbols represent the same definitions as aforementioned definitions.

<Step 1B-1>

10 [0047] This is a step for oxidation of a compound (1f) to a compound (1g). Hydrogen peroxide, peracetic acid, methaperiodate, 3-chloroperbenzoic acid or the like can be used as an oxidizing agent. Methanol, water, dichloromethane, chloroform or the like can be used as
15 a solvent. The reaction can be performed for a time of 10 minutes to 30 hours at a temperature of 0 °C to reflux temperature.

[Production method 2]

20 Another production method of the compound (1c), which is the compound (1c) having a halogen atom, a formyl group, or a cyano group as a substituent at the 3-position in the indole ring



wherein, R_{201} represents a halogen atom, a formyl group or a cyano group; other symbols represent the same definitions as the aforementioned definitions.

5

<Step 2-1>

[0048] This is a step for obtaining a compound (2b) by the condensation of a pyrimidine or pyridine derivative (1a) and a indole derivative (2a) not having a substituent on the 3-position. The compound (2b) can be obtained under the same conditions as <Step 1A-1>.

10

<Step 2-2>

[0049] This is a step in which a substituent is introduced into 3-position of indole in a compound (2b) to obtain a compound substituted at the 3-position of indole (2c). A compound (2c) substituted with a halogen atom, a formyl group, an amino group or the like as the 3-position substituent can be obtained by reacting a compound (2b) with halogenation agents such as N-chlorosuccinimide, N-bromosuccinimide or a mixed reagent of phosphorous oxychloride or thionyl chloride


15

20

with N,N-dimethylformamide, or after converting the compound into a N-chlorosulfonylcarbamoyl derivative by allowing chlorosulfonyl isocyanate to react with the compound, followed by allowing triethylamine to react with the derivative or the like as reported in Tetrahedron 50, 6549 (1994). As a reaction solvent, 2-propanol, N,N-dimethylformamide, tetrahydrofuran, acetonitrile or the like can be used, and the reaction can be performed for a time of 10 minutes to 30 hours at a temperature of 0 °C to reflux temperature.

[Production method 3]

[0050] Another production method of the compound (1d) via the compounds (3c), (3d), (3g) or (3h)



wherein, P represents a protecting group; other symbols represent the same definitions as in the aforementioned definitions.

5

<Step 3-1><Step 3-2><Step 3-3>

[0051] These are steps for obtaining an indole derivative (3c) or an indoline derivative (3d), both being introduced a carboxamide group at the 1-position, via a compound (3b) from an indole derivative (3a).

10

[0052] <Step 3-1> is a step for conducting carboxamidation of the 1-position of an indole derivative (3a) to obtain a compound (3b), and can be performed in a similar way as <Step 1A-2>. A methyl group, a benzyl group, a substituted benzyl group, a benzyloxycarbonyl group can be used as a protecting group, for example.

[0053] <Step 3-2> is a step for obtaining a compound (3c) by deprotecting an indole derivative (3b). Specifically, for example, in the case that Y₁ is an oxygen atom, the methods used for ordinary deprotection such as demethylation by using boron tribromide, debenzylation by using trifluoroacetic acid-thioanisole, debenzylation or the debenzyloxycarbonylation by catalytic reduction can be used.

[0054] <Step 3-3> is a step for reduction of an indole derivative (3c) to an indoline derivative (3d). Catalytic hydrogenation reaction in the presence of palladium catalyst under ordinary pressure or under pressurization or the like can be applied. Methanol, N,N-dimethylformamide, tetrahydrofuran or the like can be used as a reaction solvent, and the reaction can be performed for a time of 10 minutes to 30 hours at a

temperature of 0 °C to reflux temperature.

<Step 3-4><Step 3-5><Step 3-6>

5 [0055] These are steps for obtaining an aminoindole derivative (3g) or an aminoindoline derivative (3h) having a carboxamide group at the 1-position via a compound (3f) from a nitroindole derivative (3e).

10 [0056] <Step 3-4> is a step conducting carboxamidation of the 1-position of a indole derivative (3e) to obtain a compound (3f), and can be performed in the same way as in <Step 1A-2>.

15 [0057] <Step 3-5> is a step for reducing a nitroindole derivative (3f) to an aminoindole derivative (3g). The conditions used for reduction reaction of a nitro group to an amino group generally utilized, specifically, for example, reduction by iron-ammonium chloride or iron-acetic acid or the like, catalytic reduction by
20 palladium hydroxide-hydrogen or the like can be applied. Methanol, ethanol, water, N,N-dimethylformamide, tetrahydrofuran or the like can be used as a reaction solvent, and the reaction can be performed at a temperature of room temperature to reflux temperature
25 for 10 minutes to 30 hours.

[0058] <Step 3-6> is a step for reducing an indole derivative (3g) to an indoline derivative (3h) and can be performed in the same way as in <Step 3-3>.

5 <Step 3-7><Step 3-8>

[0059] These are steps for condensing an indole derivative (3c or 3g) or an indoline derivative (3d or 3h) and a compound (1a) to obtain an indole derivative (1d) or an indoline derivative (3i), and can be
10 performed in the same way as in <Step 1A-1>.

<Step 3-9>


[0060] This is a step for oxidizing an indoline derivative (3i) to an indole derivative (1d). For
15 example, 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) or the like can be used as an oxidizing agent, and 1,4-dioxane, toluene, benzene or the like can be used as a solvent. Alternatively, a method in which manganese acetate (III) is used as an oxidizing agent
20 or the like as reported in Tetrahedron Lett. 29, 2151 (1988) can be applied.

In addition, in the case that Y_1 is the formula $-NR_{Y1}-$ and R_{Y1} is a hydrogen atom in compounds (3g), (3h), (3c) or (3d), a compound (1d), wherein Y_1 is the
25 formula $-NR_{Y1}-$ and R_{Y1} is a C_{1-6} alkyl group, can be also obtained by converting the hydrogen atom into a C_{1-6}

alkyl group by a reductive amination reaction with aldehyde or ketone, and by using these for the respective following reactions. In addition, in the case that Y_1 is the formula $-NR_{Y1}-$ and R_{Y1} is a hydrogen atom in compounds (3i) or (1d), the compounds can be also similarly converted into the compounds (3i) or (1d), wherein Y_1 is the formula $-NR_{Y1}-$ and R_{Y1} is a C_{1-6} alkyl group. In this case, sodium cyanoborohydride, sodium trimethoxyborohydride, sodium triacetoxymethylborohydride or the like can be used as a reducing agent, and methanol, tetrahydrofuran, dichloromethane, dichloroethane or the like can be used as a reaction solvent. In addition, a method in which a benzotriazole derivative is prepared and is reduced by sodium borohydride as reported in Tetrahedron 47,2683 (1991) or the like can be applied.

[Production method 4]

[0061] Another production method of the compound (1e)



wherein each symbol represents the same definition as the aforementioned definition.

<Step 4-1>

[0062] This is a step for converting a compound (1c) to a compound (4a), and can be performed in the same way as in <Step 1A-3>.

5

<Step 4-2>

[0063] This is a step for conducting carboxamidation of the 1-position of indole in a compound (4a) to obtain a compound (1e), and can be performed in the same way as in <Step 1A-2>.


10

It is to be noted that, as described in [Production method 1-A], a substituent conversion can be also performed in R_2 , R_9 , R_{101} and R_{102} by properly performing oxidation reaction, reduction reaction, reductive amination reaction, ester formation reaction, amide formation reaction, protecting group introduction reaction, deprotection reaction, hydrolysis reaction or the like generally utilized after these steps.

15

20 [Production method 5]

[0064] Another production method of a compound (1e)



wherein, each symbol represents the same definition as

the aforementioned definition.

<Step 5-1>

5 [0065] This is a step for converting a pyrimidine or pyridine derivative (1a) into a corresponding urea derivative (5a), and can be performed in the same way as in <Step 1A-3>.

<Step 5-2>


10 [0066] This is a step for obtaining a compound (1e) from a pyrimidine or pyridine derivative (5a) having urea. A method in which the same operations as in <Step 1A-1> and <Step 1A-2> are sequentially performed, a method in which the same operations as in <Step 2-1>, 15 <Step 2-2> and <Step 1A-2> are sequentially performed, a method as in <Step 3-7>, a method in which the same operations as in <Step 3-8> and <Step 3-9> are sequentially performed or the like can be applied.

20 [0067] It is to be noted that, as described in [Production method 1-A], a substituent conversion can be also performed in R₂, R₉, R₁₀₁ and R₁₀₂ by properly performing oxidation reaction, reduction reaction, reductive amination reaction, ester formation reaction, 25 amide formation reaction, protecting group introduction reaction, deprotection reaction, hydrolysis reaction or

the like generally utilized after these steps.

[Production method 6]

[0068] Another manufacturing method of compounds (1c),
(1d), (3i)



wherein, L_2 represents a leaving group; each symbol represents the same definition as the aforementioned definition.

<Step 6-1><Step 6-2><Step 6-3>

[0069] These are steps for condensing a pyrimidine or pyridine derivative having leaving groups L_1 and L_2 and an indole or indoline derivative. In these steps, it is preferable that L_1 is a substituent having higher

reactivity than that of L_2 . Specifically, for example, a combination of L_1 being a nitro group and L_2 being a chlorine atom or the like comes under the category. By using an indole derivative (1b), indole derivatives (3c), (3g) having a carboxamide group at the 1-position, indoline derivatives (3d), (3h) having a carboxamide group at the 1-position, each compound (6b), (6c) and (6d) can be obtained under the same conditions as in <Step 1A-1>.

<Step 6-4>

[0070] This is a step for conducting carboxamidation of the 1-position of indole in a compound (6b) to obtain a compound (6c), and can be performed in the same way as in <Step 1A-2>.

<Step 6-5>

[0071] This is a step for oxidizing an indoline derivative (6d) to an indole derivative (6c). The same method as in <Step 3-9> can be used.

<Step 6-6><Step 6-7><Step 6-8>


[0072] These are steps in which the leaving group L_2 of pyrimidine or pyridine derivatives (6b), (6c), or (6d) is converted into a group represented by the formula - NHR_2 , wherein R_2 represents the same definition as the

aforementioned definition, to obtain compounds (1c),
 (1d), or (3i), respectively. For example, an ammonia-
 ethanol solution or a corresponding primary amine is
 used, and the reaction can be performed in a sealed
 5 tube for a time of 10 minutes to 100 hours at a
 temperature of 60 °C to reflux temperature.

[Production method 7]

[0073] Another production method of a compound (7j),
 10 which is the compound represented by the formula (1a),
 wherein Y is an oxygen atom and both 2- and 3-positions
 of indole (R_8 , R_7) are hydrogen atoms

[production method 7-A]



15 wherein each symbol represents the same definition as
 the aforementioned definition.

<Step 7A-1>

[0074] This is a step for obtaining a compound (7a) by introducing an aminophenoxy group into a compound (6a). It is preferable that in the compound (6a), L₁ is a substituent having higher reactivity than that of L₂. Specifically, for example, a combination of L₁ being a nitro group and L₂ being a chlorine atom comes under the category. A compound (7a) can be obtained by using a compound (6a) and an aminophenol derivative in the same method as in <Step 1A-1>. In addition, after these compounds are condensed by using a nitrophenol derivative in the same way as in <Step 1A-1>, a method for reducing a nitro group by catalytic hydrogenation reaction using palladium catalyst or the like, or metal reduction reaction using iron-ammonium chloride, iron-acetic acid or the like can be applied. In the reduction reaction of the nitro group, methanol, ethanol, tetrahydrofuran, N,N-dimethylformamide or the like can be used as a reaction solvent, and the catalytic hydrogenation reaction can be performed at ordinary pressure or under pressurization. The reaction can be performed at a temperature of room temperature to reflux temperature for 10 minutes to 30 hours.

<Step 7A-2>

[0075] This is a step for protecting amino group of a compound (7a) to obtain a compound (7b). As a protecting group, for example, a benzyloxycarbonyl group or the like can be introduced by using a corresponding chlorocarbonate ester.

<Step 7A-3>

[0076] This is a method for obtaining a compound (7c) from a compound (7b). t-Butyl bromoacetate as a reagent, sodium hydride or the like as a base, N,N-dimethylformamide, tetrahydrofuran, dimethyl sulfoxide or the like as a reaction solvent can be used. The reaction can be performed at a temperature of room temperature to reflux temperature for 10 minutes to 30 hours.

<Step 7A-4>

[0077] This is a step for deprotecting a compound (7c) to obtain a compound (7d). There may be mentioned, for example, deprotection reaction by the catalytic hydrogenation reaction of benzyloxycarbonyl group or the like.

<Step 7A-5>

[0078] This is a step for obtaining a compound (7e) by introducing a carboxamide group to a compound (7d). As

a reagent, an isocyanate derivative, a carbamate derivative or the like can be used. As a reaction solvent, N,N-dimethylformamide, tetrahydrofuran, dimethyl sulfoxide, toluene or the like can be used, and organic bases such as triethylamine or pyridine can be added thereto as requested. The reaction can be performed for a time of 10 minutes to 30 hours and at a temperature of 0 °C to reflux temperature.

10 <Step 7A-6>

[0079] This is a step for obtaining a compound (7f) from a compound (7e) by cyclization reaction. The reaction is performed in an acidic condition, specifically, for example, in trifluoroacetic acid-trifluoroacetic anhydride or the like. The reaction can be performed for a time of 10 minutes to 30 hours and at a temperature of 0 °C to reflux temperature.

<Step 7A-7><Step 7A-8>


20 [0080] These are steps for converting into an indole derivative (7h) via a compound (7g) from a 3-oxoindoline derivative (7f). A 3-hydroxyindoline derivative (7g) is prepared by reduction of a carbonyl group using sodium borohydride as a reagent, in tetrahydrofuran, methanol, ethanol or the like as a reaction solvent, thereafter a compound (7h) can be

obtained by performing dehydration by using camphor sulfonic acid or the like as a reagent, and toluene, dichloroethane or the like as a reaction solvent.

5 <Step 7A-9><Step 7A-10>

[0081] Thereafter, it is possible to lead to a step in which a compound (7j) is prepared under the same conditions in each of <Step 6-6>, <Step 1A-3>

[Production method 7-B]



10

wherein, each symbol represents the same definition as the aforementioned definition.

<Step 7B-1>

15

[0082] This is a step for obtaining a compound (7k) from a compound (5a), and can be performed in the same way as in <Step 7A-1>.

<Step 7B-2>

[0083] This is a step for protecting an amino group of a compound (7k) to obtain a compound (7l), and can be performed in the same way as in <Step 7A-2>.

<Step 7B-3>

[0084] This is a method for obtaining a compound (7m) from a compound (7l), and can be performed in the same way as in <Step 7A-3>.

<Step 7B-4>

[0085] This is a step for deprotecting a compound (7m) to obtain a compound (7n), and can be performed in the same way as in <Step 7A-4>.

<Step 7B-5>

[0086] This is a step for introducing a carboxamide group to a compound (7n) to obtain a compound (7o), and can be performed in the same way as in <Step 7A-5>.

<Step 7B-6>

[0087] This is a step for obtaining a cyclized compound (7p) from a compound (7o), and can be performed in the same way as in <Step 7A-6>.


<Step 7B-7><Step 7B-8>

[0088] These are steps for converting into an indole derivative (7j) via a compound (7q) from a 3-oxoindoline derivative (7p), and can be performed in the same as in <Step 7A-7><Step 7A-8>.

[Production method 8]

[0089] Another production method of a compound (8g), which is the compound represented by the formula (Ia), wherein both 2- and 3-positions of indole (R_8, R_7) are hydrogen atoms

[Production method 8-A]



wherein, each symbol represents the same definition as the aforementioned definition.

<Step 8A-1>

[0090] This is a coupling reaction of a compound (6a) with a nitrobenzene derivative. A compound (8a) can be obtained under the same conditions as in <Step 1A-1>.

5 <Step 8A-2>

[0091] This is a step for obtaining a compound (8b) from a compound (8a). The reaction can be performed under the conditions as described in Tetrahedron Lett. 39, 71 (1998). Specifically, a dimethylacetal compound
10 can be derived by condensing a nitrotoluene derivative and dimethylformamide dimethylacetal in N,N-dimethylformamide at a temperature of room temperature to reflux temperature for 10 minutes to 30 hours, and by sequentially performing the reaction of the compound
15 in methanol under acidic condition at a temperature of room temperature to reflux temperature for 10 minutes to 30 hours.

 <Step 8A-3>

20 [0092] This is a step for reducing a compound (8b) to a compound (8c). Reduction by iron-ammonium chloride, iron-acetic acid or the like can be used. As a reaction solvent, methanol, ethanol, tetrahydrofuran, N,N-dimethylformamide or the like can be used. The
25 reaction can be performed at a temperature of room temperature to reflux temperature for 10 minutes to 30

hours.

<Step 8A-4>

[0093] This is a step for converting a compound (8c)
5 into a urea derivative to obtain a compound (8d), and
can be performed in the same way as in <Step 7A-5>.
Alternatively, tetrahydrofuran or N,N-dimethylformamide
is used as a reaction solvent, for example, after a
carbamate derivative is prepared by using phenyl
10 chlorocarbonate or the like, and urea can be also
introduced by allowing the derivative to react with an
amine at a temperature of room temperature to reflux
temperature for 10 minutes to 30 hours, while N,N-
dimethylformamide, dimethyl sulfoxide are used as a
15 reaction solvent.

<Step 8A-5>

[0094] This is a step for cyclizing a compound (8d) to
obtain a compound (8e). The reaction can be performed
20 under the conditions as described in Tetrahedron Lett.
39, 71 (1998). Specifically, there may be mentioned a
method in which reflux is performed in solvents such as
benzene in the presence of catalytic amounts of camphor
sulfonic acid and quinoline.

25


<Step 8A-6>

[0095] This is a step for obtaining a compound (8f) from a compound (8e), and can be performed in the same way as in <Step 6-6>.

5 <Step 8A-7>

[0096] This is a step for obtaining a compound (8g) from a compound (8f), and can be performed in the same way as in <Step 7A-10>.

10 [0097] [Production method 8-B]



wherein, each symbol represents the same definition as in the aforementioned definition.

15 <Step 8B-1>

[0098] This is a step for obtaining a compound (8h) by performing coupling reaction of a compound (1a) with a nitrobenzene derivative, and can be performed in the same way as in <Step 1A-1>.

5

<Step 8B-2>

[0099] This is a step for introducing urea to a compound (8h) to obtain a compound (8i), and can be performed in the same way as in <Step 1A-3>.

10

<Step 8B-3>

[0100] This is a step for condensing a nitrotoluene derivative (8i) and dimethylformamide dimethylacetal, subsequently, for deriving the compound to dimethylacetal compound (8j). The step can be performed in the same way as in <Step 8A-2>.

15

<Step 8B-4>

[0101] This is a step for reducing a nitro group of a compound (8j) to obtain a compound (8k), and can be performed in the same way as in <Step 8A-3>.

20

<Step 8B-5>

[0102] This is a step for obtaining a compound (8l) from a compound (8k) by introducing urea, and can be performed in the same way as in <Step 8A-4>.

25


<Step 8B-6>

[0103] This is a step for cyclizing a compound (8l) to obtain a compound (8g), and can be performed in the same way as in <Step 8A-5>.

[Production method 9]

[0104] Another production method of a compound (7j)

[Production method 9-A]



wherein, each symbol represents the same definition as the aforementioned definition.

<Step 9A-1>

[0105] This is a step for obtaining a compound (9a) by coupling of a compound (6a) with a phenol derivative. Specifically, for example, a corresponding condensed compound can be obtained under the same conditions as in <Step 1A-1>, by using 4-amino-3-iodophenol obtained from t-butyl (2-iodo-4-

((triisopropylsilyl)oxy)phenyl)carbamate obtained by a method as described in J. Org. chem., 62, 6507 (1997) by allowing n-butylammonium fluoride or the like to react therewith.

5

<Step 9A-2>

[0106] This is a step for converting a compound (9a) into a urea derivative to obtain a compound (9b), and can be performed in the same way as in <Step 8A-4>.

10

<Step 9A-3>

[0107] This is a step for obtaining an acetylene derivative (9c) from an iodo compound (9b) using trimethylsilylacetylene. The condensation can be performed in the presence of tetrakis(triphenylphosphine)palladium or dichlorobis(triphenylphosphine)palladium, cuprous iodide. N,N-dimethylformamide or the like can be used as a reaction solvent, and the reaction can be performed at a temperature of room temperature to reflux temperature for 10 minutes to 30 hours.

15

20

<Step 9A-4>


[0108] This is a step for performing cyclization by heating an acetylene derivative (9c) in the presence of cuprous iodide to obtain an indole derivative (7h).

25

N,N-dimethylformamide or the like can be used as a reaction solvent, and the reaction can be performed at a temperature of 80 °C to reflux temperature for 5 minutes to 10 hours.

5 Subsequently, a compound (7h) can be converted into a compound (7j) as described in <Step 7A-9>, <Step 7A-10>.

[0109] [Production method 9-B]



10 wherein each symbol represents the same definition as in the aforementioned definition.

<Step 9B-1>

[0110] This is a step for coupling a compound (5a) with
 15 a phenol derivative to obtain a compound (9d), and can be performed in the same way as in <Step 9A-1>.

<Step 9B-2>

[0111] This is a step for converting a compound (9d)

into a urea derivative to obtain a compound (9e), and can be performed in the same way as in <Step 7A-5>.

<Step 9B-3>

5 [0112] This is a step for obtaining an acetylene derivative (9f) from an iodo compound (9e) by using trimethylsilylacetylene, and can be performed in the same way as in <Step 9A-3>.

10 <Step 9B-4>

[0113] This is a step for cyclizing an acetylene derivative (9f) by heating in the presence of cuprous iodide to obtain an indole derivative (7j). The same conditions as in <Step 9A-4> can be applied.


15

[Production method 10]

[0114] A typical production method of a compound (10g), which is the compound represented by the formula (Ia), wherein Y is an oxygen atom, a sulfur atom or a group represented by the formula -NR_Y- (wherein R_Y represents a hydrogen atom or a C₁₋₆ alkyl group), X₁ is a group represented by the formula -C(CN)=, X₂ is a group represented by the formula -CH=, R₂ is a hydrogen atom, R₃ is a hydrogen atom, an optionally substituted C₁₋₆ alkyl group or an optionally substituted C₃₋₈ cycloalkyl group

20

25



wherein, X_{201} represents a chlorine atom or a bromine atom, X_{202} represents a bromine atom or an iodine atom; R_{301} represents a hydrogen atom, an optionally substituted C_{1-6} alkyl group, or an optionally substituted C_{3-8} cycloalkyl group; it is preferable that as a combination of X_{201} and X_{202} , X_{202} is an iodine atom or a bromine atom if X_{201} is a chlorine atom, X_{202} is an iodine atom if X_{201} is a bromine atom; other symbols represent the same definition as the aforementioned definition.

<Step10-1>

[0115] This is a step for bromination or iodination of

the 5-position of a 2-aminopyridine derivative (10a) having a chlorine atom or a bromine atom at the 4-position to obtain a compound (10b). For example, halogenation agents such as iodine, N-bromosuccinimide, bromine, N-iodosuccinimide can be used. As a reaction solvent, for example, N,N-dimethylformamide, dimethyl sulfoxide, methylene chloride and acetonitrile can be used. The reaction can be performed at a temperature of 0 °C to reflux temperature for 10 minutes to 48 hours.

<Step 10-2>

[0116] This is a step for converting X_{202} of a compound (10b) into a cyano group to obtain a compound (10c).

For example, 0.5 to 0.6 equivalent of zinc cyanide, 1.0 to 1.2 equivalent of potassium cyanide, or trimethylsilylcyanide is reacted with a compound (10b) in the presence of a palladium catalyst such as tetrakis(triphenylphosphine)palladium or dichlorobis(triphenylphosphine)palladium. As a reaction solvent, for example, N,N-dimethylformamide, dioxane and tetrahydrofuran can be used. The reaction can be performed at a temperature of room temperature to reflux temperature for 10 minutes to 10 hours.

<Step 10-3><Step 10-4><Step 10-5>

[0117] These are steps for condensing a pyridine derivative (10c) and an indole or indoline derivative. Compounds (10d), (10e) and (10f) can be obtained, respectively, by using an indole derivative (1b),
5 indole derivatives (3c), (3g) having a carboxamide group at the 1-position, and indoline derivatives (3d), (3h) having a carboxamide group at the 1-position under the same conditions as in <Step 1A-1>.

10 <Step 10-6>

[0118] This is a step for conducting carboxamidation of 1-position of indole of a compound (10d) to obtain a compound (10e), and can be performed in the same way as in <Step 1A-2>.

15

<Step 10-7>

[0119] This is a step for oxidizing an indoline derivative (10f) to an indole derivative (10e), and can be performed in the same way as in <Step 3-9>.

20


<Step 10-8>

[0120] This is a step for converting a compound (10e) into a compound (10g), and can be performed in the same way as in <Step 1A-3>.

25

[Production method 11]

[0121] Another production method of a compound (10g)



wherein each symbol represents the same definition as the aforementioned definition.

5

<Step 11-1><Step 11-2>

[0122] These are steps for converting aminopyridine derivatives (10c), (10a) into corresponding urea derivatives (11e), (11c), respectively, and can be performed in the same way as in <Step 1A-3>.

10

<Step 11-3>

[0123] This is a step for iodination or bromination of the 5-position of a 2-ureidopyridine derivative (11c) having a chlorine atom or a bromine atom at the 4-position to obtain a compound (11d), and can be performed in the same way as in <Step 10-1>.

15

<Step 11-4>

[0124] This is a step for converting X₂₀₂ of a compound

20


(11d) into a cyano group to obtain a compound (11e), and can be performed in the same way as in <Step 10-2>.

<Step 11-5>

5 [0125] This is a step for obtaining a compound (10g) from a pyridine derivative (11e) having urea, and can be performed in the same way as in <Step 5-2>.

[Production method 12]

10 A production method of a compound (12b), which is the compound represented by the formula (Ia), wherein Y is a sulfinyl group or a sulfonyl group, X_1 is a group represented by the formula $-C(CN)=$, and R_2 is a hydrogen atom




15 wherein other symbols represent the same definitions as the aforementioned definitions.

<Step 12-1>

20 [0126] This is a step for oxidizing a compound (12a) to a compound (12b), and can be performed in the same way as in <Step 1B-1>.

[Production method 13]

[0127] A production method of a compound (13l), which is the compound represented by the formula (Ia), wherein Y is an oxygen atom, a sulfur atom or the formula $-NR_Y-$ (wherein R_Y represents a hydrogen or a C_1 to C_6 alkyl group), and X_1 is a group represented by the formula $-C(CN)=$



wherein, X_{203} represents a chlorine atom, a bromine atom or an iodine atom; other symbols represent the same definition as the aforementioned definition.

5 <Step 13-1>

[0128] This is a step for converting X_{203} of 4,6-dihalogenated nicotinic acid or its analogous compound (13a) such as 4,6-dichloronicotinic acid as reported in Acad. Nauk Ukr. SSSR, 1986, page 36 into an amino group
10 to obtain a compounds (13b). The reaction can be performed at a temperature of 0 °C to reflux temperature for 10 minutes to 100 hours by using, for example, an ammonia-ethanol solution or the like.

15 <Step 13-2>

[0129] This is a step for obtaining a compound (13c) by converting a carboxyl group of a compound (13b) into a carbamoyl group. For example, a method in which, after oxalyl chloride or thionyl chloride is allowed to react
20 with the compound at a temperature of 0 °C to reflux temperature for 10 minutes to 24 hours, ammonia is allowed to react with the compound, or a method in which diethylcyanophosphate, ammonium chloride, triethylamine are employed as disclosed in Synthesis
25 [SYNTBF], 1998, 1467 - 1475 or the like can be used.

<Step 13-3><Step 13-4><Step 13-5>

[0130] These are steps for condensing a pyridine or pyrimidine derivative (13c) and an indole or indoline derivative. Compounds (13d), (13e), (13f) can be
5 obtained, respectively, by using an indole derivative (1b), indole derivatives (3c), (3g) having a carboxamide group at the 1-position, or indoline derivatives (3d), (3h) having a carboxamide group at the 1-position under the same conditions as in <Step
10 1A-1>.

<Step 13-6><Step 13-11>

[0131] These are steps for conducting carboxamidation of the 1-position of indole of compounds (13d), (13g)
15 to obtain compounds (13e), (13j) and can be performed in the same way as in <Step 1A-2>.

<Step 13-8><Step 13-12><Step 13-15>

[0132] These are steps for converting a carbamoyl group
20 of compounds (13d), (13h), (13k) into a cyano group to obtain compounds (13g), (13l). For example, a method in which phosphorus oxychloride, thionyl chloride, trifluoroacetic anhydride are allowed to react with the compounds at a temperature of 0 °C to reflux
25 temperature for 10 minutes to 24 hours can be used.

<Step 13-9><Step 13-10><Step 13-14>


[0133] These are steps for converting aminopyridine or aminopyrimidine derivatives (13e), (13f), (13j) into corresponding urea derivatives (13h), (13i), (13l), and
5 can be performed in the same way as in <Step 1A-3>.

<Step 13-7><Step 13-13>

[0134] These are steps for oxidizing indoline derivatives (13f), (13i) to indole derivatives (13e),
10 (13k), and can be performed in the same way as in <Step 3-9>.

[Production method 14]

[0135] A typical production method of compounds (14d),
15 (14e), (14f) by halogenation of compounds (14a), (14b), (14c)



wherein, each symbol represents the same definition as the aforementioned definition.

5 <Step 14-1><Step 14-2><Step 14-3>


[0136] These are steps for substituting a substituent in 6-membered heterocycle from a hydrogen atom to a halogen atom. A compound can be obtained from a corresponding compound respectively: a compound (14d) from a compound (14a), a compound (14e) from a compound (14b) and a compound (14f) from a compound (14c) as in <Step 10-1>.

10

[Production method 15]

15 [0137] A typical production method of compounds (15a), (15b), (15c) by substituting a halogen atom in 6-

membered heterocycle of compounds (14d), (14e), (14f)
to a cyano group



wherein, each symbol represents the same definition as
the aforementioned definition.

<Step 15-1><Step 15-2><Step 15-3>

[0138] These are steps for obtaining a compound from a
corresponding compound respectively: a compound (15a)
from a compound (14d), a compound (15b) from a compound
(14e) and a compound (15c) from a compound (14f) by
substituting a substituent in 6-membered heterocycle
from a halogen atom to a cyano group. 0.5 to 2.0
equivalent of zinc cyanide or 1.0 to 3.0 equivalent of
copper(I) cyanide, potassium cyanide, sodium cyanide,
trimethylsilylcyanide or the like to compounds (14d),

(14e) and (14f) can be used. In order to accelerate the reaction, as a catalyst, for example, a palladium catalyst such as tetrakis(triphenylphosphine)palladium or dichlorobis(triphenylphosphine)palladium, copper(I) iodide, copper(0) or the like can be used. As a reaction solvent, for example, N,N-dimethylformamide, N-methylpyrrolidone, dimethyl sulfoxide, dioxane, tetrahydrofuran or the like can be used. The reaction can be performed at a temperature of room temperature to reflux temperature for 10 minutes to 2 days.

[0139] After completing the aforementioned reactions, purification can be performed by an ordinary treatment method, for example, column chromatography using silica gel or adsorbent resins or the like, or recrystallization from a suitable solvent.

[0140] The compounds of the invention, salts thereof or hydrates of the foregoing may be formulated as tablets, powders, fine particles, granules, coated tablets, capsules, syrups, lozenges, inhalants, suppositories, injections, ointments, eye salves, eye drops, nasal drops, ear drops, paps, lotions and the like, by any common methods. The formulation may employ any commonly used excipients, binders, lubricants, coloring agents, corrective coatings, and if necessary,

stabilizers, emulsifiers, absorbefaciants, surfactants, pH adjustors, preservatives, antioxidants, or the like, in combination with various components that are ordinarily used as raw materials for pharmaceutical formulations. For example, an oral formulation may be prepared by combining a compound of the invention or pharmacologically acceptable salt thereof with an excipient, if necessary adding a binder, disintegrator, lubricant, coloring agent, corrective coating or the like, and forming a powder, fine particles, granules, tablets, coated tablets, capsules, etc. by a common method. As such components there may be mentioned animal and vegetable oils such as soybean oil, beef tallow and synthetic glycerides; hydrocarbons such as liquid paraffin, squalane and solid paraffin; ester oils such as octyldodecyl myristate and isopropyl myristate; higher alcohols such as cetostearyl alcohol and behenyl alcohol; silicone resins; silicone oils; surfactants such as polyoxyethylene fatty acid esters, sorbitan fatty acid esters, glycerin fatty acid esters, polyoxyethylene sorbitan fatty acid esters, polyoxyethylene hydrogenated castor oil and polyoxyethylene-polyoxypropylene block copolymer; water-soluble polymers such as hydroxyethylcellulose, polyacrylic acid, carboxyvinyl polymer, polyethylene glycol, polyvinylpyrrolidone and methylcellulose; lower

alcohols such as ethanol and isopropanol; polyhydric alcohols such as glycerin, propylene glycol, dipropylene glycol and sorbitol; sugars such as glucose and sucrose; inorganic powders such as silicic acid anhydride, magnesium aluminum silicate and aluminum silicate, purified water, and the like. Examples of excipients which may be used include lactose, corn starch, white soft sugar, glucose, mannitol, sorbit, crystalline cellulose and silicon dioxide, examples of binders which may be used include polyvinyl alcohol, polyvinyl ether, methylcellulose, ethylcellulose, gum arabic, tragacanth, gelatin, shellac, hydroxypropylmethylcellulose, hydroxypropylcellulose, polyvinylpyrrolidone, polypropylene glycol/polyoxyethylene block polymer and meglumine, examples of disintegrators which may be used include starch, agar, gelatin powder, crystalline cellulose, calcium carbonate, sodium bicarbonate, calcium citrate, dextrin, pectin and carboxymethylcellulose calcium, examples of lubricants which may be used include magnesium stearate, talc, polyethylene glycol, silica and hydrogenated vegetable oils, examples of coloring agents which may be used include those approved for addition to drugs, and examples of corrective coatings which may be used include cocoa powder, menthol, aromatic powders, mentha oil, borneol and powdered

cinnamon. The tablets or granules may also be sugar coated or provided with another type of suitable coating if necessary. For preparation of a liquid formulation such as a syrup or injection, a common method may be used to formulate a compound of the invention or a pharmacologically acceptable salt thereof with a pH adjustor, solubilizer, isotonizing agent or the like, as well as a solubilizing aid, stabilizer etc. if necessary. There are no particular restrictions on the method of preparing an external agent, and any common method may be employed. That is, it may be prepared using as base materials any of various raw materials which are ordinarily used in drugs, quasi drugs, cosmetics and the like. As examples of specific base materials there may be mentioned raw materials such as animal and vegetable oils, mineral oils, ester oils, waxes, higher alcohols, fatty acids, silicone oils, surfactants, phospholipids, alcohols, polyhydric alcohols, water-soluble polymers, clay minerals, purified water and the like, and if necessary pH adjustors, antioxidants, chelating agents, antiseptics and fungicides, coloring agents, aromas and the like may also be added, although the base materials for external agents according to the invention are not limited to these. If necessary, there may also be included components such as ingredients having

differentiation-inducing activity, circulation
promoters, microbicides, antiphlogistic agents, cell
activators, vitamins, amino acids, humectants,
keratolytic agents and the like. The amounts of the
5 aforementioned base materials may be the concentrations
established for preparation of ordinary external agents.

[0141] There are no particular restrictions on the
compound of the invention, the salt thereof or the
10 hydrate thereof when administered, and either oral or
parenteral administration may be carried out according
to ordinary methods. For example, it may be prepared
and administered in the form of a tablet, powder, a
granule, a capsule, syrup, lozenge, inhalant,
15 suppository, injection, ointment, eye salve, eye drop,
nasal drop, ear drop, pap, lotion or the like.

[0142] Although the dosage of a drug according to the
invention will differ depending on severity of symptoms,
20 age, gender, body weight, form of administration, type
of disease, etc., it will be generally 100 μ g - 10 g
per day for an adult and such dosages may be
administered once or divided over several.

25 [0143] The administration form of the medicine
according to the present invention is not particularly

restricted, and can be an oral administration or a parenteral administration by a generally employed method.

5 [0144] The biochemical activity and actions and effects (angiogenesis inhibition activity, antitumor activity or the like) as a medicine of the compounds according to the present invention can be evaluated by the following methods.

10

[0145] The following is a list of abbreviations used in the pharmacological test examples described below.

<List of Abbreviations>

DNA (deoxyribonucleic acid)

15 VEGFR2 (vascular endothelial growth factor receptor 2)

Hepes (N-[2-Hydroxyethyl]piperazine-N'-[2-ethanesulfonic acid], HEPES (buffer solution))

MgCl₂ (Magnesium Chloride)

MnCl₂ (Manganese Chloride)

20 Na₃VO₄ (Sodium Orthovanadate(V))

ATP (Adenosine 5'-Triphosphate)

EDTA (Ethylenediaminetetraacetic acid)

HTRF (Homogenous Time-Resolved Fluorescence)

FGFR1 (Fibroblast growth factor receptor 1)

25 PDGFR β (Platelet derived growth factor receptor β)

HGFR (Hepatocyte growth factor receptor)

EGFR (Epidermal growth factor receptor)

Tris (Tris (hydroxymethyl)aminomethane, Tris (buffer solution))

NaCl (sodium Chloride)

5 BSA (Bovine Serum Albumin)

HRP (Horseradish peroxidase)

EGTA (Ethylene glycol bis(2-aminoethyl ether)-N,N,N',N'-tetraacetic acid)

SDS (Sodium Dodecylsulphate)

10 NP-40 (Nonidet P-40)

PCR: polymerase chain reaction

RT-PCR: reverse transcription-polymerase chain reaction

RNA: ribonucleic acid

cDNA: complementary DNA

15 cRNA: complementary RNA

dNTP: a mixture composed of dATP, dCTP, dGTP, and dTTP

UTP: Uridine 5'-triphosphate

CTP: Cytidine 5'-triphosphate

dATP: 2'-Deoxyadenosine 5'-triphosphate

20 dCTP: 2'-Deoxycytidine 5'-triphosphate

dGTP: 2'-Deoxyguanosine 5'-triphosphate

dUTP: 2'-Deoxyuridine 5'-triphosphate

GAPDH: glyceraldehyde 3-phosphate dehydrogenase

FBS: Fetal bovine serum

25 PBS: Phosphate buffered saline

MTT: (3-[4,5-Dimethylthiazol-2-yl]-2,5-

diphenyltetrazolium bromide; Thiazolyl blue)

DMSO: Dimethyl sulfoxide

PDGF: Platelet derived growth factor

EGF: Epidermal growth factor

5 FGF2: Fibroblast growth factor 2

VEGF: Vascular endothelial growth factor

HGF: Hepatocyte growth factor

TNF- α : Tumor Necrosis factor alpha

FCS: Fetal Calf Serum

10 EGM-2: Endothelial Cell Growth Medium-2

Pharmacological Test Example 1: Inhibition against
sandwich tube formation by vascular endothelial cells
in response to stimulation by angiogenesis factor

15 [0146] Human Umbilical Vein Endothelial Cells (HUVECs)
were isolated according to a reported method
(Shinseikagaku Jikken Koza [New Biochemistry Experiment
Lectures], "Saibo Baiyo Gijutsu" [Cell Culturing
Techniques], p.197-202),

20 and were cultured in a 5% CO₂ incubator (37°C) using
EGM-2 medium (purchased from Clonetics Corp.) until the
cells reached confluency.

25 [0147] An ice-cooled mixture of collagen: 5x RPMI 1640:
reconstitution buffer (all purchased from Nitta Gelatin,
Inc.) at 7:2:1 was dispensed at 0.4 ml into each well

of a 24-well plate. After the solution was gelled by
being stationed for 40 minutes in a 5% CO₂ incubator
(37°C), HUVEC cell suspension was added at 0.4 ml to
each well (using 1 to 1.2 x 10⁵ cells, though the
5 numbers of cells differ slightly according to the HUVEC
lot), the HUVEC cell suspension being in human
endothelial serum free medium (SFM, purchased from
GIBCO BRL) containing added angiogenesis factors [20
ng/ml FGF2 (purchased from GIBCO BRL) and 10 ng/ml EGF
10 (purchased from GIBCO BRL), or 25 ng/ml VEGF (purchased
from Wako Pure Chemical Industries Co., Ltd.) and 10
ng/ml EGF, or 30 ng/ml HGF (purchased from R&D Co.) and
10 ng/ml EGF], and cultured overnight in a 5% CO₂
incubator (37°C). On the following day, the medium on
15 the upper layer was aspirated off, and then 0.4 ml of
an ice-cooled mixture of collagen: 5x RPMI 1640:
reconstitution buffer (all purchased from Nitta Gelatin,
Inc.) at 7:2:1 was superposed into each well prior to
stationing for 4 hours in a 5% CO₂ incubator (37°C) for
20 gelling. After adding 1.5 ml of an SFM solution
containing each of the aforementioned angiogenesis
factors and a diluted test substance onto the upper
layer, culturing was performed in a 5% CO₂ incubator
(37°C). Upon aspirating off the culture supernatant in
25 each well on the 4th day after addition of the test
substance, 0.4 ml of a 3.3 mg/ml MTT solution dissolved

in PBS (purchased from Sigma Corp.) was added to each well and culturing was performed for approximately 2 hours in a 5% CO₂ incubator (37°C). The tubes formed in the collagen gel of each well were stained by MTT, the tube images were loaded into a computer (Macintosh), and the total length of the tubes was determined by image analysis software "MacScope" (purchased from Mitani Corp.). The ratio of the total length of the tubes formed in the well containing the test substance with respect to the total length of the tubes formed in the well containing no test substance was expressed as a percentage, and the concentration of each test substance required for 50% inhibition of tube formation (IC₅₀) was determined from the ratio value. The results are shown in Table 1.

[0148] [Table 1]

Example No.	VEGF-stimulated tube formation IC ₅₀ (nM)	FGF2-stimulated tube formation IC ₅₀ (nM)
39	5.1	470
41	2.1	250
46	7.0	470
47	5.8	120
53	6.7	440
78	3.0	450

Pharmacological Test Example 2: Measurement of
inhibition against receptor tyrosine kinase activity

5 [0149] This assay is used to determine inhibition of a
test substance on tyrosine kinase activity. DNA coding
for the cytoplasmic domain of VEGFR2 is obtained by
total cDNA synthesis (Edwards M, International
Biotechnology Lab 5(3), 19-25, 1987) or by cloning.
10 Expression in an appropriate expression system can
produce a polypeptide with tyrosine kinase activity.
The cytoplasmic domain of VEGFR2 obtained by expression
of recombinant protein in, for example, insect cells
have been found to exhibit intrinsic tyrosine kinase
15 activity. For VEGFR2 (GenBank Accession No. L04947),
the 1.7 kb DNA fragment described by Terman et al.
(Oncogene, 6(9), 1677-1683, 1991), coding for the

cytoplasmic domain, beginning with lysine 791 and including the termination codon, was isolated from a human placental cDNA library (purchased from Clontech Laboratories, Inc.) and cloned in a Baculovirus expression vector (pFastBacHis, purchased from GIBCO BRL). The recombinant construct was transfected into insect cells (*Spondoptea frugiperda*9 (Sf9)) to prepare a recombinant Baculovirus. (Instructions for preparation and use of recombinant Baculovirus may be found in standard texts, such as "Bac-To-Bac Baculovirus Expression System" (GIBCO BRL).) The cytoplasmic fragment starting from lysine 398 (FGFR1, GenBank Accession No. X52833), the cytoplasmic fragment starting from lysine 558 (PDGFR β , GenBank Accession No. M21616) or the cytoplasmic fragment starting from lysine 974 (HGFR, GenBank Accession No. J02958) may be cloned and expressed by the same method for use in assays for other tyrosine kinases. EGFR was purchased from Sigma Co. (Product No. E-2645).

[0150] For expression of the VEGFR2 tyrosine kinase, Sf9 cells were infected with the VEGFR2 recombinant virus and collected after 48 hours. The collected cells were rinsed with ice-cooled phosphate buffered saline (PBS) and then resuspended using 20 ml of ice-cooled Lysis Buffer (50 mM Tris-HCl (pH 8.5), 5 mM 2-

mercaptoethanol, 100 mM KCl, 1 mM phenylmethylsulfonyl fluoride, 1% (v/v) NP-40) per 1.5×10^8 cells. The suspension was centrifuged at 12,000 rpm for 30 minutes at 4°C and the supernatant was obtained. The
5 supernatant was added to a Ni-NTA agarose column (3 ml, purchased from Qiagen) equilibrated with Buffer A {20 mM Tris-HCl (pH 8.5), 5 mM 2-mercaptoethanol, 500 mM KCl, 20 mM imidazole, 10% (v/v) glycerol}. The column was washed with 30 ml of Buffer A, and then with 6 ml
10 of Buffer B {20 mM Tris-HCl (pH 8.5), 5 mM 2-mercaptoethanol, 1M KCl, 10% (v/v) glycerol}, and finally with 6 ml of Buffer A. After washing, it was eluted with 6 ml of Buffer C {20 mM Tris-HCl (pH 8.5), 5 mM 2-mercaptoethanol, 100 mM KCl, 100 mM imidazole,
15 10% (v/v) glycerol}. The eluate was placed on a dialysis membrane (purchased from Spectrum Laboratories) and dialyzed with a dialysis buffer {20 mM Tris-HCl (pH 7.5), 10% (v/v) glycerol, 1 mM dithiothreitol, 0.1 mM Na_3VO_4 , 0.1 mM EGTA}. After
20 dialysis, it was supplied for SDS-electrophoresis, and the recombinant protein (His6-VEGFR2, cytoplasmic domain of VEGFR2 fused with 6 histidine residues at the N-terminus) detected at a molecular weight of approximately 100 kDa with Coomassie Brilliant Blue
25 staining was assayed using BSA (bovine serum albumin, purchased from Sigma Co.) as the standard substance,

and stored at -80°C until use. Using the same method for the cytoplasmic domains of FGFR1, PDGFR β and HGFR yielded respective recombinant proteins fused with 6 histidine residues at the N-terminal (His6-FGFR1, His6-PDGFR β or His6-HGFR).

[0151] The tyrosine kinase reaction was conducted as follows. In the case of VEGFR2, for example, 10 μl of a kinase reaction solution {200 mM Hepes (pH 7.4), 80 mM MgCl_2 , 16 mM MnCl_2 , 2 mM Na_3VO_4 }, 250 ng of biotin-bound poly(Glu4:Tyr1) (biotin-poly(GT), purchased from CIS Diagnostics Co.) (6 μl of a 15-fold dilution with distilled water), 15 ng of His6-VEGFR2 (10 μl of a 240-fold dilution with 0.4% BSA solution) and the test substance dissolved in dimethyl sulfoxide (4 μl of a 100-fold dilution with 0.1% BSA solution) were added into each well of a 96-well round-bottom plate (NUNC Co., Product No. 163320), to a total of 30 μl . Next, 10 μl of 4 μM ATP (diluted with distilled water) was added prior to incubation at 30°C for 10 minutes, and then 10 μl of 500 mM EDTA (pH 8.0) was added.

[0152] The tyrosine phosphorylated biotin-poly(GT) was measured by the Homogenous Time-Resolved Fluorescence (HTRF) method (Analytical Biochemistry, 269, 94-104, 1999). Specifically, the kinase reaction solution was

transferred to a 96-well black half-plate (Product No. 3694, Coster, Inc.), 7.5 ng of europium cryptate-labeled anti-phosphotyrosine antibody (Eu(K)-PY20, purchased from CIS Diagnostics Co.) (25 μ l of a 250-fold dilution with 20 mM Hepes (pH 7.0), 0.5 M KF, 0.1% BSA solution) and 250 ng of XL665-labeled streptavidin (XL665-SA, purchased from CIS Diagnostics Co.) (25 μ l of a 62.5-fold dilution with 20 mM Hepes (pH 7.0), 0.5 M KF and 0.1% BSA solution) were added thereto, the mixture was allowed to stand at room temperature for 30 minutes, and then the fluorescent intensity was measured at 665 nm and 620 nm under irradiation with an excitation wavelength of 337 nm using a Discovery HTRF Microplate Analyzer (Packard Co.). The tyrosine phosphorylation rate for the biotin-poly(GT) was expressed as the delta F% value as described in the HTRF Standard Experiment Methods text by CIS Diagnostics Co. The delta F% value in the presence of the test substance was determined as a ratio (%) with the delta F% value with addition of His6-VEGFR2 in the absence of the test substance defined as 100% and the delta F% value in the absence of both the test substance and His6-VEGFR2 defined as 0%. This ratio (%) was used to calculate the test substance concentration required for 50% inhibition of VEGFR2 kinase activity (IC_{50}).

[0153] Measurement of inhibition against FGFR1, EGFR and HGFR kinase activity was conducted using 15 ng of His6-FGFR1, 23 ng of EGFR and 30 ng of His6-HGFR, respectively, according to the tyrosine kinase reaction and HTRF method described above. Measurement of inhibition against PDGFR β kinase activity was conducted using 50 ng of His6-PDGFR β according to the tyrosine kinase reaction described above, followed by detection of tyrosine phosphorylated biotin-poly(GT) by the following method. Specifically, the kinase reaction solution was added to a 96-well streptavidin-coated plate (Product No. 15129, Pierce Chemical) and incubated at room temperature for 30 minutes. After rinsing 3 times with 150 μ l of a rinsing solution {20 mM Tris-HCl (pH 7.6), 137 mM NaCl, 0.05% Tween-20, 0.1% BSA}, 70 μ l of anti-phosphotyrosine (PY20)-HRP conjugate (Product No. P-11625, Transduction Laboratories) {2000-fold dilution with 20 mM Tris-HCl (pH 7.6), 137 mM NaCl, 0.05% Tween-20, 1% BSA} was added thereto and incubation was performed at room temperature for 1 hour. After incubation, it was rinsed 3 times with 150 μ l of the rinsing solution, and 100 μ l of TMB Membrane Peroxidase Substrate (Product No. 50-5077-03, Funakoshi Co., Ltd.) was added to initiate the reaction. After stationing at room temperature for

10 minutes, 100 μ l of 1 M phosphoric acid was added to suspend the reaction, and the absorbance at 450 nm was measured with a microplate reader (BIO KINETICS READER EL304, Bio-Tek Instruments). The absorbance ratio in the presence of the test substance was determined with respect to 100% as the absorbance with addition of His6-PDGFR β and no test substance, and 0% as the absorbance without addition of both the test substance and His6-PDGFR β . This absorbance ratio was used to calculate the test substance concentration required for 50% inhibition of PDGFR β kinase activity (IC_{50}). The results are shown in Table 2.

[0154] [Table 2]

Example No.	VEGFR2 kinase IC ₅₀ (nM)	FGFR1 kinase IC ₅₀ (nM)	Example No.	VEGFR2 kinase IC ₅₀ (nM)	FGFR1 kinase IC ₅₀ (nM)
7	8.0	26	68	37	52
11	3.0	47	79	9.8	25
18	3.0	70	81	12	38
28	4.5	4.1	82	15	24
32	9.3	16	88	14	24
33	7.1	12	104	3.9	19
34	8.4	22	116	14	87
36	3.4	16	119	21	120
37	4.8	1.2	139	6.3	190
39	4.5	6.3	206	4.1	3.5
40	5.7	6.9	207	4.6	12
41	6.1	3.2	208	7.7	6.8
43	6.4	18	209	17	29
44	7.7	14	210	8.1	40
46	32	12	211	45	36
47	40	21	212	8.6	19
50	5.0	13	213	10	330
53	3.8	2.1			

Pharmacological Test Example 3: Evaluation of *in vivo* angiogenesis-inducing activity using mouse dorsal air

sac model

[0155] (1) Construction of VEGF (Vascular Endothelial Growth Factor) expression vector

5 PCR was conducted using a human placenta cDNA library (Toyobo Co., Ltd.) as the template and the SEQ ID NO:1 (5'CCGGATCCATGAACTTTCTGCTG3') and SEQ ID NO:2 (5'GTGAATTCTGTATCGATCGTT3') of VEGF as primers. After completion of the PCR reaction, the 5' ends were phosphorylated and an approximately 600 bp DNA band was
10 separated by 1.2% agarose gel electrophoresis. After polymerization by self-ligation, the cDNA was cut with *EcoRI* and *BamHI* and inserted into the *EcoRI* and *BamHI* sites of vector pUC19. This was used to transform *E. coli* JM83, and plasmids were recovered from the
15 transformed clones. A VEGF cDNA fragment was cut out of the plasmids with *HindIII* and *EcoRI* and then inserted into pIRES2-rsGFP vector to yield pIRES2-rsGFP/VEGF for protein expression.

20 [0156] (2) Preparation of VEGF high-expressing strain

After overnight culturing of KP-1 human pancreatic cancer cells (3×10^6 cells) with 10% FCS-containing RPMI 1640 medium, an Effectene Transfection Reagent Kit (Qiagen) was used for introduction of 3 μ g
25 of pIRES2-rsGFP/VEGF into the KP-1 cells. After culturing in 10% FCS-containing RPMI 1640 medium

containing 600 μ g/ml of Geneticin, drug-resistant cells were selected. Furthermore, GFP high-expressing cells were collected by cell sorter (Becton Dickinson) as VEGF high-expressing KP-1 cells (KP-1/VEGF).

5

[0157] (3) Measurement of VEGF level in culture supernatant

The KP-1/VEGF cells were prepared to 5×10^5 cells/ml, and 0.5 ml thereof was dispensed into each well of a 24-well plate and cultured at 37°C for 24 hours. The culture supernatants were collected and the VEGF levels thereof measured using a VEGF measuring kit (IBL Co., Ltd.) for confirmation of high expression.

15 [0158] (4) Evaluation of *in vivo* angiogenesis-inducing activity using mouse dorsal air sac model

Millipore rings (Nihon Millipore) were sealed with 0.45 μ m Durapore filter membranes (Nihon Millipore) to create chambers. KP-1/VEGF human pancreatic cancer cells (3×10^6) suspended in 0.17 ml of collagen gel were injected into each chamber through the injection port, and the chambers were sealed. Approximately 10 ml of air was then injected in the dorsal skin of 6-week-old C57BL/6N female mice under anesthesia to produce pouches, and the prepared chambers were transplanted therein. About 6 hours

after completing transplantation, a test substance suspended in 0.5% methyl cellulose was orally administered (0.1 ml/10 g body weight), and this was continued once a day for the next 4 days.

5

[0159] On the 4th day after transplanting the chambers, 0.2 ml of ^{51}Cr (Amersham Pharmacia)-labeled mouse erythrocytes (2.5×10^6 cpm/ml) were injected through the caudal veins of each of the mice with the transplanted chambers. After a prescribed period, the skin in contact with the chamber was excised and frozen, the section in direct contact with the chamber was precisely cut off, and the radioactivity was measured with a γ -counter. The blood volume was calculated from the radioactivity and used as an index of the *in vivo* angiogenesis-inducing activity. The angiogenesis volume was recorded as this measured blood volume minus the blood volume obtained with transplantation of a chamber containing only collagen gel. The experiment was conducted using 10 mice in the control (solvent-administered) group and 5 mice in each compound-administered group. The proportions (%) of the angiogenesis amount after administering a test substance to that of control are shown in Table 3.

25

[0160] [Table 3]

Example No.	Dose (mg/kg/day)	
	30	100
46	59.9%	43.7%
47	87.4%	39.5%
53	43.3%	44.4%

Pharmacological Test Example 4: Evaluation of antitumor activity on KP-1/VEGF cells in subcutaneous xenograft models

[0161] VEGF high-expressing pancreatic cancer cells (KP-1/VEGF) suspended in PBS at a concentration of 1×10^7 cells/ml were transplanted under the right flank skin of 6-week-old female Balb/c (nu/nu) mice in a volume of 0.1 ml. When the tumor volume reached approximately 100 mm³, the test substance was orally administered twice a day over a period of 2 weeks with a schedule of 5 days per week. The test substance was suspended in 0.5% methyl cellulose for an administered volume of 0.1 ml/10 g body weight. The tumor size was measured twice a week using a micrometer caliper. The tumor volume was determined by measuring the long and short diameters of the tumor with a micrometer caliper, and calculating $1/2 \times (\text{long diameter} \times \text{short diameter} \times \text{short diameter})$. The experiment was conducted using 10 mice in the control (solvent-

administered) group and 5 mice in each test substance-administered group. The proportions (%) of the tumor volume after administering a test substance to that of control are shown in Table 4.

5

[0162] [Table 4]

Example No.	Dose (mg/kg/day)			
	6	20	60	200
46	77.7%	59.3%	60.0%	36.2%
47	80.5%	64.6%	45.3%	33.6%
53	73.1%	60.4%	48.4%	37.9%

Pharmacological Test Examples 5: Evaluation of angiogenesis inhibition activity in mouse angiogenesis model by using Matrigel

10

[0163] The experiment was performed according to the method as already reported in the method (Lab. Invest., 67(4), 519 - 528, 1992). Specifically, 10 μ g/ml of recombinant FGF-2 (purchased from Invitrogen Corporation) dissolved in PBS was added to Matrigel Matrix (purchased from BD Biosciences) to prepare 1 μ g/ml. After that, a 300 μ l of this mixture of Matrigel Matrix and Recombinant FGF-2 was injected into a subcutaneous tissue on the median line of the abdomen of a 6-week-old Balb/c (nu/nu) mouse.

20

[0164] Subsequently, the test substance suspended in a 0.5% methyl cellulose or the like had been orally administered in succession once a day or twice a day for 7 days.

5

[0165] After 7 days, the implanted Matrigel was taken out, 300 μ l of water was added thereto, and cut into pieces with scissors. The resultant substance was allowed to stand at a cool dark place overnight. After hemoglobin in Matrigel was fully extracted, 100 μ l of the supernatant obtained by centrifugation and 100 μ l of Drabkin's solution (purchased from Sigma Chemical Co., Ltd) were allowed to react at room temperature at a dark place for 1 hour. After that, the absorbance of the reaction solution was measured with wavelength of 550nm and reference wavelength of 660 nm. The hemoglobin quantity (g/ml) in Matrigel was calculated from the calibration curve established by use of hemoglobin as a standard.

15
20

[0166] The experiment was conducted using 8 mice in the control (solvent-administered) group and 6 mice in each compound-administered group.

[Examples]

25

[0167] The compounds according to the present invention

can be prepared by the methods as described in the following examples, for example. These are, however, exemplary, and the compounds according to the present invention are not limited to the specific examples mentioned below in any cases.

Example 1

N1-Ethyl-5-(2-((methoxylamino)carbonyl)amino-4-pyrimidyl)oxy-1H-1-indolecarboxamide

[0168] Similarly to Production example 27-2, a crude product of phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyl)oxy-2-pyrimidyl)carbamate (546 mg, 1.31 mmol, 56.3%) was obtained as pale brown powder from N1-ethyl-5-(2-amino-4-pyrimidyl)oxy-1H-1-indolecarboxamide (691 mg, 2.32 mmol) and phenyl chlorocarbonate. The crude carbamate product (273 mg, 0.65 mmol) was dissolved in tetrahydrofuran (7.0 ml); and triethylamine (0.91 ml, 6.53 mmol) and methoxylamine hydrochloride (273 mg, 3.27 mmol) was added thereto while stirred at room temperature. After the reaction mixture was stirred overnight, the reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with water and brine, and was dried over anhydrous sodium sulfate. The solvent was distilled off, and the residue was purified by silica gel column chromatography (eluent; ethyl acetate: hexane = 1: 1).

The crystals were precipitated from ethyl acetate-hexane (1: 10), filtered off, and dried under aeration to yield the title compound (52.5 mg, 0.14 mmol, 21.7%) as white crystals.

5 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.17 (3H, t, J=7.2 Hz), 3.20-3.40 (2H, m), 3.68 (3H, s), 6.45 (1H, d, J=5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 7.09 (1H, dd, J=2.4, 8.8 Hz), 7.43 (1H, d, J=2.4 Hz), 7.54 (1H, d, J=5.6 Hz), 7.89 (1H, d, J=3.6 Hz), 8.21 (1H, m), 8.26 (1H, d, J=8.8 Hz), 8.34 (1H, d, J=5.6 Hz), 9.31 (1H, d, J=10.0 Hz).

[0169] The starting materials were synthesized by the following methods.

15 Production example 1-1

4-Chloro-6-(1H-5-indolyloxy)-2-pyrimidinamine

Sodium hydride (1.0 g, 25 mmol) was suspended in dimethyl sulfoxide (40 ml) under nitrogen atmosphere, and 5-hydroxyindole (3.33 g, 25 mmol) was gradually added while the reaction mixture was stirred at room temperature. After 20 minutes, 2-amino-4,6-dichloropyrimidine (3.28 g, 20 mmol) was added. The reaction mixture was heated at 100 °C and was stirred for 3 hours. After the reaction mixture was cooled to room temperature, the reaction mixture was partitioned between ethyl acetate and 10% aqueous ammonia solution.

The organic layer was washed with water and brine, and dried over anhydrous sodium sulfate. After the solvent was distilled off, the residue was purified by NH silica gel column chromatography (eluent; ethyl acetate: hexane = 2: 1). The crystals were precipitated from ethyl acetate, filtered off, and dried under aeration to yield the title compound (1.15 g, 4.41 mmol, 22.0%) as white crystals.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 5.09 (2H, brs), 6.07 (1H, s), 6.57 (1H, m), 6.95 (1H, dd, J=2.4, 8.8 Hz), 7.29 (1H, m), 7.37 (1H, m), 7.41 (1H, d, J=8.8 Hz), 8.28 (1H, brs).

Production example 1-2

4-(1H-5-Indolyloxy)-2-pyrimidinamine

[0170]

4-Chloro-6-(1H-5-indolyloxy)-2-pyrimidinamine (1.15 g, 4.41 mmol) was dissolved in tetrahydrofuran (50 ml)-triethylamine (3.07 ml), 10% palladium on carbon (50% wet, 500 mg) was added, and the reaction mixture was stirred overnight under hydrogen atmosphere at atmospheric pressure.

The reaction was purged with nitrogen. After methanol (50 ml) was added and stirred, the catalyst was filtered out. The resultant solution was concentrated under reduced pressure, thus the title compound (826 mg, 3.65 mmol, 82.8%) was obtained as

pale gray powder.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 4.96 (2H, brs), 6.06 (1H, d, J=5.6 Hz), 6.56 (1H, m), 6.97 (1H, dd, J=2.4, 8.8 Hz), 7.26-7.28 (1H, m), 7.38-7.42 (2H, m), 8.08 (1H, d, J=8.8 Hz), 8.29 (1H, brs).

Production example 1-3

N1-Ethyl-5-(2-amino-4-pyrimidyl)oxy-1H-1-indolecarboxamide

[0171] Sodium hydride (157 mg, 3.93 mmol) was suspended in N,N-dimethylformamide (10 ml) under nitrogen atmosphere, and 4-(1H-5-indolyloxy)-2-pyrimidinamine (826 mg, 3.65 mmol) was gradually added while the reaction mixture was stirred at room temperature. After 10 minutes, the reaction mixture was cooled with an ice-water bath, phenyl N-ethylcarbamate (633 mg, 3.83 mmol) was added, the reaction mixture was heated to room temperature, and the solution was stirred for 4 hours. The reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with water and brine and dried over anhydrous sodium sulfate. The solvent was distilled off, and the residue was purified by silica gel chromatography (eluent; ethyl acetate: hexane = 3: 1 to 4: 1) to yield the title compound (691 mg, 2.32 mmol, 63.7%) as white powder.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 1.32 (3H, t, J=7.2 Hz), 3.54 (2H, m), 4.94 (2H, brs), 5.50 (1H, brs), 6.11 (1H, dd, J=2.4, 5.6 Hz), 6.62 (1H, d, J=3.6 Hz), 7.09 (1H, dd, J=2.4, 8.8 Hz), 7.34 (1H, d, J=2.4 Hz), 7.46 (1H, d, J=3.6 Hz), 8.11 (1H, d, J=5.6 Hz), 8.15 (1H, d, J=8.8 Hz).

Example 2

5-(6-(3-(3-Diethylaminopropylamino)ureido)pyrimidin-4-yloxy)-1H-indole-1-carboxylic acid methylamide
[0172] Phenyl (6-(1-methylcarbamoyl-1H-indol-5-yloxy)pyrimidin-4-yl)carbamate (161 mg, 0.400 mmol) was dissolved in N,N-dimethylformamide (1.0 ml), and 3-(diethylamino)propylamine (130 mg, 1.00 mmol) was added while the reaction mixture was stirred at room temperature. After the reaction mixture was stirred overnight, the reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with water and brine, and dried over anhydrous sodium sulfate. The solvent was distilled off, and the residue was purified by NH silica gel column chromatography (eluent; ethyl acetate: methanol = 50:1). The crystals were precipitated from ethyl acetate-hexane, filtered off, and dried under aeration to yield the title compound (123 mg, 0.280 mmol, 70%) as white crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.93 (6H, t, J=7.0 Hz), 1.52 (2H, m), 2.32-2.46 (6H, m), 2.84 (3H, d, J=3.6 Hz), 3.12 (2H, m), 6.69 (1H, d, J=3.6 Hz), 6.98 (1H, s), 7.06 (1H, dd, J=2.2, 8.8 Hz), 7.37-7.46 (2H, m), 7.88 (1H, d, J=3.6 Hz), 8.18 (1H, m), 8.27 (1H, d, J=8.8 Hz), 8.37 (1H, s), 9.49 (1H, brs).

[0173] The starting materials were synthesized by the following methods.

10 Production example 2-1

Phenyl N-methylcarbamate

Methylamine hydrochloride (16.9 g, 250 mmol) was dissolved in N,N-dimethylformamide (250 ml), pyridine (44 ml, 275 mmol) was added thereto, and the reaction mixture was stirred. The reaction mixture was cooled with ice, phenyl chloroformate (35 ml, 275 mmol) was added dropwise thereto, and the reaction mixture was then stirred at room temperature for 24 hours. The reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and the solvent was distilled off under reduced pressure. The obtained crystals were suspended in diethylether, diluted with hexane, filtered off, washed with the diethylether: hexane = 1: 1, and dried by evacuation, to yield the title compound (22.3 g, 147 mmol, 59.1%) as colorless

crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.64 (3H, d, J=3.6 Hz), 7.07 (2H, d, J=8.0 Hz), 7.17 (1H, t, J=8.4 Hz), 7.35 (2H, dd, J=8.0 Hz, 8.4 Hz), 7.58 (1H, d, J=3.6 Hz).

5 [0174]

Production example 2-2

6-(1H-Indol-5-yloxy)pyrimidin-4-ylamine

Sodium hydride (400 mg, 10.0 mmol) was suspended in dimethyl sulfoxide (20 ml) under nitrogen atmosphere, and 5-hydroxyindole (1.33 g, 10.0 mmol) was gradually added while the reaction mixture was stirred at room temperature. After 20 minutes, 6-chloropyrimidin-4-ylamine (1.04 g, 8.00 mmol) was added thereto, the reaction mixture was heated at 100 °C and stirred for 1 hour. After the reaction mixture was naturally cooled to room temperature, the reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with water and brine, and was dried over anhydrous magnesium sulfate. The solvent was distilled off, and the residue was purified by silica gel column chromatography (eluent; ethyl acetate: hexane = 3: 1) to yield the title compound (1.07 g, 4.73 mmol, 59%) as a brown oil.

25 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 5.54 (1H, s), 6.43 (1H, m), 6.71 (2H, brs), 6.85 (1H, dd, J=2.4, 8.8 Hz), 7.29 (1H, d, J=2.4 Hz), 7.40-7.45 (2H, m), 8.06 (1H, s),

11.20 (1H, brs).

[0175]

Production example 2-3

5-(6-Aminopyrimidin-4-yloxy)-1H-indol-1-carboxylic acid
methanamide

Sodium hydride (199 mg, 4.97 mmol) was suspended in N,N-dimethylformamide (10 ml) under nitrogen atmosphere, 6-(1H-indol-5-yloxy)pyrimidin-4-ylamine (1.07 g, 4.73 mmol) synthesized in Production example 2-2 was gradually added while the reaction mixture was stirred at room temperature. After 30 minutes, the reaction mixture was cooled with an ice water bath, then phenyl N-methylcarbamate (751 mg, 4.97 mmol) synthesized in Production example 2-1 was added. The reaction mixture was heated to room temperature and stirred for 1 hour. The reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with water and brine, and was dried over anhydrous magnesium sulfate. After the solvent was distilled off, the residue was purified by silica gel column chromatography (eluent; ethyl acetate) to yield the title compound (847 mg, 2.99 mmol, 63%) as white crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.85 (3H, d, J=4.0 Hz), 5.62 (1H, s), 6.68 (1H, d, J=3.6 Hz), 6.77 (2H, brs), 7.04 (1H, dd, J=2.4, 9.2 Hz), 7.36 (1H, d, J=2.4

Hz), 7.87 (1H, d, J=3.6 Hz), 8.07 (1H, s), 8.15 (1H, q, J=4.0 Hz), 8.27 (1H, d, J=9.2 Hz).

Production example 2-4

5 Phenyl (6-(1-methylcarbamoyl-1H-indol-5-yloxy)pyrimidin-4-yl)carbamate

[0176] 5-(6-Aminopyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide (847 mg, 2.99 mmol) synthesized in Production example 2-3 was dissolved in N,N-dimethylformamide (10 ml) under nitrogen atmosphere. Pyridine (0.290 ml, 11.5 mmol) and phenyl chlorocarbonate (0.394 ml, 3.15 mmol) were sequentially added dropwise thereto while cooling with an ice water bath. After the reaction mixture was stirred for 30 minutes, triethylamine (0.417 ml, 2.99 mmol) was added, and the reaction mixture was heated to room temperature while stirred. After 30 minutes, the reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with water and brine, and was dried over anhydrous magnesium sulfate. The solvent was distilled off, and then the residue was purified by silica gel column chromatography (eluent; ethyl acetate: hexane = 3: 1). The crystals were precipitated from ethyl acetate-hexane, filtered off, and dried under aeration to yield the title compound (504 mg, 1.25 mmol, 42%) as white crystals.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 3.05 (3H, d, J=4.8 Hz),
5.53 (1H, q, J=4.8 Hz), 6.58 (1H, d, J=4.0 Hz), 7.08
(1H, dd, J=2.4, 8.8 Hz), 7.13-7.19 (2H, m), 7.23-7.29
(1H, m), 7.34 (1H, d, J=2.4 Hz), 7.36-7.44 (3H, m),
5 7.52 (1H, s), 8.14 (1H, d, J=8.8 Hz), 8.59 (1H, s),
9.99 (1H, brs).

Example 3

10 5-(6-(((4-Hydroxypiperidin-1-yl)carbonyl)amino)pyrimidin-4-yloxy)-1H-indole-1-
carboxylic acid methylamide

[0177] Similarly to Example 2, the title compound (100
mg, 0.231 mmol, 58%) was obtained as white powder from
phenyl (6-(1-methylcarbamoyl-1H-indol-5-
15 yloxy)pyrimidin-4-yl)carbamate (161 mg, 0.400 mmol) and
4-hydroxypiperidine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.24-1.34 (2H, m),
1.64-1.73 (2H, m), 2.85 (3H, d, J=4.0 Hz), 3.02-3.12
(2H, m), 3.64 (1H, m), 3.72-3.80 (2H, m), 4.69 (1H, d,
20 J=4.0 Hz), 6.68 (1H, d, J=3.6 Hz), 7.06 (1H, dd, J=2.4,
8.8 Hz), 7.20 (1H, s), 7.40 (1H, d, J=2.4 Hz), 7.88 (1H,
d, J=3.6 Hz), 8.17 (1H, q, J=4.0 Hz), 8.27 (1H, d,
J=8.8 Hz), 8.40 (1H, s), 9.72 (1H, brs).

25 Example 4

5-(6-((4-(Pyrrolidin-1-yl)piperidin-1-

yl)carbonylamino)pyrimidin-4-yloxy)-1H-indole-1-
carboxylic acid methylamide

[0178] Similarly to Example 2, the title compound (141 mg, 0.304 mmol, 76%) was obtained as white crystals from phenyl (6-(1-methylcarbamoyl-1H-indol-5-yloxy)pyrimidin-4-yl)carbamate (161 mg, 0.400 mmol) and 4-(1-pyrrolidynyl)piperidine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.23-1.36 (2H, m), 1.63-1.70 (4H, m), 1.74-1.84 (2H, m), 2.08-2.18 (1H, m), 2.42-2.50 (4H, m), 2.82-2.95 (5H, m), 3.90-3.98 (2H, m), 6.68 (1H, d, J=3.6 Hz), 7.06 (1H, dd, J=2.4, 8.8 Hz), 7.20 (1H, s), 7.40 (1H, d, J=2.4 Hz), 7.88 (1H, d, J=3.6 Hz), 8.17 (1H, q, J=4.0 Hz), 8.27 (1H, d, J=8.8 Hz), 8.40 (1H, s), 9.71 (1H, brs).

Example 5

5-(2-(3-((1R)-1-Carbamoyl-2-phenylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0179] Phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (104 mg, 0.200 mmol) and triethylamine (1 ml) were dissolved in N,N-dimethylformamide (3 ml), and (2R)-2-amino-3-phenylpropionamide hydrochloride (201 mg, 1.00 mmol) was added, and the reaction mixture was stirred for 18 hours. The reaction mixture was partitioned between ethyl acetate and the saturated aqueous

solution of ammonium chloride. The organic layer was washed with water and brine, and was dried over anhydrous magnesium sulfate. The solvent was distilled off, and the residue was purified by silica gel column chromatography (eluent; ethyl acetate: methanol = 50: 1). The crystals were precipitated from a solvent mixture of ethyl acetate-hexane, filtered off, and dried under aeration to yield the title compound (77.2 mg, 0.152 mmol, 76%) as white crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.81 (1H, dd, J=8.0, 13.2 Hz), 2.84 (3H, d, J=4.4 Hz), 3.01 (1H, dd, J=4.8, 13.2 Hz), 4.38 (1H, m), 6.52 (1H, dd, J=2.4, 6.0 Hz), 6.69 (1H, d, J=3.2 Hz), 6.86 (1H, s), 7.01-7.07 (2H, m), 7.15-7.30 (5H, m), 7.37 (1H, d, J=2.4 Hz), 7.50 (1H, s), 7.88 (1H, d, J=3.2 Hz), 8.02 (1H, d, J=6.0 Hz), 8.18 (1H, q, J=4.4 Hz), 8.22-8.34 (2H, m), 9.11 (1H, s).

[0180] The starting material, Phenyl N-(4-(1-(methylamino)carbonyl-1H-indol-5-yloxy)pyridin-2-yl)-N-(phenoxycarbonyl) carbamate, was synthesized as follows.

Production example 5-1

N1-Methyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide

Sodium hydride (430 mg, 10.75mmol) was suspended in N,N-dimethylformamide (25 ml) under nitrogen atmosphere, and 4-(1H-5-indolyloxy)-2-pyridinamine

(2.253 g, 10.00 mmol, CAS No. 417722-11-3) described in WO 02/32872 was gradually added while stirred at room temperature. After 10 minutes, the reaction mixture was cooled with an ice water bath, and then phenyl N-methylcarbamate (1.587 g, 10.50 mmol) was added. The reaction mixture was heated to room temperature and stirred for 2 hours. The reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with water and brine, and was dried over anhydrous sodium sulfate. The solvent was removed by distilled off. The crystals were precipitated from ethyl acetate, filtered off, and dried under aeration to yield the title compound (2.163 g, 7.66 mmol, 76.6%) as pale brown crystals.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 3.09 (3H, d, J=4.8 Hz), 4.36 (2H, m), 5.49 (1H, m), 5.92 (1H, d, J=2.0 Hz), 6.30 (1H, dd, J=2.0, 6.0 Hz), 6.61 (1H, d, J=3.6 Hz), 7.07 (1H, dd, J=2.4, 8.8 Hz), 7.30 (1H, d, J=2.4 Hz), 7.45 (1H, d, J=3.6 Hz), 7.92 (1H, d, J=6.0 Hz), 8.17 (1H, d, J=8.8 Hz).

Production example 5-2

phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate

[0181] N1-Methyl-5-(2-amino-pyridyl)oxy-1H-1-indolecarboxamide (2.0 g, 7.1 mmol) was suspended in

tetrahydrofuran (140 ml) and N,N-dimethylformamide (1.4 ml) at room temperature, and triethylamine (2.2 ml, 16 mmol) was added while stirred. The reaction mixture was cooled with an ice, and phenyl chloroformate (1.8 ml, 15 mmol) was added, and the reaction mixture was stirred at room temperature for 1.5 hours. Phenyl chloroformate (0.5 ml) was further added, and the reaction mixture was stirred at room temperature for 0.5 hours. Brine was added to the reaction mixture; and this was subjected to extraction with ethyl acetate, washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. Diethyl ether was added to the residue, then the precipitated crystals were filtered off, washed with diethyl ether, and dried under aeration to yield the title compound (3.3 g, 6.3 mmol, 89%) as pale brown crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 3.30 (3H, d, J=4.4 Hz), 6.66 (1H, d, J=3.6 Hz), 6.95 (1H, dd, J=2.4, 6.0 Hz), 7.10 (1H, dd, J=2.4, 8.8 Hz), 7.15-7.18 (4H, m), 7.27-7.31 (2H, m), 7.40-7.45 (5H, m), 7.52 (1H, d, J=2.4 Hz), 7.88 (1H, d, J=3.6 Hz), 8.17 (1H, q, J=4.4 Hz), 8.31 (1H, d, J=8.8 Hz), 8.41 (1H, d, J=6.0 Hz).

[0182] N1-methyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide described in Production example 5-1,

can be also synthesized as follows.

N1-Methyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide

5- (2-Aminopyridin-4-yloxy) -2,3-dihydro-1H-indole-1-carboxylic acid methylamide (40 mg, 0.14 mmol) was dissolved in acetic acid (0.9 ml), manganese (III) acetate (29 mg, 0.17 mmol) was added thereto and the reaction mixture was stirred at 70 °C for 3.5 hours. Manganese (III) acetate (29 mg, 0.17 mmol) was further added, and the reaction mixture was further stirred at 70 °C for 0.5 hours. After naturally cooled to room temperature, the reaction mixture was partitioned between ethyl acetate and saturated aqueous solution of sodium hydrogencarbonate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The obtained crystals were suspended in diethyl ether: acetone = 3:1, filtered off, washed with diethyl ether, and dried under aeration to yield the title compound (24 mg, 0.085 mmol, 61%) as colorless crystals.

[0183] The starting material, 5-(2-Aminopyridin-4-yloxy)-2,3-dihydro-1H-indole-1-carboxylic acid methylamide was synthesized as follows.

Production example 5-3

5-Benzyloxy-1H-indole-1-carboxylic acid methylamide

Sodium hydride (2.212 g, 55.30 mmol, 60% in oil) was suspended in N,N-dimethylformamide (100 ml), 5-benzyloxyindole (10.29 g, 46.09 mmol) was added thereto while stirred at room temperature, and the reaction mixture was stirred at room temperature for 40 minutes. The reaction mixture was cooled with an ice water bath, and phenyl N-methylcarbamate (8.360 g, 55.30 mmol) was added. After the reaction mixture was stirred for 30 minutes, the solution was stirred at room temperature for 2.5 hours. After water was added to the reaction mixture and the reaction mixture was stirred at room temperature for 1 hour, the obtained crystals were filtered off, then these crystals were sequentially washed with water and diethyl ether, and dried under aeration to yield the title compound (12.07 g, 43.06 mmol, 93.41%) as pale yellow crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.80 (3H, d, J=4.4 Hz), 5.10 (2H, s), 6.56 (1H, d, J=3.8 Hz), 6.93 (1H, dd, J=2.4, 9.0 Hz), 7.16 (1H, d, J=2.4 Hz), 7.30 (1H, t, J=7.2 Hz), 7.37 (2H, t, J=7.2 Hz), 7.45 (2H, d, J=7.2 Hz), 7.74 (1H, d, J=3.8 Hz), 8.00 (1H, m), 8.11 (1H, d, J=9.0 Hz).

Production example 5-4

5-Hydroxy-2,3-dihydro-1H-indole-1-carboxylic acid methylamide

[0184] 5-Benzyloxy-1H-indole-carboxylic acid
methanamide (10.00 g, 35.67 mmol) was dissolved in
methanol (200 ml) and tetrahydrofuran (150 ml), 10%
palladium on carbon (0.9 g) was added, and the reaction
5 mixture was stirred at room temperature under hydrogen
atmosphere for 9 hours. After the catalyst was removed
by filtration, the solvent was distilled off under
reduced pressure. The residue was dissolved in ethanol
(400 ml), 10% palladium on carbon (0.9 g) was added,
10 then the reaction mixture was stirred at room
temperature under hydrogen atmosphere for 26 hours.
After the catalyst was removed by filtration, the
solvent was distilled off under reduced pressure. The
obtained crystals were suspended in diethyl ether,
15 filtered off, washed with diethyl ether, and dried
under aeration to yield the title compound (6.522 g,
33.93 mmol, 95.12%) as grayish crystals.
¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.61 (3H, d, J=4.4
Hz), 2.99 (2H, t, J=8.6 Hz), 3.76 (2H, t, J=8.6 Hz),
20 6.33 (1H, d, J=4.4 Hz), 6.43 (1H, dd, J=2.4, 8.4 Hz),
6.54 (1H, d, J=2.4 Hz), 7.58 (1H, d, J=8.4 Hz), 8.82
(1H, s).

Production example 5-5

25 5-(2-Aminopyridin-4-yloxy)-2,3-dihydro-1H-indole-1-
carboxylic acid methanamide

[0185] Sodium hydride (202 mg, 3.89 mmol, 60% in oil) was suspended in dimethyl sulfoxide (5.0 ml), then 5-hydroxy-2,3-dihydro-1H-indole-1-carboxylic acid methylamide (971 mg, 5.06 mmol) and 2-amino-4-chloropyridine (500 mg, 3.89 mmol) were added at room temperature under nitrogen atmosphere, and the reaction mixture was heated and stirred at 160 °C for 12 hours under nitrogen atmosphere. After naturally cooled down to room temperature, the reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, was dried over anhydrous magnesium sulfate, and was concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Fuji Silysia BW-300; eluent: ethyl acetate, ethyl acetate: methanol = 85: 10 in this order). The fractions containing the desired compound were concentrated, and the residue was further purified by silica gel column chromatography (Fuji Silysia NH, eluent; from ethyl acetate to ethyl acetate: methanol = 90: 10). The obtained crystals were suspended in diethyl ether: acetone = 3: 1, filtered off, washed with diethyl ether, and dried under aeration to yield the title compound (51 mg, 0.18 mmol, 4.6%) as pale green crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.65 (3H, d, J=4.4 Hz), 3.09 (2H, t, J=8.6 Hz), 3.86 (2H, t, J=8.6 Hz),

5.75 (1H, d, J=2.0 Hz), 5.85 (2H, brs), 6.07 (1H, dd, J=2.0, 6.0 Hz), 6.56 (1H, d, J=4.4 Hz), 6.81 (1H, dd, J=2.4, 8.4 Hz), 6.90 (1H, d, J=2.4 Hz), 7.73 (1H, d, J=6.0 Hz), 7.83 (1H, d, J=8.4 Hz).

5

Example 6

5-(2-(3-((1S)-1-Carbamoyl-2-phenylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0186] N1-Methyl-5-((2-amino-4-pyridyl)oxy)-1H-1-indolcarboxamide (100 mg, 0.354 mmol) synthesized in Production example 5-1 and triethylamine (0.3 ml) were dissolved in N,N-dimethylformamide (3 ml). Phenyl chlorocarbonate (0.0888 ml, 0.708 mmol) was added dropwise thereto at room temperature and the reaction mixture was stirred for 30 minutes. (2S)-2-Amino-3-phenylpropionamide (290 mg, 1.77 mmol) was added and the reaction mixture was stirred for 3 days. The reaction mixture was partitioned between a solvent mixture of ethyl acetate-tetrahydrofuran and water. The organic layer was washed with water and brine, dried over anhydrous magnesium sulfate. The solvent was distilled off, and the residue was purified by silica gel column chromatography (eluent; ethyl acetate: methanol = 20: 1). The crystals were precipitated from a solvent mixture of ethyl acetate-hexane, filtered off, and dried under aeration to yield

the title compound (69.4 mg, 0.147 mmol, 41%) as white crystals.

Example 7

5 5-(2-(3-(2-Oxo-2-(pyrrolidin-1-yl)ethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0187] tert-Butoxycarbonylaminoacetic acid (876 mg, 5.00 mmol) and N-methylmorpholine (506 mg, 5.00 mmol) were dissolved in tetrahydrofuran (20 ml). After
10 isobutyl chloroformate (683 mg, 5.00 mmol) was added dropwise at below -15 °C and the reaction mixture was stirred for 30 minutes, pyrrolidine (782 mg, 11.0 mmol) was added at below -15 °C and the reaction mixture was further stirred at 0 °C for 30 minutes. The reaction
15 mixture was partitioned between ethyl acetate and 1N aqueous solution of sodium hydroxide. The organic layer was washed with 1N hydrochloric acid, a saturated aqueous solution of sodium hydrogencarbonate and brine, and was dried over anhydrous magnesium sulfate. The
20 solvent was distilled off, and the obtained residue was dissolved in a solvent mixture of ethyl acetate (10 ml)-tetrahydrofuran (5 ml). 4N hydrochloric acid Ethyl acetate solution (5 ml) was added and the reaction mixture was stirred at room temperature for 18 hours.
25 After the solvent was distilled off, ethyl acetate was added to the crude product to precipitate crystals; and

the crystals were filtered off and dried under aeration to yield 2-amino-1-(pyrrolidin-1-yl)ethanone hydrochloride (573 mg, 4.16 mmol, 84%) as white crystals.

5 The title compound (74.7 mg, 0.171 mmol, 86%) was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxy carbonyl)carbamate (104 mg, 0.200 mmol) synthesized in Production example 5-2 and the
10 previously obtained 2-amino-1-(pyrrolidin-1-yl)ethanone hydrochloride (165 mg, 1.00 mmol) similarly to Example 5.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.71-1.81 (2H, m), 1.83-1.93 (2H, m), 2.85 (3H, d, J=4.0 Hz), 3.26-3.40
15 (4H, m), 3.90 (2H, d, J=4.4 Hz), 6.55 (1H, dd, J=2.0, 6.0 Hz), 6.69 (1H, d, J=3.4 Hz), 6.94 (1H, d, J=2.0 Hz), 7.06 (1H, dd, J=2.0, 9.0 Hz), 7.38 (1H, d, J=2.0 Hz), 7.89 (1H, d, J=3.4 Hz), 8.05 (1H, d, J=6.0 Hz), 8.12-8.26 (2H, m), 8.30 (1H, d, J=9.0 Hz), 9.28 (1H, s).

20 [0188]

Example 8

5-(2-(3-(2-(4-Hydroxy-4-methylpiperidin-1-yl)-2-oxoethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

25 4-Hydroxy-4-methylpiperidine hydrochloride (113 mg, 0.745 mmol) was suspended in N,N-dimethylformamide

(3 ml), then triethylamine (1 ml) was added; benzotriazole-1-isooxytris(dimethylamino)phosphonium hexafluorophosphate (201 mg, 0.454 mmol) and ((4-(1-methylcarbamoyl-1H-indol-5-yloxy)pyridin-2-yl)aminocarbonylamino)acetic acid (145 mg, 0.378 mmol) were added thereto; and the reaction mixture was stirred at room temperature for 2 hours. After water was added to the reaction mixture, extraction was performed with ethyl acetate-tetrahydrofuran. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Fuji Silysia NH silica gel; ethyl acetate, ethyl acetate: methanol = 20: 1, 10: 1 in this order). After concentration under reduced pressure, the product was solidified with diethyl ether, suspended, filtered off, washed with diethyl ether, and dried under aeration to yield the title compound (137 mg, 0.285 mmol, 75.4%) as a colorless amorphous solid.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.10 (3H, s), 1.38-1.44 (4H, m), 2.83 (3H, d, J=3.6 Hz), 3.02 (2H, m), 3.90 (2H, m), 3.96 (2H, d, J=4.0 Hz), 4.37 (1H, s), 6.52 (1H, d, J=5.6 Hz), 6.67 (1H, d, J=3.2 Hz), 6.91 (1H, s), 7.04 (1H, d, J=9.0 Hz), 7.36 (1H, s), 7.87 (1H, d, J=3.2 Hz), 8.03 (1H, d, J=5.6 Hz), 8.17 (2H, m), 8.28 (1H, d, J=9.0 Hz), 9.27 (1H, s).

[0189] The starting materials were synthesized as follows.

Production example 8-1

5 ((4-(1-Methylcarbamoyl-1H-indol-5-yloxy)pyridin-2-yl)aminocarbonylamino)acetic acid

 Methyl aminoacetate hydrochloride (300 mg, 2.3 mmol) was dissolved in N,N-dimethylformamide (4 ml), and then triethylamine (1 ml) was added. Phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxy carbonyl) carbamate (250 mg, 0.48 mmol) synthesized in Production example 5-2 was added thereto. The reaction mixture was stirred at room temperature for 22 hours. After water was added to the reaction mixture, extraction was performed with a solvent mixture of ethyl acetate-tetrahydrofuran. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Fuji Silysia BW-300, ethyl acetate). The obtained pale yellow oil was dissolved in a solvent mixture of tetrahydrofuran (2 ml)-methanol (1 ml), then 4N aqueous solution of lithium hydroxide (0.48 ml) was added, and the reaction mixture was stirred at room temperature for 1 hour. After that, 1N hydrochloric acid (2 ml) was added, and this was

10
15
20
25

subjected to extraction with ethyl acetate-tetrahydrofuran. The organic layer was washed with brine and dried over anhydrous magnesium sulfate, and concentrated under reduced pressure to yield the title compound (145 mg, 0.38 mmol, 79%) as colorless crystals. ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.83 (3H, d, J=3.6 Hz), 3.81 (2H, d, J=5.6 Hz), 6.57 (1H, m), 6.68 (1H, d, J=3.6 Hz), 6.84 (1H, s), 7.05 (1H, dd, J=2.0, 9.2 Hz), 7.38 (1H, d, J=2.0 Hz), 7.88 (1H, d, J=3.6 Hz), 8.05 (1H, d, J=5.6 Hz), 8.16-8.30 (3H, m), 9.33 (1H, brs). [0190]

Production example 8-2

Benzyl (4-hydroxy-4-methylpiperidin-1-yl)carboxylate

Benzyl (4-oxopiperidin-1-yl)carboxylate (4.7 g, 20 mmol) was dissolved in tetrahydrofuran (200 ml); methyllithium-diethylether solution (9.0 ml (1.02 M) + 11.6 ml (1.14 M), total 22 mmol) was added dropwise thereto (internal temperature: -60 °C or below) while stirred at -78 °C under nitrogen atmosphere; and then the reaction mixture was stirred for 1.5 hours as it stands. On the other hand, a similar reaction was performed by using piperidin-4-one-1-carboxylate (1.1 g, 5.0 mmol) in another container. After the saturated aqueous solution of ammonium chloride was added to each reaction mixture, the two reaction mixtures were mixed. Extraction was performed with ethyl acetate, washed

with brine, dried over anhydrous magnesium sulfate, concentrated under reduced pressure, and purified by silica gel column chromatography (Fuji Silysia BW-300, hexane-ethyl acetate system) to yield the title compound (4.5 g, 18 mmol, 73%) as colorless crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.10 (3H, s), 1.32-1.44 (4H, m), 3.17 (2H, m), 3.61 (2H, dt, J=3.6, 9.2 Hz), 4.34 (1H, s), 5.04 (2H, s), 7.27-7.37 (5H, m).

[0191]

10 Production example 8-3

4-Hydroxy-4-methylpiperidine monohydrochloride

Benzyl (4-hydroxy-4-methylpiperidin-1-yl)carboxylate (4.5 g, 18 mmol) was dissolved in methanol (90 ml), 10% palladium on carbon powder (0.60 g) was added, and the reaction mixture was stirred at room temperature under hydrogen atmosphere overnight. The catalyst was removed by filtration and the resultant solution was concentrated under reduced pressure to yield a crude product of 4-hydroxy-4-methylpiperidine as a pale yellow oil (2.1 g). After the product was dissolved in methanol, 1N hydrochloric acid (17.5 ml) was added and the solvent was distilled off under reduced pressure. The obtained crystals were suspended in acetone, the crystals were filtered off, washed with acetone, and dried under aeration to yield the title compound (2.1 g, 14 mmol, 77%) as colorless

crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.14 (3H, s), 1.55-1.69 (4H, m), 3.00 (4H, m), 4.68 (1H, brs), 8.77 (1H, brs), 8.89 (1H, brs).

5

Example 9

5-(2-(3-((1S)-1-Carbamoylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0192] N1-Methyl-5-((2-amino-4-pyridyl)oxy)-1H-1-indolecarboxamide (100 mg, 0.354 mmol) synthesized in Production example 5-1 and triethylamine (1 ml) were dissolved in tetrahydrofuran (3 ml), then phenyl chlorocarbonate (0.0888 ml, 0.708 mmol) was added dropwise at room temperature, and the reaction mixture was stirred for 2 hours. After the solvent was distilled off under reduced pressure, the residue was dissolved in N,N-dimethylformamide (3 ml). (2S)-2-Aminopropionamide hydrochloride (220 mg, 1.77 mmol) and triethylamine (1 ml) were added and the reaction mixture was stirred for 18 hours. The reaction mixture was partitioned between ethyl acetate and a saturated aqueous solution of ammonium chloride. The organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was distilled off and the residue was purified by silica gel column chromatography (eluent; ethyl acetate: methanol = 20:

10

15

20

25

1). The crystals were precipitated from a solvent mixture of ethyl acetate-hexane, filtered off, and dried under aeration to yield the title compound (38.5 mg, 0.0971 mmol, 27%) as white crystals.

5 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.21 (3H, d, J=6.8 Hz), 2.85 (3H, d, J=4.0 Hz), 4.17 (1H, m), 6.55 (1H, d, J=5.2 Hz), 6.70 (1H, d, J=3.6 Hz), 6.93 (1H, s), 7.02 (1H, s), 7.06 (1H, dd, J=2.0, 8.8 Hz), 7.39 (1H, d, J=2.0 Hz), 7.46 (1H, s), 7.90 (1H, d, J=3.6 Hz), 8.06
10 (1H, d, J=5.2 Hz) 8.11 (1H, brs), 8.20 (1H, q, J=4.0 Hz), 8.30 (1H, d, J=8.8 Hz), 9.21 (1H, brs).

Example 10

15 5-(2-(3-((1S)-1-Carbamoyl-3-methylbutyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0193] Similarly to Example 9, the title compound (59.5 mg, 0.135 mmol, 38%) was obtained as white crystals from N1-methyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (100 mg, 0.354 mmol) synthesized in
20 Production example 5-1 and (2S)-2-amino-4-methylpentanamide hydrochloride (295 mg, 1.77 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.83-0.91 (6H, m), 1.35-1.50 (2H, m), 1.58 (1H, m), 2.85 (3H, d, J=4.4 Hz), 4.17 (1H, m), 6.53 (1H, dd, J=2.4, 6.0 Hz), 6.69 (1H, d, J=3.8 Hz), 6.92-7.01 (2H, m), 7.06 (1H, dd, J=2.4, 8.8
25 Hz), 7.38 (1H, d, J=2.4 Hz), 7.48 (1H, s), 7.89 (1H, d,

J=3.8 Hz) 7.98-8.12 (2H, m), 8.19 (1H, q, J=4.4 Hz), 8.30 (1H, d, J=8.8 Hz), 9.09 (1H, s).

Example 11

5 5-(2-(3-Carbamoylmethylureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0194] Similarly to Example 5, the title compound (52.8 mg, 0.138 mmol, 69%) was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (104 mg, 0.200 mmol) synthesized in Production example 5-2 and glycine hydrochloride (111 mg, 1.00 mmol).
1H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.85 (3H, d, J=4.0 Hz), 3.70 (2H, d, J=5.2 Hz), 6.53 (1H, dd, J=2.4, 5.8 Hz), 6.69 (1H, d, J=3.4 Hz), 6.92 (1H, d, J=2.4 Hz), 7.01 (1H, s), 7.06 (1H, dd, J=2.4, 9.2 Hz), 7.34-7.42 (2H, m), 7.89 (1H, d, J=3.4 Hz), 8.05 (1H, d, J=5.8 Hz), 8.14-8.26 (2H, m), 8.30 (1H, d, J=9.2 Hz), 9.21 (1H, s).

20 Example 12

5-(2-(3-Cyclopropylcarbamoylmethylureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0195] Similarly to Example 5, the title compound (50.7 mg, 0.120 mmol, 60%) was obtained as white powder from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (104 mg, 0.200

mmol) synthesized in Production example 5-2 and 2-amino-N-cyclopropylacetamide hydrochloride (151 mg, 1.00 mmol) obtained from tert-butoxycarbonylaminoacetic acid and cyclopropylamine by the method similar to Example 7.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.36-0.42 (2H, m), 0.57-0.63 (2H, m), 2.60 (1H, m), 2.85 (3H, d, J=4.4 Hz), 3.68 (2H, d, J=5.2 Hz), 6.53 (1H, dd, J=2.0, 6.0 Hz), 6.69 (1H, d, J=3.6 Hz), 6.91 (1H, d, J=2.0 Hz), 7.06 (1H, dd, J=2.4, 9.0 Hz), 7.38 (1H, d, J=2.4 Hz), 7.89 (1H, d, J=3.6 Hz), 8.00 (1H, d, J=4.0 Hz), 8.06 (1H, d, J=6.0 Hz) 8.14-8.26 (2H, m), 8.30 (1H, d, J=9.0 Hz), 9.21 (1H, s).

Example 13

5-(2-(3-((1S)-1-Carbamoyl-2-hydroxyethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0196] Similarly to Example 9, the title compound (52.1 mg, 0.126 mmol, 36%) was obtained as white crystals from N1-methyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (100 mg, 0.354 mmol) synthesized in Production example 5-1 and (2S)-2-amino-3-hydroxypropionamide hydrochloride (249 mg, 1.77 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.85 (3H, d, J=4.4 Hz), 3.52 (1H, dd, J=4.8, 6.4 Hz), 3.62 (1H, dd, J=4.8,

6.4 Hz), 4.13 (1H, m), 4.94 (1H, brs), 6.53 (1H, dd, J=2.4, 6.0 Hz), 6.69 (1H, d, J=3.6 Hz), 6.99 (1H, s), 7.02-7.10 (2H, m), 7.35 (1H, s), 7.38 (1H, d, J=2.4 Hz), 7.89 (1H, d, J=3.6 Hz), 8.05 (1H, d, J=6.0 Hz), 8.10-
 5 8.26 (2H, m), 8.30 (1H, d, J=8.8 Hz), 9.22 (1H, s).

Example 14

5-(2-(3-((1R)-1-Carbamoyl-2-hydroxyethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0197] Similarly to Example 9, the title compound (56.0 mg, 0.136 mmol, 68%) was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate
 15 (104 mg, 0.200 mmol) synthesized in Production example 5-2 and (2R)-2-amino-3-hydroxypropioamide hydrochloride (167 mg, 1.00 mmol) obtained from (2R)-2-(tert-butoxycarbonylamino)-3-hydroxypropionic acid and aqueous ammonia by the method similar to Example 7.

20

Example 15

(2S)-2-(3-(4-(1-Methylcarbamoyl-1H-indol-5-yloxy)pyridin-2-yl)ureido)-1,5-pentanedicarboxylic acid diamide

[0198] Similarly to Example 6, the title compound (82.5 mg, 0.189 mmol, 51%) was obtained as white powder from

N1-methyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (100 mg, 0.354 mmol) synthesized in Production example 5-1 and (2S)-2-amino-1,5-pentanedicarboxylic acid diamide hydrochloride (321 mg, 1.77 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.66-2.28 (4H, m), 2.85 (3H, d, J=4.4 Hz), 4.17 (1H, m), 6.53 (1H, dd, J=2.4, 6.0 Hz), 6.69 (1H, d, J=3.6 Hz), 6.72 (1H, s), 6.97 (1H, s), 7.01-7.10 (2H, m), 7.30 (1H, s), 7.38 (1H, d, J=2.4 Hz), 7.49 (1H, s), 7.76 (1H, s), 7.89 (1H, d, J=3.6 Hz), 8.06 (1H, d, J=6.0 Hz), 8.18 (1H, q, J=4.4 Hz), 8.30 (1H, d, J=8.8 Hz), 9.13 (1H, s).

Example 16

(2S)-2-(3-(4-(1-Methylcarbamoyl-1H-indol-5-yloxy)pyridin-2-yl)ureido)succinamide

[0199] Similarly to Example 6, the title compound (65.7 mg, 0.150 mmol, 42%) was obtained as white crystals from N1-methyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (100 mg, 0.354 mmol) synthesized in Production example 5-1 and (2S)-2-aminosuccinamide hydrochloride (297 mg, 1.77 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.45 (2H, d, J=6.8 Hz), 2.85 (3H, d, J=3.6 Hz), 4.40 (1H, m), 6.53 (1H, dd, J=2.4, 6.0 Hz), 6.69 (1H, d, J=3.6 Hz), 6.88 (1H, s), 6.95 (1H, s), 7.00 (1H, d, J=2.4 Hz), 7.06 (1H, dd,

J=2.4, 9.2 Hz), 7.28 (1H, s), 7.35 (1H, s), 7.38 (1H, d, J=2.4 Hz), 7.89 (1H, d, J=3.6 Hz), 8.04 (1H, d, J=6.0 Hz), 8.18 (1H, q, J=4.0 Hz), 8.26 (1H, brs), 8.30 (1H, d, J=9.2 Hz), 9.19 (1H, s).

5

Example 17

5-(2-(3-((1S)-1-Cyclopropylcarbamoyl-2-hydroxyethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

10 [0200] Similarly to Example 5, the title compound (72.0 mg, 0.159 mmol, 80%) was obtained as white powder from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (104 mg, 0.200 mmol) synthesized in Production example 5-2 and (2S)-2-amino-N-cyclopropyl-3-hydroxypropionamide hydrochloride
15 (181 mg, 1.00 mmol) obtained from (2S)-2-(tert-butoxycarbonylamino)-3-hydroxypropionic acid and cyclopropylamine by the method similar to Example 7.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.35-0.44 (2H, m),
20 0.54-0.63 (2H, m), 2.62 (1H, m), 2.85 (3H, d, J=4.0 Hz), 3.45-3.58 (2H, m), 4.09 (1H, m), 4.91 (1H, t, J=5.2 Hz), 6.53 (1H, dd, J=2.0, 6.0 Hz), 6.69 (1H, d, J=3.6 Hz), 6.99 (1H, d, J=2.0 Hz), 7.04 (1H, dd, J=2.4, 8.8 Hz), 7.38 (1H, d, J=2.4 Hz), 7.89 (1H, d, J=3.6 Hz), 7.98
25 (1H, d, J=4.4 Hz), 8.05 (1H, d, J=6.0 Hz), 8.09-8.24 (2H, m), 8.30 (1H, d, J=8.8 Hz), 9.18 (1H, s).

Example 18

5-(2-(3-((1S)-1-Hydroxymethyl-2-oxo-2-pyrrolidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0201] Similarly to Example 5, the title compound (67.6 mg, 0.145 mmol, 73%) was obtained as white powder from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (104 mg, 0.200 mmol) synthesized in Production example 5-2 and (2S)-2-amino-3-hydroxy-1-(pyrrolidin-1-yl)propan-1-one hydrochloride (165 mg, 0.848 mmol) obtained from (2S)-2-(tert-butoxycarbonylamino)-3-hydroxypropionic acid and pyrrolidine by the method similar to Example 7.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.72-1.81 (2H, m), 1.81-1.90 (2H, m), 2.85 (3H, d, J=4.4 Hz), 3.22-3.36 (2H, m), 3.46-3.60 (4H, m), 4.54 (1H, m), 4.98 (1H, brs), 6.54 (1H, dd, J=2.0, 5.6 Hz), 6.69 (1H, d, J=3.6 Hz), 6.97 (1H, d, J=2.0 Hz), 7.05 (1H, dd, J=2.4, 8.8 Hz), 7.38 (1H, d, J=2.4 Hz), 7.89 (1H, d, J=3.6 Hz), 8.05 (1H, d, J=5.6 Hz), 8.13-8.23 (2H, m), 8.30 (1H, d, J=8.8 Hz), 9.18 (1H, s).

Example 19

5-(2-(3-((1R)-1-Hydroxymethyl-2-oxo-2-pyrrolidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic

acid methylamide

[0202] Similarly to Example 5, the title compound (305 mg, 0.654 mmol, 93%) was obtained as white powder from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (366 mg, 0.700 mmol) synthesized in Production example 5-2 and (2R)-2-amino-3-hydroxy-1-(pyrrolidin-1-yl)propan-1-one hydrochloride obtained from (2R)-2-(tert-butoxycarbonylamino)-3-hydroxypropionic acid and pyrrolidine by the method similar to Example 7.

Example 20

5-(2-(3-((1S)-1-Hydroxymethyl-2-oxo-2-piperidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0203] Similarly to Example 5, the title compound (124 mg, 0.258 mmol, 86%) was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (157 mg, 0.300 mmol) synthesized in Production example 5-2 and (2S)-2-amino-3-hydroxy-1-(piperidin-1-yl)propan-1-one hydrochloride (312 mg, 1.50 mmol) obtained from (2S)-2-(tert-butoxycarbonylamino)-3-hydroxypropionic acid and piperidine by the method similar to Example 7.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.36-1.61 (6H, m),

2.85 (3H, d, J=4.4 Hz), 3.40-3.53 (6H, m), 4.76 (1H, m),
4.92 (1H, brs), 6.54 (1H, dd, J=2.4, 6.0 Hz), 6.69 (1H,
d, J=3.6 Hz), 6.97 (1H, d, J=2.4 Hz), 7.06 (1H, dd,
J=2.4, 9.0 Hz), 7.38 (1H, d, J=2.4 Hz), 7.89 (1H, d,
5 J=3.6 Hz), 8.05 (1H, d, J=6.0 Hz), 8.10-8.26 (2H, m),
8.30 (1H, d, J=9.0 Hz), 9.21 (1H, s).

Example 21

10 5-(2-(3-((1R)-1-Hydroxymethyl-2-oxo-2-piperidin-1-
ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic
acid methylamide

[0204] (2R)-2-Benzylloxycarbonylamino-3-hydroxypropionic
acid (1.91 g, 8.00 mmol) and N-methylmorpholine (809 mg,
8.00 mmol) were dissolved in tetrahydrofuran (20 ml).
15 After isobutyl chloroformate (1.09 g, 8.00 mmol) was
added dropwise at -15 °C or below, the reaction mixture
was stirred for 30 minutes. Then, pyrrolidine (1.13 g,
16.0 mmol) was added at -15 °C or below, and the
reaction mixture was further stirred at 0 °C for 30
20 minutes. The reaction mixture was partitioned between
ethyl acetate and water. The organic layer was washed
with 1N hydrochloric acid, 1N aqueous solution of
sodium hydroxide, a saturated aqueous solution of
sodium hydrogencarbonate, and brine, and dried over
25 anhydrous magnesium sulfate. The solvent was distilled
off, and the obtained residue was dissolved in a

solvent mixture of methanol (15 ml)-tetrahydrofuran (15 ml). Then, 10% palladium on carbon (wet) (300 mg) was added, and the reaction mixture was stirred at room temperature under the stream of hydrogen for 90 minutes.

5 After the catalyst was removed by filtration, the solvent of the filtrate was distilled off under reduced pressure to yield (2R)-2-amino-3-hydroxy-1-(piperidin-1-yl)propan-1-one (684 mg, 3.97 mmol, 50%) as a colorless oil. Similarly to Example 5, the title

10 compound (107 mg, 0.223 mmol, 74%) was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (157 mg, 0.300 mmol) synthesized in Production example 5-2 and previously

15 obtained (2R)-2-amino-3-hydroxy-1-(piperidin-1-yl)propan-1-one (228 mg, 1.32 mmol).

Example 22

20 5-(2-(3-((1S)-1-Hydroxymethyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0205] Similarly to Example 5, the title compound (118 mg, 0.238 mmol, 69%) was obtained as white powder from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (179 mg, 0.343

25 mmol) synthesized in Production example 5-2 and (2S)-2-

amino-3-hydroxy-1-(4-hydroxypiperidin-1-yl)propan-1-one hydrochloride (385 mg, 1.71 mmol) obtained from (2S)-2-(tert-butoxycarbonylamino)-3-hydroxypropionic acid and 4-hydroxypiperidine by the method similar to Example 7.

5 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.16-1.40 (2H, m), 1.61-1.80 (2H, m), 2.85 (3H, d, J=4.0 Hz), 2.98-3.50 (5H, m), 3.63-3.95 (3H, m), 4.76 (1H, m), 4.92 (1H, brs), 6.55 (1H, dd, J=2.0, 6.0 Hz), 6.69 (1H, d, J=3.6 Hz), 6.96 (1H, d, J=2.0 Hz), 7.06 (1H, dd, J=2.4, 8.8 Hz), 7.38 (1H, d, J=2.4 Hz), 7.90 (1H, d, J=3.6 Hz), 8.05 (1H, d, J=6.0 Hz), 8.08-8.26 (2H, m), 8.30 (1H, d, J=8.8 Hz), 9.26 (1H, s).

Example 23

15 5-(2-(3-((1S)-1-Hydroxymethyl-2-(morpholin-4-yl)-2-oxoethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0206] Similarly to Example 5, the title compound (121 mg, 0.251 mmol, 84%) was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (157 mg, 0.300 mmol) synthesized in Production example 5-2 and (2S)-2-amino-3-hydroxy-1-(morpholin-4-yl)propan-1-one hydrochloride (316 mg, 1.50 mmol) obtained from (2S)-2-(tert-butoxycarbonylamino)-3-hydroxypropionic acid and morpholine by the method

similar to Example 7.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.85 (3H, d, J=4.4 Hz), 3.36-3.62 (10H, m), 4.74 (1H, m), 4.92 (1H, brs), 6.54 (1H, dd, J=2.4, 6.0 Hz), 6.69 (1H, d, J=3.6 Hz), 6.96 (1H, d, J=2.4 Hz), 7.06 (1H, dd, J=2.4, 8.8 Hz), 7.38 (1H, d, J=2.4 Hz), 7.89 (1H, d, J=3.6 Hz), 8.05 (1H, d, J=6.0 Hz), 8.14-8.28 (2H, m), 8.30 (1H, d, J=8.8 Hz), 9.25 (1H, s).

10 Example 24

5-(2-(3-(2-Cyclopropylcarbamoyl)ethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0207] Similarly to Example 5, the title compound (117 mg, 0.268 mmol, 89%) was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (157 mg, 0.300 mmol) synthesized in Production example 5-2 and 3-amino-N-cyclopropylpropionamide hydrochloride (247 mg, 1.50 mmol) obtained from 3-(tert-butoxycarbonylamino)propionic acid and cyclopropylamine by the method similar to Example 7.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.32-0.38 (2H, m), 0.54-0.60 (2H, m), 2.19 (2H, t, J=6.4 Hz), 2.60 (1H, m), 2.85 (3H, d, J=4.4 Hz), 3.25-3.33 (2H, m), 6.53 (1H, dd, J=2.0, 6.0 Hz), 6.69 (1H, d, J=3.6 Hz), 6.90 (1H, d, J=2.0 Hz), 7.05 (1H, dd, J=2.4, 9.0 Hz), 7.38 (1H, d,

J=2.4 Hz), 7.89 (1H, d, J=3.6 Hz), 7.93 (1H, d, J=4.0 Hz), 7.96-8.06 (2H, m), 8.18 (1H, q, J=4.4 Hz), 8.30 (1H, d, J=9.0 Hz), 9.08 (1H, s).

5 Example 25

5-(2-(3-(3-oxo-3-(pyrrolidin-1-yl)propyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0208] Similarly to Example 5, the title compound (122 mg, 0.270 mmol, 90%) was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (157 mg, 0.300 mmol) synthesized in Production example 5-2 and 3-amino-1-(pyrrolidin-1-yl)propan-1-one hydrochloride (268 mg, 1.50 mmol) obtained from 3-(tert-butoxycarbonylamino)propionic acid and pyrrolidine by the same method similar to Example 7.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.70-1.78 (2H, m), 1.80-1.88 (2H, m), 2.40 (2H, t, J=6.2 Hz), 2.85 (3H, d, J=4.4 Hz), 3.24-3.38 (6H, m), 6.52 (1H, dd, J=2.0, 5.6 Hz), 6.69 (1H, d, J=3.6 Hz), 6.92 (1H, d, J=2.0 Hz), 7.05 (1H, dd, J=2.4, 9.0 Hz), 7.38 (1H, d, J=2.4 Hz), 7.89 (1H, d, J=3.6 Hz), 7.98-8.10 (2H, m), 8.18 (1H, q, J=4.4 Hz), 8.30 (1H, d, J=9.0 Hz), 9.10 (1H, s).

25

Example 26

5-(2-(3-(3-(4-Hydroxy-4-methylpiperidin-1-yl)-3-oxopropyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0209] The title compound (177 mg, 0.358 mmol, 71.1%)
5 was obtained as colorless crystals by performing the
reaction similar to Example 8 using 3-(3-(4-(1-methylcarbamoyl-1H-indol-5-yloxy)pyridin-2-yl)ureido)propionic acid (200 mg, 0.503 mmol) and 4-hydroxy-4-methylpiperidine monohydrochloride (114 mg,
10 0.755 mmol, Production example 8-3).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.07 (3H, s), 1.23-
1.41 (4H, m), 2.44 (2H, d, J=4.8 Hz), 2.83 (3H, d, J=4.4 Hz), 2.98 (1H, m), 3.23-3.30 (3H, m), 3.46 (1H, m), 3.93 (1H, m), 4.32 (1H, s), 6.49 (1H, dd, J=2.0,
15 6.0 Hz), 6.67 (1H, d, J=3.4 Hz), 6.90 (1H, s), 7.03 (1H, dd, J=2.0, 8.8 Hz), 7.35 (1H, d, J=2.0 Hz), 7.87 (1H, d, J=3.4 Hz), 8.00 (2H, m), 8.15 (1H, d, J=4.4 Hz), 8.28 (1H, d, J=8.8 Hz), 9.06 (1H, s).

20 [0210] The starting material was synthesized by the following methods.

Production example 26-1

3-(3-(4-(1-Methylcarbamoyl-1H-indol-5-yloxy)pyridin-2-yl)ureido)propionic acid

25 Ethyl 4-aminopropionate hydrochloride (588 mg, 3.8 mmol) was suspended in N,N-dimethylformamide (3.0

ml), and then 5N aqueous solution of sodium hydroxide (0.77 ml, 3.8 mmol) was added, and the reaction mixture was stirred at room temperature. Phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (400 mg, 0.77 mmol, Production example 5-2) was added thereto, and the reaction mixture was stirred at room temperature for 0.75 hours. Water was added to the reaction mixture, and this was subjected to extraction with ethyl acetate-tetrahydrofuran, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Fuji Silysia BW-300, ethyl acetate) to yield a pale brown oil. This oil was dissolved in tetrahydrofuran (4.0 ml) and methanol (2.0 ml), 4N aqueous solution of lithium hydroxide (0.77 ml) was added at room temperature, and the reaction mixture was stirred at room temperature for 1.5 hours. To the reaction mixture, 1N hydrochloric acid (3.1 ml) was added while stirred at room temperature; and this was subjected to extraction with ethyl acetate-tetrahydrofuran, washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. A small amount of acetone was added to the obtained amorphous solid, and this solution was diluted with diethyl ether. The crystals were filtered

off, washed with diethyl ether, and dried under aeration to yield the title compound (200 mg, 0.50 mmol, 66%) as colorless crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.39 (2H, t, J=6.2 Hz), 2.84 (3H, d, J=4.0 Hz), 3.30 (2H, m), 6.51 (1H, d, J=5.8 Hz), 6.68 (1H, d, J=3.2 Hz), 6.87 (1H, s), 7.05 (1H, d, J=9.0 Hz), 7.37 (1H, s), 7.88 (1H, d, J=3.2 Hz), 8.01 (1H, d, J=5.8 Hz), 8.16 (1H, m), 8.17 (1H, d, J=4.0 Hz), 8.29 (1H, d, J=9.0 Hz), 9.10 (1H, s), 12.24 (1H, s).

Example 27

N1-Ethyl-5-(2-(((2-ethoxyethyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0211] Phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate (100 mg, 0.24 mmol) was dissolved in N,N-dimethylformamide (1.0 ml), and 2-ethoxyethylamine (0.063 ml, 0.6 mmol) was added while stirred at room temperature. After 1 hour, the reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with water and brine, and dried over anhydrous sodium sulfate. After the solvent was distilled off, the crystals were precipitated from ethyl acetate-hexane (1: 5), filtered off, and dried under aeration to yield the title compound (100 mg, 0.24 mmol, quantitative) as white

crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.09 (3H, t, J=7.2 Hz), 1.17 (3H, t, J=7.2 Hz), 3.21-3.45 (8H, m), 6.50 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 6.87 (1H, brs), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, d, J=2.4 Hz), 7.91 (1H, d, J=3.6 Hz), 8.01 (1H, d, J=5.6 Hz), 8.12 (1H, m), 8.22 (1H, t, J=4.8 Hz), 8.28 (1H, d, J=8.8 Hz), 9.08 (1H, s).

[0212] The starting materials were synthesized by the following methods.

Production example 27-1

N1-Ethyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide

Sodium hydride (573 mg, 14.32 mmol) was suspended in N,N-dimethylformamide (30 ml) under nitrogen atmosphere. 4-(1H-5-Indolyloxy)-2-pyridinamine (3.00 g, 13.32 mmol, CAS No. 417722-11-3) described in WO 02/32872 was gradually added thereto while stirred at room temperature. After 10 minutes, the reaction mixture was cooled with an ice water bath, and phenyl N-ethylcarbamate (2.31 g, 13.98 mmol) was added. The reaction mixture was heated to room temperature and was stirred for 2 hours. The reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with water and brine, and

dried over anhydrous sodium sulfate. The solvent was distilled off, then the crystals were precipitated from ethyl acetate, filtered off, and dried under aeration to yield the title compound (3.168 g, 10.69 mmol, 80.3%) as pale brown crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.32 (3H, t, J=7.2 Hz), 2.40-2.50 (2H, m), 5.74 (1H, d, J=2.4 Hz), 5.83 (2H, brs), 6.12 (1H, dd, J=2.4, 5.6 Hz), 6.66 (1H, d, J=3.6 Hz), 7.01 (1H, dd, J=2.4, 8.8 Hz), 7.32 (1H, d, J=2.4 Hz), 7.75 (1H, d, J=5.6 Hz), 7.88 (1H, d, J=3.6 Hz), 8.19 (1H, t, J=5.6 Hz), 8.26 (1H, d, J=8.8 Hz).

Production example 27-2

Phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate

[0213] N1-ethyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (3.168 g, 10.69 mmol) synthesized in Production example 27-1 was dissolved in N,N-dimethylformamide (30 ml) under nitrogen atmosphere. Pyridine (1.25 ml, 15.40 mmol) and phenyl chlorocarbonate (1.61 ml, 12.83 mmol) were sequentially added dropwise while cooled with an ice water bath. The reaction mixture was heated to room temperature while stirred. After 1 hour, the reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with water and brine, and

dried over anhydrous sodium sulfate. The solvent was distilled off, and the crystals were precipitated from ethyl acetate, filtered off, and dried under aeration to yield the title compound (1.530 g, 3.67 mmol, 34.4%) as white crystals.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 1.32 (3H, t, J=7.2 Hz), 3.53 (2H, m), 5.48 (1H, m), 6.58 (1H, d, J=4.0 Hz), 6.62 (1H, dd, J=2.4, 5.6 Hz), 7.06 (1H, dd, J=2.4, 8.8 Hz), 7.15 (2H, m), 7.20-7.27 (1H, m), 7.30 (1H, d, J=2.4 Hz), 7.37 (2H, m), 7.45 (1H, d, J=4.0 Hz), 7.52 (1H, d, J=2.4 Hz), 8.10-8.15 (3H, m).

Example 28

N1-Methyl-5-(2-((4-(2-hydroxy-2-methylpropionyl)piperazin-1-yl)carbonyl)amino-4-pyridyl)oxy-1H-indolecarboxamide

[0214] N1-Methyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (150 mg, 0.53 mmol) synthesized in Production example 5-1 was dissolved in tetrahydrofuran (3 ml). Triethylamine (0.37 ml, 2.66 mmol) and phenyl chlorocarbonate (0.15 ml, 1.2 mmol) were sequentially added dropwise at room temperature, and the reaction mixture was stirred for 30 minutes. 1-(2-Hydroxy-2-methylpropionyl)piperazine (412 mg, 2.39 mmol) and N,N-dimethylformamide (3 ml) were added and the reaction mixture was stirred for 3 days. The reaction mixture

was partitioned between ethyl acetate and water. The organic layer was washed with water and brine, and dried over anhydrous sodium sulfate. The solvent was distilled off, and the residue was purified by silica gel column chromatography (eluent; ethyl acetate: methanol = 95: 5). The crystals were precipitated from diethyl ether-hexane (1 : 2), filtered off, and dried under aeration to yield the title compound (189.4 mg, 0.39 mmol, 74.2%) as white crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.28 (6H, s), 2.83 (3H, d, J=4.0 Hz), 3.10-3.50 (8H, m), 5.43 (1H, s), 6.56 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.30 (1H, d, J=2.4 Hz), 7.36 (1H, d, J=2.4 Hz), 7.87 (1H, d, J=3.6 Hz), 8.08 (1H, d, J=5.6 Hz), 8.16 (1H, q, J=4.0 Hz), 8.28 (1H, d, J=8.8 Hz), 9.21 (1H, s).

[0215] 1-(2-Hydroxy-2-methylpropionyl)piperazine was synthesized by the following methods.

20 Production example 28-1

Benzyl 4-(2-hydroxy-2-methylpropionyl)piperazine-1-carboxylate

Benzyl piperazine-1-carbamate (2.203 g, 10.0 mmol) was dissolved in tetrahydrofuran (50 ml); 2-hydroxy-2-methylpropionic acid (1.25 g, 12.0 mmol), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide

hydrochloride (2.30 g, 12.0 mmol), 1-hydroxy-1H-benzotriazole monohydrate (1.84 g, 12.0 mmol) and triethylamine (3.35 ml, 24.0 mmol) were added; and the reaction mixture was stirred at room temperature for 7
5 hours. The reaction mixture was partitioned between ethyl acetate and 1N hydrochloric acid. The organic layer was washed with water, a saturated aqueous solution of sodium hydrogencarbonate and brine, and dried over anhydrous sodium sulfate. The solvent was
10 distilled off, and dried under reduced pressure to yield the title compound (2.823 g, 9.21 mmol, 92.1%) as a colorless oil.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 1.50 (6H, s), 3.52-3.55 (4H, m), 3.60-3.70 (4H, m), 3.93 (1H, s), 5.16 (2H, s),
15 7.34-7.38 (5H, m).

Production example 28-2

1-(2-Hydroxy-2-methylpropionyl)piperazine

[0216] Benzyl 4-(2-hydroxy-2-methylpropionyl)piperazine-1-carbamate (2.82 g, 9.20
20 mmol) synthesized in Production example 28-1 was dissolved in methanol (100 ml) under nitrogen atmosphere; 10% palladium on carbon (50% wet, 1.96 g) was added thereto, the reaction system was purged with
25 hydrogen at atmospheric pressure; and the reaction mixture was stirred overnight. After the reaction

system was purged with nitrogen, the catalyst was filtered out, and washed with methanol, then the solvent, together with the filtrate and the washing solution, was distilled off. The residue was dried under reduced pressure to yield the title compound (1.58 g, 9.20 mmol, quantitative) as a colorless oil.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 1.49 (6H, s), 2.84-2.94 (4H, m), 3.49 (1H, s), 3.62-3.70 (4H, m).

[0217]

10 Example 29

N1-Methyl-5-(2-((3-diethylamino)propylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

Similarly to Example 27, the title compound (96.4 mg, 0.22 mmol, 73.3%) was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate (121 mg, 0.30 mmol) and 3-(diethylamino)propylamine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.91 (6H, t, J=7.2 Hz), 1.50 (2H, m), 2.30-2.44 (6H, m), 2.83 (3H, d, J=4.4 Hz), 3.23 (2H, m), 6.50 (1H, dd, J=2.4, 6.0 Hz), 6.68 (1H, d, J=3.6 Hz), 6.82 (1H, s), 7.04 (1H, dd, J=2.4, 8.8 Hz), 7.37 (1H, d, J=2.4 Hz), 7.87 (1H, d, J=3.6 Hz), 8.01 (1H, d, J=6.0 Hz), 8.10-8.17 (2H, m), 8.29 (1H, d, J=8.8 Hz), 9.04 (1H, s).

[0218]The starting material was synthesized as follows.

Production example 29-1

Phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate

5 N1-Methyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (2.163 g, 7.66 mmol) synthesized in Production example 5-1 was dissolved in N,N-dimethylformamide (50 ml) under nitrogen atmosphere; pyridine (0.93 ml, 11.5 mmol), triethylamine (2.4 ml, 10 17.24 mmol) and phenyl chlorocarbonate (1.44 ml, 11.5 mmol) were sequentially added dropwise while cooled with an ice water bath; and the reaction mixture was heated to room temperature while stirred. After 1 hour, the reaction mixture was partitioned between ethyl 15 acetate and water. The organic layer was washed with water and brine, and dried over anhydrous sodium sulfate. The solvent was distilled off, and then the residue was purified by silica gel column chromatography (eluent; ethyl acetate), precipitated 20 from ethyl acetate-hexane (1: 10), filtered off, and dried under aeration to yield the title compound (2.731 g, 6.79 mmol, 88.6%) as white crystals.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 3.09 (3H, d, J=4.8 Hz), 5.52 (1H, m), 6.62 (1H, d, J=3.6 Hz), 6.98 (1H, dd, 25 J=2.4, 5.6 Hz), 7.01 (1H, d, J=2.4 Hz), 7.11 (1H, dd, J=2.4, 8.8 Hz), 7.14-7.40 (7H, m), 7.47 (1H, d, J=3.6

Hz), 8.24 (1H, d, J=8.8 Hz), 8.41 (1H, d, J=5.6 Hz).

Example 30

N1-Methyl-5-(2-(((3-4-

5 hydroxypiperidino)propyl)amino)carbonyl)amino-4-
pyridyl)oxy-1H-1-indolecarboxamide

[0219] Similarly to Example 27, the title compound
 (51.3 mg, 0.11 mmol, 29.5%) was obtained as white
 crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-
 10 5-indolyl)oxy-2-pyridyl)carbamate (150 mg, 0.37 mmol,
 Production example 29-1) and 1-(3-aminopropyl)-4-
 hydroxypiperidine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.29-1.38 (2H, m),
 1.50-1.55 (2H, m), 1.64-1.68 (2H, m), 1.88-1.92 (2H, m),
 15 2.20-2.24 (2H, m), 2.62-2.66 (2H, m), 2.83 (3H, d,
 J=4.4 Hz), 3.06-3.12 (2H, m), 3.39 (1H, m), 4.49 (1H, d,
 J=4.0 Hz), 6.50 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d,
 J=3.6 Hz), 6.84 (1H, s), 7.03 (1H, dd, J=2.4, 8.8 Hz),
 7.36 (1H, s), 7.87 (1H, d, J=3.6 Hz), 8.01 (1H, d,
 20 J=5.6 Hz), 8.05 (1H, m), 8.16 (1H, q, J=4.4 Hz), 8.28
 (1H, d, J=8.8 Hz), 9.02 (1H, s).

Example 31

N1-Methyl-5-(2-(((3-(4-methylpiperazin-1-

25 yl)propyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-
indolecarboxamide

[0220] Similarly to Example 27, the title compound (133.2 mg, 0.29 mmol, 76.8%) was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate (150 mg, 0.37 mmol, Production example 29-1) and 1-(3-aminopropyl)-4-methylpiperazine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.53 (2H, m), 2.11 (3H, s), 2.11-2.40 (10H, m), 2.83 (3H, d, J=4.0 Hz), 3.09 (2H, m), 6.50 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 6.84 (1H, s), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, d, J=2.4 Hz), 7.87 (1H, d, J=3.6 Hz), 8.01 (1H, d, J=5.6 Hz), 8.05 (1H, m), 8.16 (1H, q, J=4.0 Hz), 8.28 (1H, d, J=8.8 Hz), 9.01 (1H, s).

15 Example 32

5-(2-(3-(4-Oxo-4-(pyrrolidin-1-yl)butyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0221] The title compound (113 mg, 0.24 mmol, 77%) was obtained as colorless crystals by performing the reaction similar to Example 8 using 4-((4-(1-methylcarbamoyl-1H-indol-5-yloxy)pyridin-2-yl)aminocarbonylamino)butyric acid (130 mg, 0.31 mmol) and pyrrolidine (0.053 ml, 0.63 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.64 (2H, m), 1.71 (2H, m), 1.82 (2H, m), 2.20 (2H, t, J=6.8 Hz), 2.83 (3H, d, J=4.0 Hz), 3.09 (2H, q, J=6.8 Hz), 3.22 (2H, t,

J=6.8 Hz), 3.33 (2H, m), 6.50 (1H, dd, J=2.4, 5.8 Hz),
 6.67 (1H, d, J=3.6 Hz), 6.86 (1H, d, J=2.4 Hz), 7.03
 (1H, dd, J=2.4, 9.0 Hz), 7.36 (1H, d, J=2.4 Hz), 7.87
 (1H, d, J=3.6 Hz), 8.00 (1H, m), 8.03 (1H, d, J=5.8 Hz),
 5 8.16 (1H, m), 8.28 (1H, d, J=9.0 Hz), 9.00 (1H, s).

[0222] The starting material was synthesized by the following methods.

Production example 32-1

10 4-((4-(1-Methylcarbamoyl-1H-indol-5-yloxy)pyridin-2-yl)aminocarbonylamino)butyric acid

Ethyl 4-aminobutyrate hydrochloride (1.0 g, 6.0 mmol) was suspended in N,N-dimethylformamide (6.7 ml), 5N aqueous solution of sodium hydroxide (1.2 ml, 6.0
 15 mmol) was added and the reaction mixture was stirred at room temperature. Phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxy-carbonyl)carbamate (700 mg, 1.3 mmol, Production example 5-2) was added thereto and the
 20 reaction mixture was stirred at room temperature for 1.2 hours. The reaction mixture was partitioned between ethyl acetate and water. The organic layer was dried over anhydrous magnesium sulfate, concentrated under reduced pressure. The residue was purified by
 25 silica gel column chromatography (Fuji Silysia BW-300, ethyl acetate) to yield a pale yellow oil. This oil

was dissolved in tetrahydrofuran (6.0 ml) and methanol (3.0 ml); 4N lithium hydroxide (1.1 ml) was added thereto at room temperature; and the reaction mixture was stirred at room temperature for 3.5 hours.

5 Moreover, 1N hydrochloric acid (4.4 ml) and water (2 ml) were added thereto while stirred at room temperature; and this was subjected to extraction with ethyl acetate, washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced

10 pressure. After the precipitated crystals were suspended in diethyl ether: hexane = 1: 1, the crystals were filtered off, washed with diethyl ether, and dried under aeration to yield the title compound (411 mg, 1.0 mmol, 75%) as colorless crystals.

15 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.63 (2H, m), 2.20 (2H, t, J=7.4 Hz), 2.83 (3H, d, J=4.0 Hz), 3.10 (2H, m), 6.52 (1H, d, J=5.4 Hz), 6.68 (1H, d, J=3.6 Hz), 6.87 (1H, s), 7.04 (1H, dd, J=2.4, 9.0 Hz), 7.37 (1H, d, J=2.4 Hz), 7.88 (1H, d, J=3.6 Hz), 8.03 (2H, m), 8.17

20 (1H, d, J=4.0 Hz), 8.29 (1H, d, J=9.0 Hz), 9.03 (1H, s), 12.05 (1H, s).

Example 33

5-(2-(3-(3-(Cyclopropylcarbamoyl)propyl)ureido)pyridin-

25 4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0223] The title compound (166 mg, 0.37 mmol, 76%) was

obtained as colorless crystals by performing the reaction similar to Example 8 using 4-((4-(1-methylcarbamoyl-1H-indol-5-yloxy)pyridin-2-yl)aminocarbonylamino)butyric acid (200 mg, 0.49 mmol, 5 Production example 32-1) and cyclopropylamine (0.028 ml, 0.58 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.33-0.37 (2H, m), 0.54-0.59 (2H, m), 1.62 (2H, m), 2.02 (2H, t, J=7.4 Hz), 2.58 (1H, m), 2.85 (3H, m), 3.08 (2H, m), 6.53 (1H, dd, 10 J=2.4, 6.0 Hz), 6.70 (1H, d, J=3.6 Hz), 6.88 (1H, d, J=2.4 Hz), 7.06 (1H, dd, J=2.4, 8.8 Hz), 7.39 (1H, d, J=2.4 Hz), 7.86 (1H, d, J=3.6 Hz), 7.90 (1H, d, J=3.6 Hz), 8.04 (1H, m), 8.05 (1H, d, J=6.0 Hz), 8.19 (1H, d, J=4.2 Hz), 8.31 (1H, d, J=8.8 Hz), 9.04 (1H, s).

Example 34

5-(2-(3-(4-(4-Hydroxy-4-methylpiperidin-1-yl)-4-oxobutyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0224] The title compound (195 mg, 0.383 mmol, 78.9%) was obtained as colorless crystals by performing the reaction similar to Example 8 using 4-((4-(1-methylcarbamoyl-1H-indol-5-yloxy)pyridin-2-yl)aminocarbonylamino)butyric acid (200 mg, 0.486 mmol, 25 Production example 32-1) and 4-hydroxy-4-methylpiperidine monohydrochloride (110 mg, 0.729 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.08 (3H, s), 1.22-1.44 (4H, m), 1.62 (2H, m), 2.27 (2H, t, J=7.4 Hz), 2.83 (3H, d, J=4.0 Hz), 2.97 (1H, m), 3.08 (2H, m), 3.29 (1H, m), 3.47 (1H, m), 3.89 (1H, m), 4.33 (1H, s),
5 6.50 (1H, d, J=6.0 Hz), 6.67 (1H, d, J=3.6 Hz), 6.87 (1H, s), 7.04 (1H, d, J=9.2 Hz), 7.36 (1H, s), 7.87 (1H, d, J=3.6 Hz), 8.01 (1H, m), 8.02 (1H, d, J=6.0 Hz), 8.16 (1H, m), 8.28 (1H, d, J=9.2 Hz), 9.00 (1H, m).

10 Example 35

5-(2-(3-(3-(Diethylcarbamoyl)propyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0225] The title compound (94 mg, 0.20 mmol, 64%) was obtained as colorless crystals by performing the
15 reaction similar to Example 8 using 4-((4-(1-methylcarbamoyl-1H-indol-5-yloxy)pyridin-2-yl)aminocarbonylamino)butyric acid (130 mg, 0.31 mmol, Production example 32-1) and diethylamine (0.066 ml, 0.63 mmol).

20 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.96 (3H, t, J=7.2 Hz), 1.04 (3H, t, J=7.2 Hz), 1.63 (2H, m), 2.25 (2H, t, J=7.2 Hz), 2.83 (3H, d, J=4.4 Hz), 3.09 (2H, m), 3.22 (4H, m), 6.51 (1H, dd, J=2.0, 5.6 Hz), 6.67 (1H, d, J=3.4 Hz), 6.86 (1H, d, J=2.0 Hz), 7.03 (1H, dd, J=2.4,
25 8.8 Hz), 7.36 (1H, d, J=2.4 Hz), 7.87 (1H, d, J=3.4 Hz), 8.02 (2H, m), 8.16 (1H, d, J=4.4 Hz), 8.29 (1H, d,

J=8.8 Hz), 9.00 (1H, s).

Example 36

5-(2-(3-(3-(Methylcarbamoyl)propyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0226] The title compound (107 mg, 0.25 mmol, 69%) was obtained as colorless crystals by performing the reaction similar to Example 8 using 4-((4-(1-methylcarbamoyl-1H-indol-5-yloxy)pyridin-2-yl)aminocarbonylamino)butyric acid (150 mg, 0.36 mmol, Production example 32-1) and methylamine hydrochloride (49 mg, 0.73 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.61 (2H, m), 2.03 (2H, t, J=7.6 Hz), 2.51 (3H, d, J=4.4 Hz), 2.83 (3H, d, J=4.0 Hz), 3.06 (2H, q, J=6.4 Hz), 6.50 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 6.86 (1H, d, J=2.4 Hz), 7.03 (1H, dd, J=2.4, 9.2 Hz), 7.36 (1H, d, J=2.4 Hz), 7.71 (1H, m), 7.87 (1H, d, J=3.6 Hz), 8.03 (2H, m), 8.16 (1H, d, J=4.4 Hz), 8.28 (1H, d, J=9.2 Hz), 9.01 (1H, s).

Example 37

N1-Methyl-5-(2-(pyrrolidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0227] Similarly to Example 5, the title compound (265 mg, 0.70 mmol, 69%) was obtained as white crystals from

phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (532 mg, 1.02 mmol) synthesized in Production example 5-2 and pyrrolidine (0.42 ml, 5.0 mmol).

5 MS Spectrum (ESI): 380 (M+1), 759 (2M+1)

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.78-1.84 (4H, m), 2.83 (3H, d, J=4.5 Hz), 3.22-3.36 (4H, m), 6.54 (1H, dd, J=2.3, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 7.03 (1H, dd, J=2.3, 8.7 Hz), 7.35 (1H, d, J=2.3 Hz), 7.41 (1H, d, J=2.3 Hz), 7.87 (1H, d, J=3.6 Hz), 8.04 (1H, d, J=5.6 Hz), 8.16 (1H, m), 8.28 (1H, t, J=8.7 Hz), 8.59 (1H, s).

Example 38

15 N1-Methyl-5-(2-(piperidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0228] Similarly to Example 5, the title compound (265 mg, 0.674 mmol, 76%) was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (463 mg, 0.885 mmol) synthesized in Production example 5-2 and piperidine (0.44 ml, 4.4 mmol).

20 MS Spectrum (ESI): 394 (M+1), 787 (2M+1)

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.37-1.57 (6H, m), 2.83 (3H, d, J=4.4 Hz), 3.26-3.45 (4H, m), 6.54 (1H, dd, J=2.4, 5.4 Hz), 6.67 (1H, d, J=3.4 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.30 (1H, d, J=2.4 Hz), 7.36 (1H, d,

J=2.4 Hz), 7.87 (1H, d, J=3.4 Hz), 8.05 (1H, d, J=5.4 Hz), 8.16 (1H, m), 8.28 (1H, t, J=8.8 Hz), 9.05 (1H, s).

Example 39

5 N1-Methyl-5-(2-((4-hydroxypiperidino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0229] Similarly to Example 27, the title compound (86.7 mg, 0.21 mmol, 21.2%) was obtained as white powder from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate (402 mg, 1.0 mmol) synthesized in Production example 29-1 and 4-hydroxypiperidine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.60-1.70 (2H, m), 1.75 (1H, m), 2.83 (3H, d, J=4.4 Hz), 2.95-3.01 (2H, m), 15 3.55-3.65 (2H, m), 3.71-3.76 (2H, m), 4.64 (1H, d, J=4.0 Hz), 6.53 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.32 (1H, d, J=2.4 Hz), 7.36 (1H, d, J=2.4 Hz), 7.87 (1H, d, J=3.6 Hz), 8.06 (1H, d, J=5.6 Hz), 8.16 (1H, q, J=4.4 Hz), 20 8.28 (1H, d, J=8.8 Hz), 9.10 (1H, s).

Example 40

N1-Methyl-5-(2-(4-oxopiperidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

25 [0230] Phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate

(440 mg, 0.841 mmol) synthesized in Production example 5-2 was dissolved in N,N-dimethylformamide (5 ml); triethylamine (0.543 ml, 3.90 mmol) and 4-piperidone hydrochloride monohydrate (0.530 g, 3.93 mmol) were added thereto; and the reaction mixture was stirred for 2 hours. The reaction mixture was partitioned between ethyl acetate and water. The organic layer was concentrated to yield the title compound (0.202 g, 0.496 mmol, 59%) as a colorless amorphous solid.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.32 (4H, t, J=4.9 Hz), 2.82 (3H, d, J=4.3 Hz), 3.68 (4H, t, J=4.9 Hz), 6.55 (1H, dd, J=2.3, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 7.03 (1H, dd, J=2.3, 8.6 Hz), 7.37 (2H, s), 7.87 (1H, d, J=3.6 Hz), 8.09 (1H, d, J=5.6 Hz), 8.17 (1H, s), 8.28 (1H, t, J=8.6 Hz), 9.37 (1H, s).

[0231]

Example 41

5-(2-(((4-Hydroxy-4-methylpiperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indole-1-carboxylic acid methylamide

4-Hydroxy-4-methylpiperidine monohydrochloride (508 mg, 3.83 mmol, Production example 8-3) was dissolved in N,N-dimethylformamide (8 ml); triethylamine (2 ml) was added; and the reaction mixture was stirred at room temperature. Phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-

(phenoxycarbonyl) carbamate (500 mg, 0.957 mmol, Production example 5-2) was added and the reaction mixture was stirred at room temperature for 8 hours. The reaction mixture was partitioned between ethyl acetate and water. The organic layer was dried over anhydrous magnesium sulfate, and concentrated under reduced pressure; and the residue was purified by silica gel column chromatography (Fuji Silysia BW-300, ethyl acetate, ethyl acetate: methanol = 20: 1 then 10: 1). The obtained amorphous solid was crystallized by adding diethyl ether: acetone = 2: 1. Thus obtained crystals were filtered off, washed with diethyl ether, and dried under aeration to yield the title compound (385 mg, 0.909 mmol, 95.0%).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.08 (3H, s), 1.33-1.40 (4H, m), 2.83 (3H, d, J=4.4 Hz), 3.14 (2H, m), 3.63 (2H, m), 4.27 (1H, s), 6.53 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.4 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.32 (1H, d, J=2.4 Hz), 7.35 (1H, d, J=2.4 Hz), 7.87 (1H, d, J=3.4 Hz), 8.06 (1H, d, J=5.6 Hz), 8.16 (1H, m), 8.28 (1H, d, J=8.8 Hz), 9.04 (1H, s).

Example 42

N1-Methyl-5-(2-((4-(1-hydroxy-1-methylethyl)piperidino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0232] Similarly to Example 28, the title compound (71.1 mg, 0.16 mmol, 29.7%) was obtained as white crystals from N1-ethyl-5-((2-amino-4-pyridyl)oxy)-1H-1-indolecarboxamide (150 mg, 0.53 mmol) synthesized in Production example 5-1 and 4-(1-hydroxy-1-methylethyl)piperidine (342 mg, 2.39 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.99 (6H, s), 1.03-1.09 (2H, m), 1.30 (1H, m), 1.60-1.64 (2H, m), 2.54-2.61 (2H, m), 2.83 (3H, d, J=4.4 Hz), 4.08 (1H, s), 4.10-4.15 (2H, m), 6.53 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.32 (1H, d, J=2.4 Hz), 7.36 (1H, d, J=2.4 Hz), 7.87 (1H, d, J=3.6 Hz), 8.06 (1H, d, J=5.6 Hz), 8.16 (1H, q, J=4.0 Hz), 8.27 (1H, d, J=8.8 Hz), 9.04 (1H, s).

[0233] 4-(1-Hydroxy-1-methylethyl)piperidine was synthesized in the following methods.

Production example 42-1

Benzyl 4-ethoxycarbonylpiperidine-1-carboxylate

4-Ethoxycarbonylpiperidine (1.572 g, 10.0 mmol) was dissolved in tetrahydrofuran (50 ml); triethylamine (2.79 ml, 20.0 mmol) and benzyl chlorocarbonate (1.71 ml, 12.0 mmol) were added dropwise while cooled with an ice water bath; and the reaction mixture was stirred at room temperature overnight. The reaction mixture was partitioned between ethyl acetate and the saturated

aqueous solution of sodium hydrogencarbonate. The organic layer was washed with brine and dried over anhydrous sodium sulfate. The solvent was distilled off, and the residue was purified by silica gel column chromatography (eluent; ethyl acetate: hexane = 1: 3) to yield the title compound (2.315 g, 7.95 mmol, 79.5%) as a colorless oil.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 1.26 (3H, t, J=7.2 Hz), 1.60-1.70 (2H, m), 1.80-2.00 (2H, m), 2.46 (1H, m), 2.80-3.00 (2H, m), 4.00-4.20 (2H, m), 4.15 (2H, q, J=7.2 Hz), 5.13 (2H, s), 7.29-7.38 (5H, m).

Production example 42-2

Benzyl 4-(1-hydroxy-1-methylethyl)piperidine-1-carboxylate

[0234] Benzyl 4-ethoxycarbonylpiperidine-1-carboxylate (2.315 g, 7.95 mmol) synthesized in Production example 42-1 was dissolved in tetrahydrofuran (25 ml) under nitrogen atmosphere; methyl magnesium bromide (0.93 M) in tetrahydrofuran (32.5 ml, 30.2 mmol) was added dropwise while cooled with an ice water bath; and the reaction mixture was stirred at room temperature overnight. The reaction mixture was partitioned between ethyl acetate and the saturated aqueous solution of ammonium chloride. The organic layer was washed with brine and dried over anhydrous sodium

sulfate. The solvent was distilled off, and the residue was purified by silica gel column chromatography (eluent; ethyl acetate: hexane = 1: 1) to yield the title compound (1.786 g, 6.44 mmol, 81%) as a colorless oil.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 1.18 (6H, s), 1.18-1.27 (2H, m), 1.40-1.48 (1H, m), 1.74-1.78 (2H, m), 2.60-2.80 (2H, m), 4.20-4.40 (2H, m), 5.13 (2H, s), 7.27-7.37 (5H, m).

Production example 42-3

4-(1-Hydroxy-1-methylethyl)piperidine

[0235] Benzyl 4-(1-hydroxy-1-methylethyl)piperidine-1-carboxylate (1.786 g, 6.44 mmol) synthesized in Production example 42-2 was dissolved in methanol (100 ml) under nitrogen atmosphere; 10% palladium on carbon (50% wet, 1.37 g) was added; the reaction system was purged with hydrogen at atmospheric pressure; and the reaction mixture was stirred overnight. After the reaction system was purged with nitrogen, the catalyst was filtered out, and washed with methanol; the solvent, together with the filtrate and the washing solution, was distilled off; and the residue was dried under reduced pressure to yield the title compound (922 mg, 6.44 mmol, quantitative) as pale gray crystals.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 1.18 (6H, s), 1.26-1.42

(3H, m), 1.74-1.80 (2H, m), 2.57-2.64 (2H, m), 3.14-3.22 (2H, m), 3.48 (1H, s).

Example 43

5-(2-(((4-(3-Methylcarbamoylpropyl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0236] 4-(1-((4-(1-Methylcarbamoyl-1H-indol-5-yloxy)pyridin-2-yl)aminocarbonyl)piperidin-4-yl)butyric acid (170 mg, 0.35 mmol) was dissolved in N,N-dimethylformamide (7.0 ml); methylamine hydrochloride (48 mg, 0.71 mmol), benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate (314 mg, 0.71 mmol) and triethylamine (0.35 ml) were added thereto; and the reaction mixture was stirred at room temperature for 2 hours. The reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Fuji Silysia NH silica gel, hexane-ethyl acetate-methanol system). After a small amount of acetone and ethyl acetate were added to the obtained amorphous solid; this solution was diluted with diethyl ether; and the solid portion was filtered off, washed with diethyl ether, and dried under aeration to yield the title compound (30 mg, 0.061 mmol,

17%) as a colorless amorphous solid.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.87-1.00 (2H, m),
 1.13 (2H, m), 1.33 (1H, m), 1.46 (2H, m), 1.57 (2H, m),
 1.99 (2H, t, J=7.4 Hz), 2.52 (3H, d, J=4.4 Hz), 2.65
 5 (2H, m), 2.83 (3H, d, J=4.0 Hz), 4.03 (2H, m), 6.53 (1H,
 d, J=6.0 Hz), 6.67 (1H, d, J=3.4 Hz), 7.03 (1H, d,
 J=9.0 Hz), 7.31 (1H, s), 7.35 (1H, s), 7.66 (1H, m),
 7.87 (1H, d, J=3.4 Hz), 8.06 (1H, d, J=4.0 Hz), 8.16
 (1H, d, J=4.0 Hz), 8.27 (1H, d, J=9.0 Hz), 9.05 (1H, s).

10

[0237] The starting materials were synthesized as follows.

Production example 43-1

tert-Butyl 4-(3-ethoxycarbonylpropyl)piperidine-1-
 15 carboxylate

tert-Butyl 4-(2-(toluene-4-sulfonyloxy)ethyl)piperidine-1-carboxylate (7.55 g,
 19.7 mmol, CAS No. 89151-45-1) as described in WO
 02/32872 was dissolved in ethanol; diethyl malonate
 20 (3.3 ml, 21.3 mmol) and sodium ethoxide (1.45 g, 21.3
 mmol) were added; and the reaction mixture was heated
 to reflux under nitrogen atmosphere for 2.5 hours.
 After naturally cooled to room temperature, the
 saturated aqueous solution of ammonium chloride was
 25 added; this was subjected to extraction with ethyl
 acetate, washed with brine, dried over anhydrous

magnesium sulfate, and concentrated under reduced pressure. After the residue was dissolved in dimethyl sulfoxide (20 ml); lithium chloride (1.7 g, 40 mmol) and water (0.36 ml, 20 mmol) were added; and the reaction mixture was stirred at 185 °C for 1.5 hours and further stirred at 195 °C for 2 hours. After naturally cooled to room temperature, the reaction mixture was partitioned between ethyl acetate-brine. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Fuji Silysia BW-300, hexane-ethyl acetate system) to yield the title compound (2.60 g, 8.7 mmol, 43%) as a pale yellow oil.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 1.02-1.13 (2H, m), 1.23-1.29 (5H, m), 1.39 (1H, m), 1.45 (9H, s), 1.62-1.69 (4H, m), 2.29 (2H, t, J=7.4 Hz), 2.67 (2H, m), 4.07 (2H, m), 4.13 (2H, q, J=7.2 Hz).

Production example 43-2

Ethyl 4-(piperidin-4-yl)butyrate

[0238] tert-Butyl 4-(3-ethoxycarbonylpropyl)piperidine-1-carboxylate (1.2 g, 4.0 mmol, Production example 43-1) was dissolved in trifluoroacetic acid (30 ml), and the reaction mixture was stirred at room temperature for 20 minutes. This was concentrated under reduced

pressure, and was further azeotropically distilled with toluene. The obtained residue was partitioned between ethyl acetate and a saturated aqueous solution of sodium hydrogencarbonate. The organic layer was dried
5 over anhydrous magnesium sulfate. In addition, the aqueous layer was concentrated under reduced pressure to dryness; the obtained solid was suspended in tetrahydrofuran; insoluble portion were filtered off, and this solution was added to the previously obtained
10 organic layer. This was purified by silica gel column chromatography (Fuji Silysia NH, hexane-ethyl acetate-methanol system) to yield the title compound (1.15g, quantitative) as a yellow oil.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 1.26 (3H, m), 1.28-1.37
15 (2H, m), 1.40-1.52 (3H, m), 1.64 (2H, m), 1.86 (2H, m), 2.29 (2H, t, J=7.4 Hz), 2.82 (2H, m), 3.35 (2H, m), 4.13 (2H, m).

Production example 43-3

20 5-(2-(((4-(3-Ethoxycarbonylpropyl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0239] Ethyl 4-(piperidin-4-yl)butyrate (650 mg, 2.0 mmol, Production example 43-2) was suspended in N,N-dimethylformamide (3.35 ml); phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-
25

(phenoxycarbonyl)carbamate (350 mg, 0.67 mmol, Production example 5-2) was added; and the reaction mixture was stirred at room temperature for 1 hour. The reaction mixture was partitioned between ethyl acetate and water. The organic layer was dried over anhydrous magnesium sulfate, concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Fuji Silysia BW-300, hexane-ethyl acetate-methanol system) to yield the title compound (271 mg, 0.54 mmol, 80%) as a pale yellow oil.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 1.05-1.16 (2H, m), 1.22-1.28 (5H, m), 1.43 (1H, m), 1.62 (2H, m), 1.71 (2H, m), 2.27 (2H, t, J=7.4 Hz), 2.80 (2H, m), 2.95 (3H, d, J=4.4 Hz), 3.99 (2H, m), 4.12 (2H, q, J=7.2 Hz), 6.09 (1H, d, J=4.4 Hz), 6.46 (1H, d, J=3.4 Hz), 6.58 (1H, dd, J=2.0, 5.6 Hz), 7.04 (1H, dd, J=2.0, 8.8 Hz), 7.24 (1H, s), 7.28 (1H, d, J=2.0 Hz), 7.32 (1H, d, J=3.4 Hz), 7.54 (1H, d, J=2.0 Hz), 8.03 (1H, d, J=5.6 Hz), 8.20 (1H, d, J=8.8 Hz).

Production example 43-4

4-(1-((4-(1-Methylcarbamoyl-1H-indol-5-yloxy)pyridin-2-yl)aminocarbonyl)piperidin-4-yl)butyric acid

[0240] 5-(2-((4-(3-Ethoxycarbonylpropyl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-indole-1-

carboxylic acid methylamide (271 mg, 0.54 mmol,

Production example 43-3) was dissolved in tetrahydrofuran (3.0 ml) and methanol (1.5 ml); 4N lithium hydroxide (0.54 ml) was added; and the reaction mixture was stirred at room temperature for 3.5 hours.

5 1N hydrochloric acid (2.2 ml) was added thereto while the stirred at room temperature. After the precipitated crystals were filtered off, the crystals were washed with water and diethyl ether sequentially, and dried under aeration to yield the title compound
10 (170 mg, 0.35 mmol, 66%) as colorless crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.93 (2H, m), 1.16 (2H, m), 1.36 (1H, m), 1.47 (2H, m), 1.58 (2H, m), 2.15 (2H, t, J=7.4 Hz), 2.66 (2H, m), 2.83 (3H, d, J=4.2 Hz), 4.02 (2H, m), 6.53 (1H, d, J=6.0 Hz), 6.67 (1H, d, J=3.4 Hz),
15 7.03 (1H, d, J=9.2 Hz), 7.31 (1H, s), 7.35 (1H, s), 7.86 (1H, d, J=3.4 Hz), 8.05 (1H, d, J=6.0 Hz), 8.15 (1H, d, J=4.2 Hz), 8.27 (1H, d, J=9.2 Hz), 9.02 (1H, s).

20 Example 44

5-(2-(((4-(3-Carbamoylpropyl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0241] 4-(Piperidin-4-yl)butanamide (547 mg, 1.41 mmol)
25 was dissolved in N,N-dimethylformamide (3 ml); phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-

pyridyl)-N-(phenoxycarbonyl)carbamate (210 mg, 0.402 mmol, the product of Production example 5-2) was added thereto; and the reaction mixture was stirred at room temperature for 1.5 hour. The reaction mixture was partitioned between ethyl acetate and water; the organic layer was dried over anhydrous magnesium sulfate, and concentrated under reduced pressure; and the residue was purified by silica gel column chromatography (Fuji Silysia NH, ethyl acetate-methanol system). The obtained amorphous solid was crystallized by adding diethyl ether. After addition of a small amount of ethanol to make a suspension, this was diluted with hexane. After separation by filtration to obtain crystals, these were rinsed with diethyl ether and dried under aeration. Thus, the title compound was obtained as colorless crystals (157 mg, 0.328 mmol, 81.7%).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.87-1.00 (2H, m), 1.10-1.16 (2H, m), 1.35 (1H, m), 1.42-1.50 (2H, m), 1.58 (2H, m), 1.98 (2H, t, J=7.4 Hz), 2.65 (2H, m), 2.83 (3H, d, J=4.0 Hz), 4.03 (2H, m), 6.53 (1H, dd, J=2.0, 5.6 Hz), 6.67 (2H, m), 7.03 (1H, dd, J=2.0, 9.0 Hz), 7.20 (1H, s), 7.31 (1H, d, J=2.0 Hz), 7.35 (1H, d, J=2.0 Hz), 7.87 (1H, d, J=3.2 Hz), 8.06 (1H, d, J=5.6 Hz), 8.16 (1H, m), 8.28 (1H, d, J=9.0 Hz), 9.05 (1H, s).

[0242] The starting materials were synthesized as follows.

Production example 44-1

tert-Butyl 4-(3-carbamoylpropyl)piperidine-1-
5 carboxylate

tert-Butyl 4-(3-ethoxycarbonylpropyl)piperidine-
1-carboxylate (0.60 g, 2.0 mmol, the product of
Production example 43-1) and formamide (0.27 ml, 6.7
mmol) were dissolved in N,N-dimethylformamide (1.0 ml);
10 sodium ethoxide (0.095 g, 1.4 mmol) was added thereto
while stirred and heated at 100 °C; the reaction
mixture was stirred for 2 hours under nitrogen
atmosphere. After cooled to room temperature, the
reaction mixture was partitioned between water and
15 ethyl acetate. The organic layer was washed with brine,
dried over anhydrous magnesium sulfate, and then the
solvent was distilled off under reduced pressure. The
residue was purified by silica gel chromatography
(eluent; hexane-ethyl acetate = 95:5 to 85:15). The
20 title compound was obtained as a colorless oil (0.38 g,
1.4 mmol, 70%).

¹H-NMR Spectrum (CDCl₃) δ(ppm): 1.03-1.14 (2H, m),
1.26-1.31(2H, m), 1.35-1.45 (1H, m), 1.46 (9H, s),
1.63-1.71 (4H, m), 2.22 (2H, t, J=7.6 Hz), 2.67 (2H, m),
25 4.07 (2H, brs), 5.30 (1H, brs), 5.39 (1H, brs).

Production example 44-24-(Piperidin-4-yl)butanamide

[0243] tert-Butyl 4-(3-carbamoylpropyl)piperidine-1-carboxylate (0.38 g, 1.4 mmol, Production example 44-1) was dissolved in trifluoroacetic acid (2 ml) and the reaction mixture was stirred at room temperature for 20 minutes. The reaction mixture was concentrated under reduced pressure and then azeotropically distilled with toluene. The residue was partitioned between tetrahydrofuran and a saturated aqueous solution of sodium hydrogencarbonate; and the organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure; and the residue was purified by silica gel chromatography (Fuji Silysia NH, ethyl acetate-methanol system) to yield the title compound (0.55 g, quantitative) as pale yellow oil.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.90-1.01 (2H, m), 1.09-1.15 (2H, m), 1.26 (1H, m), 1.45 (2H, m), 1.55 (2H, m), 1.98 (2H, t, J=7.4 Hz), 2.43 (2H, m), 2.91 (2H, m), 6.65 (1H, s), 7.20 (1H, s).

Example 455-(2-((4-Pyrrolidin-1-yl)carbonyl)piperidin-1-yl)carbonylamino)pyridin-4-yloxy)-1H-indole-1-carboxylic acid methanamide

[0244] Similarly to Example 5, the title compound (134

mg, 0.273 mmol, 91%) was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (157 mg, 0.300 mmol) synthesized in Production example 5-2 and (piperidin-4-yl)-(pyrrolidin-1-yl)methanone (328 mg, 1.50 mmol) obtained from N-benzyloxycarbonylisonipecotic acid and pyrrolidine by the method similar to Example 21.

¹H NMR Spectrum (DMSO-d₆) δ (ppm): 1.35-1.48 (2H, m), 1.56-1.65 (2H, m), 1.71-1.80 (2H, m), 1.82-1.91 (2H, m), 2.61 (1H, m), 2.73-2.84 (2H, m), 2.85 (3H, d, J=4.4 Hz), 3.22-3.28 (2H, m), 3.44-3.50 (2H, m), 4.04-4.12 (2H, m), 6.56 (1H, d, J=6.0 Hz), 6.69 (1H, d, J=3.6 Hz), 7.06 (1H, dd, J=2.4, 9.2 Hz), 7.34 (1H, s), 7.38 (1H, d, J=2.4 Hz), 7.89 (1H, d, J=3.6 Hz), 8.09 (1H, d, J=6.0 Hz), 8.18 (1H, q, J=4.4 Hz), 8.30 (1H, d, J=9.2 Hz), 9.16 (1H, s).

Example 46

N1-Methyl-5-(2-(((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide

[0245] Similarly to Example 27, the title compound (88.5 mg, 0.19 mmol, 63.8%) was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate (121 mg, 0.30 mmol,

Production example 29-1) and 4-(1-pyrrolidinyl)piperidine.

Phenyl N1-methyl-5-(2-(((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide may be synthesized by the following methods.

Phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (12.1 g, 23.2 mmol) synthesized in Production example 5-2 was dissolved in dimethylformamide (150 ml); 4-(1-pyrrolidinyl)piperidine (14.4 g, 93.3 mmol) was added thereto; and the reaction mixture was stirred at room temperature for 16 hours. The reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine and concentrated to about 100 ml. The residue was allowed to be kept cool at 5 °C for overnight to precipitate crystals. The crystals were filtered off, washed with ethyl acetate to yield the title compound (7.8 g, 16.9 mmol, 73%) as white crystals.

¹H NMR Spectrum (DMSO-d₆) δ (ppm): 1.20-1.33 (2H, m), 1.60-1.70 (4H, m), 1.70-1.80 (2H, m), 2.40-2.60 (5H, m), 2.77-2.84 (5H, m), 3.90-4.00 (2H, m), 6.54 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz) 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.31 (1H, s), 7.35 (1H, d, J=2.4 Hz), 7.87 (1H, d, J=3.6 Hz), 8.06 (1H, d, J=5.6 Hz), 8.16

(1H, m), 8.28 (1H, d, J=8.8 Hz), 9.11 (1H, s).

Example 47

N1-Methyl-5-(2-(((4-(piperidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide

[0246] Similarly to Example 27, the title compound (94.6 mg, 0.20 mmol, 66.2%) as white crystals was obtained from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate (121 mg, 0.30 mmol, Production example 29-1) and 4-piperidinopiperidine.

N1-Methyl-5-(2-(((4-(piperidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide may be prepared by the following methods.

Phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (15.5 g, 29.7 mmol) synthesized in Production example 5-2 was dissolved in dimethylformamide (180 ml); 4-piperidinopiperidine (20.0 g, 119 mmol) was added thereto; and the reaction mixture was stirred at room temperature for 9 hours. The reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine and concentrated to about 100 ml. The residue was allowed to be kept cool at 5 °C overnight to precipitate crystals. The

crystals were filtered off and washed with ethyl acetate to yield the title compound (4.0 g, 8.4 mmol, 28%) as white crystals.

¹H NMR Spectrum (DMSO-d₆) δ (ppm): 1.20-1.65 (10H, m),
5 2.31-2.40 (5H, m), 2.66 (2H, m), 2.83 (3H, d, J=4.4 Hz),
4.08 (2H, m), 6.53 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d,
J=3.6 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.31 (1H, d,
J=2.4 Hz), 7.35 (1H, d, J=2.4 Hz), 7.87 (1H, d, J=3.6
Hz), 8.06 (1H, d, J=5.6 Hz), 8.16 (1H, q, J=4.4 Hz),
10 8.28 (1H, d, J=8.8 Hz), 9.09 (1H, s).

Example 48

N1-Methyl-5-(2-((4-ethylpiperazin-1-yl)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

15 [0247] Similarly to Example 27, the title compound
(73.2 mg, 0.17 mmol, 57.8%) was obtained as white
powder from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-
indolyl)oxy-2-pyridyl)carbamate (121 mg, 0.30 mmol,
Production example 29-1) and 1-ethylpiperazine.

20 ¹H NMR Spectrum (DMSO-d₆) δ (ppm): 0.97 (3H, t, J=7.2
Hz), 2.25-2.32 (6H, m), 2.83 (3H, d, J=4.0 Hz), 3.20-
3.40 (4H, m), 6.55 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d,
J=3.6 Hz), 7.04 (1H, dd, J=2.4, 8.8 Hz), 7.31 (1H, d,
J=2.4 Hz), 7.36 (1H, d, J=2.4 Hz), 7.87 (1H, d, J=3.6
25 Hz), 8.07 (1H, d, J=5.6 Hz), 8.16 (1H, q, J=4.0 Hz),
8.28 (1H, d, J=8.8 Hz), 9.13 (1H, s).

Example 49N1-Methyl-5-(2-((4-(2-hydroxyethyl)piperazin-1-yl)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

5 [0248] Similarly to Example 27, the title compound (97.6 mg, 0.22 mmol, 59.7%) was obtained as pale pink powder from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate (150 mg, 0.37 mmol, Production example 29-1) and 1-(2-hydroxyethyl)piperazine.

¹H NMR Spectrum (DMSO-d₆) δ (ppm): 2.30-2.40 (6H, m), 2.83 (3H, d, J=4.0 Hz), 3.20-3.40 (4H, m), 3.46 (2H, m), 4.39 (1H, t, J=5.6 Hz), 6.55 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.31 (1H, d, J=2.4 Hz), 7.35 (1H, d, J=2.4 Hz), 7.87 (1H, d, J=3.6 Hz), 8.06 (1H, d, J=5.6 Hz), 8.16 (1H, q, J=4.0 Hz), 8.27 (1H, d, J=8.8 Hz), 9.12 (1H, s).

Example 50

20 N1-Methyl-5-(2-((3-methylsulfonylpropylamino)carbonyl)amino-4-pyridyl)-oxy-1H-1-indolecarboxamide

[0249] Similarly to Example 28, the title compound (166.8 mg, 0.37 mmol, 70.5%) was obtained as white crystals from N1-methyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (150 mg, 0.53 mmol, Production

example 5-1) and 3-methylsulfonylpropylamine hydrochloride (410 mg, 2.36 mmol).

¹H NMR Spectrum (DMSO-d₆) δ (ppm): 1.70-1.90 (2H, m), 2.83 (3H, d, J=4.4 Hz), 2.94 (3H, s), 3.04-3.09 (2H, m), 3.17-3.24 (2H, m), 6.52 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 6.86 (1H, s), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, s), 7.87 (1H, d, J=3.6 Hz), 8.03 (1H, d, J=5.6 Hz), 8.10-8.17 (2H, m), 8.28 (1H, d, J=8.8 Hz), 9.07 (1H, s).

Example 51

N1-Methyl-5-(2-((4-(2-dimethylaminoacetyl)piperazin-1-yl)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0250] Similarly to Example 28, the title compound (189.8 mg, 0.40 mmol, 74.5%) was obtained as white powder from N1-methyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (150 mg, 0.53 mmol, Production example 5-1) and 1-(2-dimethylaminoacetyl)piperazine (500 mg, 2.92 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.14 (6H, s), 3.04 (3H, d, J=4.0 Hz), 3.29 (2H, s), 3.20-3.49 (8H, m), 6.56 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz) 7.30 (1H, d, J=2.4 Hz), 7.36 (1H, d, J= 2.4 Hz), 7.87 (1H, d, J=3.6 Hz), 8.08 (1H, d, J=5.6 Hz), 8.16 (1H, q, J=4.0 Hz), 8.28 (1H, d, J=8.8 Hz), 9.24 (1H, s).

[0251] 1-(2-Dimethylaminoacetyl)piperazine was prepared by the following methods.

Production example 51-1

5 Benzyl 4-(2-dimethylaminoacetyl)piperazine-1-carboxylate

Benzyl piperazine-1-carbamate (2.203 g, 10.0 mmol) was dissolved in tetrahydrofuran (50 ml); 2-dimethylaminoacetic acid (1.24 g, 12.0 mmol), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide
10 hydrochloride(2.30 g, 12.0 mmol), 1-hydroxy-1H-benzotriazole monohydrate (1.84 g, 12.0 mmol) and triethylamine (3.35 ml, 24.0 mmol) were added thereto; and the reaction mixture was stirred at room temperature for 7 hours. The reaction mixture was
15 partitioned between ethyl acetate and a saturated aqueous solution of sodium hydrogencarbonate. The organic layer was washed with a saturated aqueous solution of sodium hydrogencarbonate, water and brine, dried over anhydrous sodium sulfate, and the residue
20 was purified by NH silica gel column chromatography (eluent; ethyl acetate: hexane = 3: 1) to yield the title compound (954 mg, 3.12 mmol, 31.2%) as a colorless oil.

¹H-NMR Spectrum (CDCl₃) δ (ppm): 2.26 (6H, s), 3.11 (2H, s), 3.45-3.65 (8H, m), 5.15 (2H, s), 7.32-7.38 (5H, m).

25

Production example 51-21-(2-Dimethylaminoacetyl)piperazine

[0252] Benzyl 4-(2-dimethylaminoacetyl)piperazine-1-carbamate (954 mg, 3.12 mmol) synthesized in Production example 51-1 was dissolved in methanol (50 ml) under nitrogen atmosphere; 10% palladium on carbon (50% wet, 665 mg) was added thereto; the reaction system was purged with hydrogen at atmospheric pressure; and the reaction mixture was stirred overnight. After the reaction system was purged with nitrogen, the catalyst was filtered out, and washed with methanol. The solvent, together with the filtrate and washing solution, was distilled off, and the residue was dried under reduced pressure to yield the title compound (508 mg, 2.97 mmol, 95.0%) as a colorless oil.

¹H-NMR Spectrum (CDCl₃) δ (ppm): 2.28 (6H, s), 2.80-2.88 (4H, m), 3.11 (2H, s), 3.52-3.62 (4H, m).

Example 52

20 N1-Methyl-5-(2-((4-cyclohexylpiperazin-1-yl)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide
[0253] Similarly to Example 27, the title compound (121.3 mg, 0.25 mmol, 68.2%) was obtained as white crystals from phenyl N-(4-(1-methylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate (150 mg, 0.37 mmol, Production example 29-1) and 1-cyclohexylpiperazine.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.00-1.20 (6H, m),
 1.53 (2H, m), 1.60-1.80 (4H, m), 2.19 (2H, m), 2.30-
 2.45 (5H, m), 2.83 (3H, d, J=4.0 Hz), 6.54 (1H, dd,
 J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 7.03 (1H, dd,
 5 J=2.4, 8.8 Hz), 7.31 (1H, d, J=2.4 Hz), 7.35 (1H, d,
 J=2.4 Hz), 7.87 (1H, d, J=3.6 Hz), 8.06 (1H, d, J=5.6
 Hz), 8.16 (1H, q, J=4.0 Hz), 8.27 (1H, d, J=8.8 Hz),
 9.09 (1H, s).

10 Example 53

N4-(4-(1-(Methylamino)carbonyl-1H-5-indolyl)oxy-2-
 pyridyl)-4-morpholinecarboxamide

[0254] Similarly to Example 27, the title compound
 (58.6 mg, 0.15 mmol, 49.4%) was obtained as white
 15 powder from phenyl N-(4-((1-((methylamino)carbonyl)-1H-
 5-indolyl)oxy-2-pyridyl)carbamate (121 mg, 0.30 mmol,
 Production example 29-1) and morpholine.

N4-(4-(1-(Methylamino)carbonyl-1H-5-indolyl)-
 oxy-2-pyridyl)-4-morpholinecarboxamide may be prepared
 20 by the following methods.

Phenyl N-(4-(1-(methylamino)carbonyl-1H-5-
 indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (20
 g, 38 mmol) synthesized in Production example 5-2 was
 dissolved in N,N-dimethylformamide (190 ml); morpholine
 25 (13.3 mg, 153 mmol) was added thereto; and the reaction
 system was stirred at room temperature for 9 hours.

The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The obtained
5 residue was dissolved in ethyl acetate and a small amount of tetrahydrofuran; this suspension was filtrated with silica gel; and ethyl acetate and three different ratio of solvent mixtures of ethyl acetate: methanol = 20: 1, 10: 1, and 5: 1 were eluted through
10 the gel. The filtrate was concentrated under reduced pressure. The residue was dissolved in diethyl ether (40 ml); hexane (200 ml) was added thereto; and precipitated insoluble syrupy portion was removed from the solution; and the resultant solution was
15 concentrated again under reduced pressure. The residue was dissolved in ethyl acetate (300 ml) and was allowed to stand at room temperature. After the crystals were precipitated, the crystals were filtered off, washed with ethyl acetate, and dried to yield the crude
20 crystals of the title compound (10.3 g). 9 g of this crude crystals was suspended in a mixture of tetrahydrofuran (3 ml) and N,N-dimethylformamide (3 ml each); this suspension was diluted with ethanol (60 ml); and the crystals were filtered off, washed with
25 ethanol and dried to yield the title compound as colorless crystals (7.70 g, 19 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.83 (3H, d, J=4.4 Hz), 3.34-3.38 (4H, m), 3.50-3.53 (4H, m), 6.56 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 7.04 (1H, dd, J=2.4, 8.8 Hz), 7.31 (1H, d, J=2.4 Hz), 7.36 (1H, d, J=2.4 Hz), 7.87 (1H, d, J=3.6 Hz), 8.08 (1H, d, J=5.6 Hz), 8.17 (1H, q, J=4.4 Hz), 8.28 (1H, d, J=8.8 Hz), 9.19 (1H, s).

Example 54

10 N1-Methyl-5-(2-((1,1-dioxothiomorpholin-4-ylcarbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide

[0255] Phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate
 15 (150 mg, 0.278 mmol, Production example 5-2) was dissolved in N,N-dimethylformamide (1.5 ml); 5N aqueous solution of sodium hydroxide (0.29 ml) and 1,1-dioxothiomorpholine hydrochloride (246 mg, 1.44 mmol) were added thereto; and the reaction mixture was
 20 stirred at room temperature for 5 hours. The reaction mixture was partitioned between ethyl acetate and water. The organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column
 25 chromatography (Fuji Silysia BW-300, ethyl acetate). Diethyl ether was added to this to allow to

crystallize; and the crystals were filtered off, washed with diethyl ether, and dried under aeration to yield the title compound as colorless crystals (100 mg, 0.226 mmol, 78.5%).

5 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.83 (3H, d, J=3.6 Hz), 3.10 (4H, m), 3.81 (4H, m), 6.57 (1H, dd, J=1.2, 5.6 Hz), 6.67 (1H, d, J=3.2 Hz), 7.03 (1H, dd, J=2.0, 9.2 Hz), 7.32 (1H, m), 7.36 (1H, d, J=2.0 Hz), 7.87 (1H, d, J=3.2 Hz), 8.09 (1H, d, J=5.6 Hz), 8.16 (1H, d, J=3.6 Hz), 8.28 (1H, d, J=9.2 Hz), 9.54 (1H, s).

10

[0256] The starting material was synthesized by the following methods.

Production examples 54-1

15 tert-Butyl thiomorpholine-4-carboxylate

Thiomorpholine (5.0 ml, 53 mmol) was dissolved in tetrahydrofuran (200 ml); triethylamine (8.1 ml, 58 mmol) was added thereto; and the reaction mixture was stirred at room temperature. tert-Butoxycarbonyl dicarbonate (13.3 ml, 58 mmol) was added thereto and the reaction mixture was stirred at room temperature for 10 hours. The reaction mixture was concentrated under reduced pressure; and the residue was purified by silica gel column (eluent; hexane: ethyl acetate = from 80: 20, 75: 25 to 70: 30) to yield the title compound as colorless crystals (10.4 g, 51 mmol).

20

25

¹H-NMR Spectrum (CDCl₃) δ (ppm): 1.46 (9H, s), 2.57 (4H, m), 3.69 (4H, m).

Production example 54-2

5 tert-Butyl 1,1-dioxothiomorpholine-4-carboxylate

[0257] tert-Butyl thiomorpholine-4-carboxylate (1.91 g, 9.42 mmol) was dissolved in dichloromethane (50 ml); m-chloroperbenzoic acid (5.0 g, 19 mmol) was gradually added while cooled with ice bath, stirred, and under
10 nitrogen atmosphere; and the reaction mixture was stirred at room temperature for 12 hours. After addition of a saturated aqueous solution of sodium thiosulfate, the reaction mixture was kept stirred for a while; and this was subjected to extraction with
15 ethyl acetate, washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. Triethylamine (8.1 ml, 58 mmol) were added to the obtained crystals; and the reaction mixture was stirred at room temperature. tert-Butoxycarbonyl dicarbonate
20 (13.3 ml, 58 mmol) was added thereto; and the reaction mixture was stirred at room temperature for 10 hours. The reaction mixture was concentrated under reduced pressure; and the obtained crystals were suspended with a solvent mixture of diethyl ether: ethanol = 10: 1,
25 filtered off, washed with diethyl ether and dried under aeration to yield the title compound as colorless

crystals (2.03 g, 8.63 mmol, 91.6%).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.40 (9H, s), 3.09 (4H, t, J=5.2 Hz), 3.72 (4H, t, J=5.2 Hz).

5 Production example 54-3

Thiomorpholine 1,1-dioxide monohydrochloride

[0258] tert-Butyl 1,1-dioxothiomorpholine-4-carboxylate (2.03 g, 8.63 mmol) was dissolved in a mixture of hydrochloric acid-methanol 10 (20 ml, purchased from Tokyo Kasei Kogyo Co., Ltd) and tetrahydrofuran (20 ml); hydrochloric acid (4.0 ml) was added thereto during stirring at room temperature; and the reaction mixture was stirred at room temperature for 3 hours. The reaction mixture was concentrated; methanol (20 ml), 15 tetrahydrofuran (20 ml) and hydrochloric acid (4.0 ml) were added to the obtained crystals. Furthermore, water (10 ml) was added to this solution to perfectly dissolve the crystals; and this solution was stirred at room temperature for 1 hour. The solvent was 20 concentrated under reduced pressure; and the obtained crystals were suspended in methanol, filtered off, washed with methanol, and dried under aeration to yield the title compound as colorless crystals (1.49 g, 8.65 mmol, quantitative).

25 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 3.54 (8H, m), 9.83 (2H, brs).

Example 55

5-(2-(3-((1R)-1-Hydroxymethyl-2-oxo-2-pyrrolidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid ethylamide

[0259] Similarly to Example 5, the title compound (118 mg, 0.246 mmol, 82%) was obtained as white crystals from phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyloxy-2-pyridyl)-N-(phenoxycarbonyl)carbamate (161 mg, 0.300 mmol), and (2R)-2-amino-3-hydroxy-1-(pyrrolidin-1-yl)propan-1-one (265 mg, 1.67 mmol) obtained by the method similar to Example 21 from (2R)-2-benzyloxycarbonylamino-3-hydroxypropionic acid and pyrrolidine.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.19 (3H, t, J=7.2 Hz), 1.70-1.90 (4H, m), 3.20-3.60 (8H, m), 4.54 (1H, m), 4.98 (1H, brs), 6.55 (1H, d, J=6.0 Hz), 6.69 (1H, d, J=3.6 Hz), 6.97 (1H, s), 7.05 (1H, dd, J=2.4, 8.8 Hz), 7.39 (1H, d, J=2.4 Hz), 7.92 (1H, d, J=3.6 Hz), 8.05 (1H, d, J=6.0 Hz), 8.08-8.28 (2H, m), 8.30 (1H, d, J=8.8 Hz), 9.21 (1H, s).

[0260] The starting material was synthesized as follows.

Production example 55-1

Phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate

The reaction similar to Production example 5-2 was performed by using N1-ethyl-5-(2-aminopyridin-4-yloxy)-1H-indolecarboxamide (2.9 g, 9.9 mmol, Production example 27-1), tetrahydrofuran, triethylamine and phenyl chloroformate; the extraction and washing was performed; the obtained residue was crystallized by addition of a solvent mixture of diethyl ether: hexane = 1: 1; and the obtained crystals were filtered off, washed with diethyl ether, and dried under aeration to yield the title compound as pale pink crystals (3.7 g, 6.9 mmol, 70%).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.17 (3H, t, J=7.2 Hz), 3.29 (2H, m), 6.66 (1H, d, J=3.4 Hz), 6.96 (1H, dd, J=2.0, 5.8 Hz), 7.09 (1H, dd, J=2.0, 8.0 Hz), 7.17 (4H, d, J=8.0 Hz), 7.29 (2H, d, J=8.0 Hz), 7.41-7.44 (5H, m), 7.51 (1H, d, J=2.0 Hz), 7.92 (1H, d, J=3.4 Hz), 8.22 (1H, m), 8.31 (1H, d, J=8.8 Hz), 8.42 (1H, d, J=5.8 Hz).

Example 56

5-(2-(3-((1S)-1-Hydroxymethyl-2-oxo-2-pyrrolidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid ethylamide

[0261] Similarly to Example 5, the title compound (132 mg, 0.275 mmol, 92%) was obtained as white crystals from phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxy carbonyl) carbamate

(161 mg, 0.300 mmol) synthesized in Production example 55-1 and (2S)-2-amino-3-hydroxy-1-(pyrrolidin-1-yl)propan-1-one hydrochloride (synthesized as an intermediate in Example 18).

5

Example 57

5-(2-(3-((1R)-1-Hydroxymethyl-2-oxo-2-piperidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid ethylamide

10 [0262] Similarly to Example 5, the title compound (127 mg, 0.257 mmol, 86%) was obtained as white crystals from phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (161 mg, 0.300 mmol) and (2R)-2-amino-3-hydroxy-1-
15 (piperidin-1-yl)propan-1-one (228 mg, 1.32 mmol, synthesized as an intermediate in Example 21).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.19 (3H, t, J=7.2 Hz), 1.38-1.61 (6H, m), 3.25-3.53 (8H, m), 4.75 (1H, m), 4.92 (1H, brs), 6.54 (1H, dd, J=2.4, 6.0 Hz), 6.69 (1H, d, J=3.6 Hz), 6.97 (1H, d, J=2.4 Hz), 7.05 (1H, dd, J=2.4, 9.0 Hz), 7.38 (1H, d, J=2.4 Hz), 7.92 (1H, d, J=3.6 Hz), 8.05 (1H, d, J=6.0 Hz), 8.08-8.27 (2H, m), 8.30 (1H, d, J=9.0 Hz), 9.21 (1H, s).

25

Example 58

5-(2-(3-((1S)-1-Hydroxymethyl-2-oxo-2-piperidin-1-

ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic
acid ethylamide

[0263] Similarly to Example 5, the title compound (54.4 mg, 0.110 mmol, 73%) was obtained as white crystals from phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxy-carbonyl)carbamate (80.1 mg, 0.150 mmol) synthesized in Production example 55-1 and (2S)-2-amino-3-hydroxy-1-(piperidin-1-yl)propan-1-one hydrochloride (156 mg, 0.748 mmol, synthesized as an intermediate in Example 20).

Example 59

5-(2-(3-(2-(4-Hydroxy-4-methylpiperidin-1-yl)-2-oxoethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic
acid ethylamide

[0264] The reaction similar to Example 5 was performed by using ((4-(1-ethylcarbamoyl-1H-indol-5-yloxy)pyridin-2-yl)aminocarbonylamino)acetic acid (149 mg, 0.37 mmol) and 4-hydroxy-4-methylpiperidine monohydrochloride (68 mg, 0.45 mmol, Production example 8-3); purification was performed by silica gel column chromatography (Fuji Silysia BW-300, eluent, ethyl acetate: methanol = 9: 1; Fuji Silysia NH, eluent, ethyl acetate: methanol = 10: 1; and again Fuji Silysia BW-300, eluent, ethyl acetate-methanol system); and the obtained crystals were suspended in diethyl ether and

filtered off, washed with diethyl ether and dried under aeration to yield the title compound as colorless crystals (40 mg, 0.081 mmol, 22%).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.10 (3H, s), 1.16
5 (3H, t, J=7.2 Hz), 1.43 (4H, m), 3.01 (2H, m), 3.36 (2H, m), 3.89 (2H, m), 3.96 (2H, d, J=4.4 Hz), 4.37 (1H, s), 6.52 (1H, d, J=5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 6.91 (1H, s), 7.03 (1H, d, J=9.0 Hz), 7.37 (1H, s), 7.90 (1H, d, J=3.6 Hz), 8.03 (1H, d, J=5.6 Hz), 8.17 (1H, m),
10 8.22 (1H, m), 8.28 (1H, d, J=9.0 Hz), 9.27 (1H, s).

[0265]The starting material was synthesized as follows.

Production example 59-1

((4-(1-Ethylcarbamoyl-1H-indol-5-yloxy)pyridin-2-yl)aminocarbonylamino)acetic acid
15

Methyl aminoacetate hydrochloride (292 mg, 2.33 mmol) was suspended in a solvent mixture of N,N-dimethylformamide (4 ml) and triethylamine (1 ml); phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (250 mg, 0.466
20 mmol, Production example 55-1) was added thereto; and the reaction mixture was stirred at room temperature for 2 days. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer
25 was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The

obtained residue was dissolved in a solvent mixture of tetrahydrofuran (2 ml) and methanol (1 ml); and 4N aqueous solution of sodium hydroxide was added thereto while stirred at room temperature; and the reaction mixture was stirred for 1.5 hour at room temperature. After 1N hydrochloric acid was added, extraction was performed with ethyl acetate-tetrahydrofuran, washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained crystals were suspended in diethyl ether, filtered off, washed with dimethyl ether, and dried under aeration to yield the title compound as colorless crystals (149 mg, 0.375 mmol, 80.5%).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.17 (3H, t, J=7.0 Hz), 3.36 (2H, d, J=7.0 Hz), 3.81 (2H, d, J=5.2 Hz), 6.54 (1H, d, J=5.6 Hz), 6.67 (1H, d, J=3.4 Hz), 6.85 (1H, s), 7.04 (1H, dd, J=2.0, 8.8 Hz), 7.37 (1H, d, J=2.0 Hz), 7.90 (1H, d, J=3.4 Hz), 8.05 (1H, d, J=5.6 Hz), 8.20-8.30 (3H, m), 9.27 (1H, s), 12.55 (1H, s).

Example 60

N1-Ethyl-5-(2-(((1-methyl-4-piperidyl)methyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0266] Similarly to Example 27, a crude product of tert-butyl 4-(((4-((1-(ethylamino)carbonyl-1H-5-

indolyl)oxy)-2-

pyridyl)amino)carbonyl)amino)methyl)piperidin-1-

carboxylate was obtained from phenyl N-(4-(1-(ethylamino)carbonyl)-1H-5-indolyl)oxy)-2-

5 pyridyl)carbamate (150 mg, 0.36 mmol, Production example 27-2) and tert-butyl 4-aminomethyl-1-piperidine carboxylate. Trifluoroacetic acid was added to this at room temperature; the solution was stirred for 30 minutes; trifluoroacetic acid was distilled off;
10 triethylamine-methanol was added to the residue to neutralize; and the solvent was distilled off again under reduced pressure. The residue was dissolved in tetrahydrofuran (4.0 ml)-methanol (4.0 ml); acetic acid (0.1 ml), 37% aqueous formaldehyde solution (0.5 ml)
15 and sodium cyanoborohydride (90.5 mg, 1.44 mmol) were added at room temperature; and the reaction mixture was stirred for 1 hour. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was washed with water and brine, dried
20 over anhydrous sodium sulfate. The solvent was distilled off, and the residue was purified by NH silica gel column chromatography (eluent; ethyl acetate: methanol = 98:2). The crystals were precipitated from diethyl ether, filtered off, and
25 dried under aeration to yield the title compound as white crystals (197.0 mg, 0.44 mmol, 60.7%).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.08-1.19 (5H, m),
1.30 (1H, m), 1.54 (2H, m), 1.75 (2H, m), 2.09 (3H, m),
2.70 (2H, m), 2.98 (2H, m), 3.20-3.40 (2H, m), 6.49 (1H,
dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 6.85 (1H,
5 s), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, d, J=3.6
Hz), 7.90 (1H, d, J=2.4 Hz), 8.02 (1H, d, J=5.6 Hz),
8.08 (1H, m), 8.22 (1H, m), 8.28 (1H, d, J=8.8 Hz),
9.00 (1H, s).

10 Example 61

N1-Ethyl-5-(2-((2-
(diethylamino)ethyl)amino)carbonyl)amino-4-pyridyl)oxy-
1H-1-indolecarboxamide

[0267] Similarly to Example 27, the title compound
15 (140.9 mg, 0.32 mmol, 89.2%) was obtained as white
crystals from phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-
indolyl)oxy-2-pyridyl)carbamate (150 mg, 0.36 mmol,
Production example 27-2) and 2-(diethylamino)ethylamine.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.93 (6H, t, J=7.2
20 Hz), 1.17 (3H, t, J=7.2 Hz), 2.40-2.49 (6H, m), 3.13
(2H, m), 3.20-3.40 (2H, m), 6.49 (1H, dd, J=2.4, 5.6
Hz), 6.67 (1H, d, J=3.6 Hz), 6.82 (1H, s), 7.03 (1H, dd,
J=2.4, 8.8 Hz), 7.36 (1H, d, J=2.4 Hz), 7.90 (1H, d,
J=3.6 Hz), 8.00 (1H, d, J=5.6 Hz), 8.20-8.25 (2H, m),
25 8.28 (1H, d, J=8.8 Hz), 9.11 (1H, s).

Example 62

N1-Ethyl-5-(2-(((2-(morpholin-4-yl)ethyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

5 [0268] Similarly to Example 27, the title compound (155.0 mg, 0.34 mmol, 95.1%) was obtained as white crystals from phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate (150 mg, 0.36 mmol, Production example 27-2) and 4-(2-aminoethyl)morpholine.

10 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.67 (3H, t, J=7.2 Hz), 2.30-2.40 (6H, m), 3.20 (2H, m), 3.20-3.40 (2H, m), 3.54-3.57 (4H, m), 6.50 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 6.84 (1H, s), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, d, J=3.6 Hz), 7.90 (1H, d, J=2.4 Hz),

15 8.02 (1H, d, J=5,6 Hz), 8.10-8.25 (2H, m), 8.28 (1H, d, J=8.8 Hz), 9.11 (1H, s).

Example 63

N1-Ethyl-5-(2-(((2-(4-hydroxypiperidino)ethyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

20 [0269] Similarly to Example 27, the title compound (49.1 mg, 0.11 mmol, 35.1%) was obtained as white crystals from phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate (125 mg, 0.30 mmol, Production example 27-2) and 1-(2-aminoethyl)-4-

25

hydroxypiperidine dihydrochloride.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.17 (3H, t, J=7.2 Hz), 1.36 (2H m), 1.66-1.70 (2H, m), 2.00 (2H, m), 2.32 (2H, m), 2.65-2.69 (2H, m), 3.16 (2H, m), 3.20-3.40 (2H, m), 3.40 (1H, m), 4.53 (1H, d, J=4.0 Hz), 6.50 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 6.83 (1H, s), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, d, J=2.4 Hz), 7.90 (1H, d, J=3.6 Hz), 8.01 (1H, d, J=5.6 Hz), 8.10-8.23 (2H, m), 8.28 (1H, d, J=8.8 Hz), 9.11 (1H, s).

Example 64

N1-Methyl-5-(2-((2-(4-hydroxypiperidino)ethyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0270] Similarly to Example 27, the title compound (114.3 mg, 0.25 mmol, 25.3%) was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate (402 mg, 1.0 mmol, Production example 29-1) and 1-(2-aminoethyl)-4-hydroxypiperidine dihydrochloride.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.32-1.38 (2H, m), 1.60-1.70 (2H, m), 1.96-2.03 (2H, m), 2.31-2.34 (2H, m), 2.60-2.70 (2H, m), 2.83 (3H, d, J=4.4 Hz), 3.15-3.18 (2H, m), 3.42 (1H, m), 4.53 (1H, d, J=4.0 Hz), 6.51 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 6.84 (1H, s), 7.04 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, d, J=2.4

Hz), 7.87 (1H, d, J=3.6 Hz), 8.01 (1H, d, J=5.6 Hz),
8.14-8.16 (2H, m), 8.28 (1H, d, J=8.8 Hz), 9.11 (1H, s).

Example 65

5 N1-Ethyl-5-(2-((3-
(diethylamino)propylamino)carbonyl)amino-4-pyridyl)oxy-
1H-1-indolecarboxamide

[0271] Similarly to Example 27, the title compound
(159.9 mg, 0.35 mmol, 98.1%) was obtained as white
10 crystals from phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-
indolyl)oxy-2-pyridyl)carbamate (150 mg, 0.36 mmol,
Production example 27-2) and 3-
(diethylamino)propylamine.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.91 (6H, t, J=7.2
15 Hz), 1.17 (3H, t, J=7.2 Hz), 1.50 (2H, m), 2.32-2.41
(6H, m), 3.10 (2H, m), 3.20-3.40 (2H, m), 6.50 (1H, dd,
J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 6.81 (1H, s),
7.03 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, d, J=3.6 Hz),
7.90 (1H, d, J=2.4 Hz), 8.00 (1H, d, J=5.6 Hz), 8.12
20 (1H, m), 8.22 (1H, t, J=5.2 Hz), 8.28 (1H, d, J=8.8 Hz),
9.03 (1H, s).

Example 66

25 N1-Ethyl-5-(2-((3-(morpholin-4-
yl)propyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-
indolecarboxamide

[0272] Similarly to Example 27, the title compound (135.0 mg, 0.29 mmol, 96.4%) was obtained as white crystals from phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate (125 mg, 0.30 mmol, Production example 27-2) and 4-(3-aminopropyl)morpholine.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.17 (3H, t, J=7.2 Hz), 1.55 (2H, m), 2.20-2.40 (6H, m), 3.11 (2H, m), 3.20-3.40 (2H, m), 3.51-3.55 (4H, m), 6.50 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 6.84 (1H, s), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, d, J=2.4 Hz), 7.90 (1H, d, J=3.6 Hz), 8.01 (1H, d, J=5.6 Hz), 8.04 (1H, m), 8.21 (1H, t, J=5.6 Hz), 8.28 (1H, d, J=8.8 Hz), 9.02 (1H, s).

Example 67

N1-Ethyl-5-(2-(((3-(4-methylpiperazin-1-yl)propyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0273] Similarly to Example 27, the title compound (141.9 mg, 0.30 mmol, 98.6%) was obtained as white crystals from phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate (125 mg, 0.30 mmol, Production example 27-2) and 1-(3-aminopropyl)-4-methylpiperazine.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.17 (3H, t, J=7.2

Hz), 1.54 (2H, m), 2.11 (3H, s), 2.11-2.40 (10H, m),
 3.08 (2H, m), 3.20-3.40 (2H, m), 6.50 (1H, dd, J=2.4,
 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 6.84 (1H, s), 7.03 (1H,
 dd, J=2.4, 8.8 Hz), 7.36 (1H, d, J=2.4 Hz), 7.90 (1H, d,
 5 J=3.6 Hz), 8.01 (1H, d, J=5.6 Hz), 8.04 (1H, m), 8.22
 (1H, t, J=5.6 Hz), 8.28 (1H, d, J=8.8 Hz), 9.01 (1H, s).

Example 68

N1-Cyclopropyl-5-(2-(((4-(pyrrolidin-1-yl)piperidin-1-
 10 yl)carbonyl)amino)pyridin-4-yl)oxy)-1H-1-
 indolecarboxamide

[0274] Tetrahydrofuran (30 ml) and triethylamine (3.87
 ml, 27.8 mmol) were added to N1-cyclopropyl-5-(2-amino-
 4-pyridyl)oxy-1H-1-indolecarboxamide (2.85 g, 9.25 mmol,
 15 CAS No. 417722-12-4) which was described in WO
 02/32872; phenyl chloroformate (2.57 ml, 20.4 mmol) was
 added thereto at 0 °C while stirred; and the reaction
 mixture was stirred at room temperature for 2 hours.
 The reaction mixture was partitioned between ethyl
 20 acetate and water; and the organic layer was
 concentrated to yield 3.30 g of the mixture of phenyl
 N-(4-(1-(cyclopropylamino)carbonyl-1H-5-indolyl)-oxy-2-
 pyridyl)carbamate and phenyl N-(4-(1-
 cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)-N-
 25 (phenoxycarbonyl)carbamate. A portion of 0.524 g of
 the mixture was dissolved in N,N-dimethylformamide (5

ml); 4-(1-pyrrolidinyl)piperidine (0.736 g, 4.80 mmol) was added thereto; the reaction mixture was stirred for 5 hours; the reaction mixture was partitioned between ethyl acetate and water; and the organic layer was concentrated to yield the title compound as white crystals (280 mg, 0.57 mmol).

MS Spectrum (ESI): 489 (M+1).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.57-0.75 (4H, m), 1.18-1.30 (2H, m), 1.58-1.80 (6H, m), 2.03-2.12 (1H, m), 2.38-2.48 (4H, m), 2.72-2.87 (3H, m), 3.88-3.96 (2H, m), 6.53 (1H, dd, J=2.7, 6.1 Hz), 6.64 (1H, d, J=3.4 Hz), 7.03 (1H, dd, J=2.7, 8.9 Hz), 7.30 (1H, d, J=2.7 Hz), 7.35 (1H, d, J=2.7 Hz), 7.86 (1H, d, J=3.4 Hz), 8.06 (1H, d, J=6.1 Hz), 8.24-8.29 (2H, m), 9.08 (1H, s).

Example 69

5-(2-(3-((1R)-1-Hydroxymethyl-2-oxo-2-pyrrolidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid cyclopropylamide

[0275] Similarly to Example 5, the title compound (113 mg, 0.229 mmol) was obtained as white crystals from a mixture (165 mg) of phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)-N-(phenoxycarbonyl)carbamate and phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate, intermediates in Example 68, and

(2R)-2-amino-3-hydroxy-1-(pyrrolidin-1-yl)propan-1-one
(265 mg, 1.67 mmol, synthesized as an intermediate in
Example 55).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.58-0.66 (2H, m),
5 0.70-0.78 (2H, m), 1.72-1.90 (4H, m), 2.78 (1H, m),
3.20-3.60 (6H, m), 4.54 (1H, m), 4.98 (1H, t, J=5.6 Hz),
6.53 (1H, dd, J=2.0, 6.0 Hz), 6.67 (1H, d, J=3.6 Hz)
6.97 (1H, d, J=2.0 Hz), 7.06 (1H, dd, J=2.4, 8.8 Hz),
7.37 (1H, d, J=2.4 Hz), 7.88 (1H, d, J= 3.6 Hz), 8.05
10 (1H, d, J=6.0 Hz), 8.16 (1H, brs), 8.25-8.34 (2H, m),
9.18 (1H, s).

Example 70

5-(2-(3-((1S)-1-Hydroxymethyl-2-oxo-2-pyrrolidin-1-
15 ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic
acid cyclopropylamide

[0276] Similarly to Example 5, the title compound (117
mg, 0.237 mmol) was obtained as white crystals from a
mixture (165 mg) of phenyl N-(4-(1-
20 cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)-N-
(phenoxycarbonyl)carbamate and phenyl N-(4-(1-
cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-
pyridyl)carbamate, intermediates in Example 68, and
(2S)-2-amino-3-hydroxy-1-(pyrrolidin-1-yl)propan-1-one
25 hydrochloride (synthesized as an intermediate in
Example 18).

Example 71

5-(2-(3-(2-Oxo-2-(pyrrolidin-1-yl)ethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid cyclopropylamide

5 [0277] Similarly to Example 5, the title compound (90.9 mg, 0.197 mmol) was obtained as white crystals from a mixture (165 mg) of phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)-N-(phenoxy-carbonyl)carbamate and phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate, intermediates in Example 68, and 2-amino-1-(pyrrolidin-1-yl)ethanone hydrochloride (247 mg, 1.50 mmol, synthesized as an intermediate in Example 7).
10 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.58-0.66 (2H, m), 0.71-0.79 (2H, m), 1.72-1.80 (2H, m), 1.83-1.91 (2H, m), 2.78 (1H, m), 3.28-3.40 (4H, m), 3.89 (2H, d, J=4.4 Hz), 6.54 (1H, dd, J=2.0, 6.0 Hz), 6.67 (1H, d, J=3.6 Hz), 6.94 (1H, d, J=2.0 Hz), 7.06 (1H, dd, J=2.4, 8.8 Hz), 7.38 (1H, d, J=2.4 Hz), 7.88 (1H, d, J=3.6 Hz), 8.05
15 (1H, d, J=6.0 Hz), 8.17 (1H, brs), 8.26-8.35 (2H, m), 9.28 (1H, s).

Example 72

25 5-(2-(3-(3-Oxo-3-(pyrrolidin-1-yl)propyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid cyclopropylamide

[0278] Similarly to Example 5, the title compound (113 mg, 0.237 mmol) was obtained as white crystals from a mixture (165 mg) of phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)-N-(phenoxy-carbonyl)carbamate and phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate, intermediates in Example 68, and 3-amino-1-(pyrrolidin-1-yl)propan-1-one hydrochloride (268 mg, 1.50 mmol, synthesized as an intermediate in Example 25).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.58-0.66 (2H, m), 0.71-0.79 (2H, m), 1.70-1.79 (2H, m), 1.79-1.88 (2H, m), 2.40 (2H, t, J=6.4 Hz), 2.78 (1H, m), 3.24-3.38 (6H, m), 6.51 (1H, dd, J=2.0, 6.0 Hz), 6.67 (1H, d, J=3.8 Hz), 6.93 (1H, d, J=2.0 Hz), 7.05 (1H, dd, J=2.4, 8.8 Hz), 7.37 (1H, d, J=2.4 Hz), 7.88 (1H, d, J=3.8 Hz), 7.98-8.10 (2H, m), 8.26-8.34 (2H, m), 9.09 (1H, s).

Example 73

5-(2-(3-((1R)-1-Hydroxymethyl-2-oxo-2-piperidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid cyclopropylamide

[0279] Similarly to Example 5, the title compound (106 mg, 0.209 mmol) was obtained as white crystals from a mixture (165 mg) of phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)-N-

(phenoxycarbonyl) carbamate and phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl) carbamate, intermediates in Example 68, and (2R)-2-amino-3-hydroxy-1-(piperidin-1-yl)propan-1-one (228 mg, 1.32 mmol, synthesized as an intermediate in Example 57).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.58-0.66 (2H, m), 0.70-0.78 (2H, m), 1.38-1.62 (6H, m), 2.79 (1H, m), 3.38-3.53 (6H, m), 4.75 (1H, m), 4.93 (1H, t, J=5.8 Hz), 6.54 (1H, dd, J=2.0, 6.0 Hz), 6.67 (1H, d, J=3.6 Hz), 6.97 (1H, d, J=2.0 Hz), 7.06 (1H, dd, J=2.4, 8.8 Hz), 7.37 (1H, d, J=2.4 Hz), 7.88 (1H, d, J=3.6 Hz), 8.05 (1H, d, J=6.0 Hz), 8.10-8.34 (3H, m), 9.20 (1H, s).

Example 74

5-(2-(3-((1S)-1-Hydroxymethyl-2-oxo-2-piperidin-1-ylethyl)ureido)pyridin-4-yloxy)-1H-indole-1-carboxylic acid cyclopropylamide

[0280] Similarly to Example 5, the title compound (66.8 mg, 0.132 mmol) was obtained as white crystals from a mixture (82.3 mg) of phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)-N-(phenoxycarbonyl) carbamate and phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl) carbamate, intermediates in Example 68, and (2S)-2-amino-3-hydroxy-1-(piperidin-1-yl)propan-1-one

hydrochloride (156 mg, 0.748 mmol, synthesized as an intermediate in Example 20).

Example 75

5 N1-Phenyl-5-(2-(((3-(diethylamino)propyl)amino)carbonyl)amino-4-pyridyl)-oxy-1H-1-indolecarboxamide

[0281] The title compound was obtained from N1-phenyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (CAS
10 No. 417721-87-0) which was written in the description of WO 02/32872 and 3-diethylaminopropylamine using a procedure analogous to that described for Example 28.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.91 (6H, t, J=7.2 Hz), 1.47-1.53 (2H, m), 2.30-2.44 (6H, m), 3.05-3.14
15 (2H, m), 6.52 (1H, dd, J=6.0, 2.0 Hz), 6.76 (1H, d, J=3.6 Hz), 6.84 (1H, d, J=2.0 H), 7.09 (1H, dd, J=9.2, 2.4 Hz), 7.13 (1H, t, J=7.6 Hz), 7.38 (2H, dd, J=7.6, 7.6 Hz), 7.42 (1H, d, J=2.4 Hz), 7.64 (2H, d, J=7.6 Hz),
8.02 (1H, d, J=6.0 Hz), 8.10-8.14 (2H, m), 8.27 (1H, d, J=9.2 Hz),
20 9.05 (1H, brs), 10.10 (1H, brs).

Example 76

N1-Phenyl-5-(2-(((3-(4-methylpiperazin-1-yl)propyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide
25

[0282] Similarly to Example 28, the title compound was

obtained from N1-phenyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (CAS No. 417721-87-0) which was described in WO 02/32872 and 1-(3-aminopropyl)-4-methylpiperazine.

5 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.52-1.59 (2H, m),
2.13 (3H, s), 2.15-2.45 (10H, m), 3.08-3.15 (2H, m),
6.54 (1H, dd, J=6.0, 2.0 Hz), 6.79 (1H, d, J=3.6 Hz),
6.89 (1H, brs), 7.10 (1H, dd, J=2.4, 9.2 Hz), 7.15 (1H,
t, J=7.6 Hz), 7.40 (2H, t, J=7.6 Hz), 7.44 (1H, d,
10 J=2.4 Hz), 7.66 (2H, d, J=7.6 Hz), 8.03-8.07 (2H, m),
8.14 (1H, d, J=3.6 Hz), 8.29 (1H, d, J=9.2 Hz), 9.05
(1H, brs), 10.10 (1H, brs).

Example 77

15 N1-Ethyl-5-(2-(((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide

[0283] Tetrahydrofuran (20 ml) and triethylamine (2.70 ml, 19.4 mmol) were added to N1-ethyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (1.91 g, 6.45 mmol, Production example 27-1); phenyl chloroformate (1.79 ml, 14.2 mmol) was added thereto at 0 °C while stirred; and the reaction mixture was stirred at room temperature for 2 hours. The reaction mixture was partitioned
25 between ethyl acetate and water; the organic layer was concentrated to yield a mixture (2.95 g) of phenyl N-

(4-(1-(ethylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate and phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-N-(phenoxycarbonyl)carbamate. A portion of 0.454 g of the mixture was dissolved in N,N-dimethylformamide (5 ml); and 4-(1-pyrrolidinyl)piperidine (0.522 g, 3.39 mmol) was added; and the reaction mixture was stirred for 5 hours. The reaction mixture was partitioned between ethyl acetate and water; the organic layer was concentrated to yield a solid; the obtained solid was washed with hexane: diethyl ether=1: 1 to yield the title compound as crystals (205 mg, 0.43 mmol).

MS Spectrum (ESI): 477 (M+1), 953 (2M+1).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.12-1.22 (5H, m), 1.57-1.81 (6H, m), 2.05-2.15 (1H, m), 2.38-2.50 (4H, m), 2.77-2.78 (2H, m), 3.28-3.37 (2H, m), 3.87-3.97 (2H, m), 6.53 (1H, dd, J=2.5, 5.4 Hz), 6.66 (1H, d, J=3.5 Hz), 7.02 (1H, dd, J=2.5, 8.9 Hz), 7.30 (1H, d, J=2.5 Hz), 7.36 (1H, d, J=2.5 Hz), 7.89 (1H, d, J=3.5 Hz), 8.05 (1H, d, J=5.4 Hz), 8.20 (1H, m), 8.27 (1H, t, J=8.9 Hz), 9.08 (1H, s).

Example 78

5-(2-(((4-Hydroxy-4-methylpiperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-indole-1-carboxylic acid ethylamide

[0284] Similarly to Example 41, the title compound was obtained as colorless crystals (124 mg, 0.283 mmol, 89.4%) from 4-hydroxy-4-methylpiperidine monohydrochloride (216 mg, 1.42 mmol, Production example 8-3) and phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (170 mg, 0.317 mmol, Production example 55-1).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.08 (3H, s), 1.17 (3H, t, J=7.2 Hz), 1.38-1.44 (4H, m), 3.13 (2H, m), 3.30 (2H, m), 3.63 (2H, m), 4.27 (1H, s), 6.53 (1H, dd, J=2.4, 6.0 Hz), 6.67 (1H, d, J=3.6 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.32 (1H, d, J=2.4 Hz), 7.35 (1H, d, J=2.4 Hz), 7.90 (1H, d, J=3.0 Hz), 8.05 (1H, d, J=6.0 Hz), 8.21 (1H, t, J=5.4 Hz), 8.27 (1H, d, J=8.8 Hz), 9.04 (1H, s).

Example 79

N1-Ethyl-5-(2-((4-hydroxypiperidin-1-yl)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0285] Similarly to Example 27, the title compound was obtained as white powder (18.7 mg, 0.044 mmol, 14.7%) from phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate (125 mg, 0.30 mmol, Production example 27-2) and 4-hydroxypiperidine.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.13-1.27 (5H, m),

1.63-1.67 (2H, m), 2.98 (2H, m), 3.20-3.40 (2H, m),
 3.60 (1H, m), 3.74 (2H, m), 4.64 (1H, d, J=4.4 Hz),
 6.53 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz),
 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.31 (1H, d, J=2.4 Hz),
 5 7.35 (1H, d, J=2.4 Hz), 7.90 (1H, d, J=3.6 Hz), 8.06
 (1H, d, J=5.6 Hz), 8.21 (1H, t, J=5.2 Hz), 8.27 (1H, d,
 J=8.8 Hz), 9.09 (1H, s).

Example 80

10 N1-Ethyl-5-(2-(piperidin-1-ylcarbonyl)amino-4-
pyridyl)oxy-1H-1-indolecarboxamide

[0286] N,N-Dimethylformamide (4 ml) and piperidine
 (0.31 ml, 3.13 mmol) were added to a mixture (0.336 g)
 of phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-
 15 indolyl)oxy-2-pyridyl)carbamate and phenyl N-(4-(1-
 (ethylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-N-
 (phenoxy carbonyl)carbamate obtained in Example 77; the
 reaction mixture was stirred overnight; the reaction
 mixture was partitioned between ethyl acetate and
 20 water; and the organic layer was concentrated to yield
 the title compound as crystals (182 mg, 0.45 mmol).

MS Spectrum (ESI): 408 (M+1), 815 (2M+1).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.18 (3H, t, J=7.6
 Hz), 1.35-1.57 (6H, m), 3.23-3.33 (6H, m), 6.52 (1H, dd,
 25 J=2.4, 5.4 Hz), 6.67 (1H, d, J=3.4 Hz), 7.03 (1H, dd,
 J=2.4, 8.7 Hz), 7.30 (1H, d, J=2.4 Hz), 7.36 (1H, d,

J=2.4 Hz), 7.90 (1H, d, J=3.4 Hz), 8.06 (1H, d, J=5.5 Hz), 8.21 (1H, t, J=5.5 Hz), 8.27 (1H, d, J=8.7 Hz), 9.05 (1H, s).

5 Example 81

N1-Ethyl-5-((2-((pyrrolidin-1-ylcarbonyl)amino)-4-pyridyl)oxy)-1H-1-indolecarboxamide

[0287] N,N-Dimethylformamide (5 ml) and pyrrolidine (0.36 ml, 4.3 mmol) were added to a mixture (0.461 g) of phenyl N-(4-(1-(ethylamino)carbonyl)-1H-5-indolyl)oxy-2-pyridyl)carbamate and phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyl)oxy)-2-pyridyl)-N-(phenoxy carbonyl)carbamate obtained in Example 77; the reaction mixture was stirred overnight; the reaction mixture was partitioned between ethyl acetate and water; and the organic layer was concentrated to yield the title compound as crystals (245 mg, 0.623 mmol).

MS Spectrum (ESI): 394 (M+1), 787 (2M+1).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.16 (3H, t, J=7.6 Hz), 1.70-1.82 (4H, m), 3.22-3.40 (6H, m), 6.54 (1H, dd, J=2.4, 5.5 Hz), 6.67 (1H, d, J=3.4 Hz), 7.03 (1H, dd, J=2.4, 8.7 Hz), 7.35 (1H, d, J=2.4 Hz), 7.41 (1H, d, J=2.4 Hz), 7.90 (1H, d, J=3.4 Hz), 8.06 (1H, d, J=5.5 Hz), 8.21 (1H, t, J=5.5 Hz), 8.27 (1H, d, J=8.7 Hz), 8.59 (1H, s).

Example 82

N4-(4-((1-(Ethylamino)carbonyl-1H-5-indolyl)oxy)-2-pyridyl)-4-morpholinecarboxamide

[0288] N,N-Dimethylformamide (5 ml) and morpholine
5 (0.326 ml, 3.73 mmol) were added to a mixture (0.401 g) of phenyl N-(4-(1-(ethylamino)carbonyl)-1H-5-indolyl)oxy-2-pyridyl)carbamate and phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyl)oxy)-2-pyridyl)-N-(phenoxy-carbonyl)carbamate obtained in Example 77; the
10 reaction mixture was stirred overnight; the reaction mixture was partitioned between ethyl acetate and water; the organic layer was concentrated; and the obtained solid was washed with a solvent mixture of hexane: diethyl ether = 1: 1 to yield the title
15 compound (255 mg, 0.62 mmol).

MS Spectrum (ESI): 410 (M+1), 819 (2M+1).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.17 (3H, t, J=7.7 Hz), 3.25-3.42 (6H, m), 3.48-3.53 (4H, m), 6.55 (1H, dd, J=2.6, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 7.02 (1H, dd, J=2.6, 8.7 Hz), 7.29 (1H, d, J=2.6 Hz), 7.35 (1H, d, J=2.4 Hz), 7.90 (1H, d, J=3.6 Hz), 8.08 (1H, d, J=5.6 Hz), 8.20 (1H, t, J=5.6 Hz), 8.28 (1H, t, J=5.6 Hz), 9.19 (1H, s).

20

25 Example 83

N1-Ethyl-5-(2-((1,1-dioxothiomorpholin-4-

ylcarbonyl)amino)pyridin-4-yloxy)-1H-1-
indolecarboxamide

[0289] Similarly to Example 54, the title compound was obtained as colorless crystals (116 mg, 0.253 mmol, 80.0%) from 1,1-dioxothiomorpholine hydrochloride (248 mg, 1.42 mmol, Production example 54-3) and phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxy carbonyl) carbamate (170 mg, 0.317 mmol, Production example 55-1).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.17 (3H, t, J=7.2 Hz), 3.10 (4H, m), 3.29 (2H, m), 3.80 (4H, m), 6.58 (1H, dd, J=2.0, 5.6 Hz), 6.67 (1H, d, J=3.4 Hz), 7.03 (1H, dd, J=2.0, 9.0 Hz), 7.31 (1H, d, J=2.0 Hz), 7.36 (1H, d, J=2.0 Hz), 7.90 (1H, d, J=3.4 Hz), 8.10 (1H, d, J=5.6 Hz), 8.22 (1H, t, J=5.4 Hz), 8.28 (1H, d, J=9.0 Hz), 9.54 (1H, s).

Example 84

N1-Ethyl-5-(2-((methoxylamino)carbonyl)amino-4-
pyridyl)oxy-1H-1-indolecarboxamide

[0290] Similarly to Example 27, the title compound was obtained as white crystals (94.3 mg, 0.26 mmol, 70.9%) from phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolylyl)oxy-2-pyridinyl) carbamate (150 mg, 0.36 mmol, Production example 27-2) and methoxylamine hydrochloride.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.17 (3H, t, J=7.2 Hz), 3.20-3.40 (2H, m), 3.59 (3H, s), 6.57 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 7.05 (1H, dd, J=2.4, 8.8 Hz), 7.16 (1H, s), 7.38 (1H, d, J=2.4 Hz), 7.90 (1H, d, J=3.6 Hz), 8.08 (1H, d, J=5.6 Hz), 8.21 (1H, m), 8.28 (1H, d, J=8.8 Hz), 8.95 (1H, s), 10.15 (1H, s).

Example 85

N1-Cyclopropyl-5-(2-((4-hydroxypiperidino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0291] N,N-Dimethylformamide (5 ml) and 4-hydroxypiperidine (433 mg, 4.29 mmol) were added to a mixture (470 mg) of phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)-N-(phenoxycarbonyl)carbamate and phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate obtained in Example 68; the reaction mixture was stirred overnight; the reaction mixture was partitioned between ethyl acetate and water; and the organic layer was concentrated to yield the title compound as white crystals (220 mg, 0.51 mmol, 39%).

MS Spectrum (ESI): 436 (M+1), 871 (2M+1).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.58-0.63 (2H, m), 0.69-0.76 (2H, m), 1.18-1.30 (2H, m), 1.60-1.70 (2H, m),

2.70-2.80 (1H, m), 2.93-3.02 (2H, m), 3.55-3.64 (1H, m),
 3.69-3.77 (2H, m), 4.63 (1H, d, J=4.4 Hz), 6.53 (1H, dd,
 J=2.4, 5.8 Hz), 6.64 (1H, d, J=3.6 Hz), 7.04 (1H, dd,
 J=2.4, 8.5 Hz), 7.31 (1H, d, J=2.4 Hz), 7.35 (1H, d,
 5 J=2.4 Hz), 7.86 (1H, d, J=3.6 Hz), 8.06 (1H, d, J=5.8
 Hz), 8.24-8.29 (2H, m), 9.08 (1H, s).

Example 86

10 N1-Cyclopropyl-5-(2-(((4-hydroxy-4-methylpiperidin-1-
 yl)carbonyl)amino)pyridin-4-yl)oxy)-1H-1-
 indolecarboxamide

[0292] Similarly to Example 41, the title compound was
 obtained as colorless crystals (109 mg, 0.242 mmol)
 from 4-hydroxy-4-methylpiperidine monohydrochloride
 15 (221 mg, 1.46 mmol, Production example 8-3) and a
 mixture (200 mg, intermediates in Example 68) of
 phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-
 indolyl)oxy-2-pyridyl)-N-(phenoxycarbonyl)carbamate and
 phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-
 20 indolyl)oxy-2-pyridyl)carbamate.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.61 (2H, m), 0.73
 (2H, m), 1.08 (3H, s), 1.30-1.41 (4H, m), 2.76 (1H, m),
 3.14 (2H, m), 3.63 (2H, m), 4.27 (1H, s), 6.53 (1H, d,
 J=5.4 Hz), 6.65 (1H, d, J=3.4 Hz), 7.03 (1H, d, J=8.8
 25 Hz), 7.32 (1H, s), 7.35 (1H, s), 7.86 (1H, d, J=3.4 Hz),
 8.06 (1H, d, J=5.4 Hz), 8.27 (2H, m), 9.04 (1H, s).

Example 87N4-(4-(1-(Cyclopropylamino)carbonyl)-1H-5-indolyl)oxy-2-pyridyl)-4-morpholinecarboxamide

5 [0293] N,N-Dimethylformamide (5 ml) and morpholine (0.373 ml, 4.28 mmol) were added to a mixture (470 mg) of phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)-N-(phenoxy-carbonyl)carbamate and phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate obtained in Example 68; and the reaction mixture was stirred overnight; the reaction mixture was partitioned between ethyl acetate and water; the organic layer was concentrated; and the obtained solid was washed with a solvent mixture of
10 hexane: diethyl ether = 1: 1 to yield the title compound (255 mg, 0.58 mmol, 95%).

MS Spectrum (ESI): 422 (M+1), 843 (2M+1).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.58-0.75 (4H, m), 2.72-2.81 (1H, m), 3.26-3.40 (4H, m), 3.50 (4H, t, J=4.8 Hz), 6.56 (1H, dd, J=2.5, 5.6 Hz), 6.65 (1H, d, J=3.4 Hz), 7.04 (1H, dd, J=2.5, 8.8 Hz), 7.30 (1H, d, J=2.5 Hz), 7.36 (1H, d, J=2.5 Hz), 7.86 (1H, d, J=3.4 Hz), 8.08 (1H, d, J=5.5 Hz), 8.24-8.30 (2H, m), 9.18 (1H, s).

25

Example 88

N1-Cyclopropyl-5-(2-((pyrrolidin-1-ylcarbonyl)amino)-4-pyridyl)oxy-1H-1-indolecarboxamide

[0294] N,N-Dimethylformamide (5 ml) and pyrrolidine (0.35 ml, 4.2 mmol) were added to a mixture (470 mg) of phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)-N-(phenoxycarbonyl)carbamate and phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate obtained in Example 68; the reaction mixture was stirred overnight; the reaction mixture was partitioned between ethyl acetate and water; and the organic layer was concentrated to yield the title compound as white crystals (200 mg, 0.49 mmol).

MS Spectrum (ESI): 406 (M+1), 811 (2M+1).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.58-0.78 (4H, m), 1.70-1.83 (4H, m), 2.73-2.81 (1H, m), 3.23-3.45 (4H, m), 6.55 (1H, dd, J=2.2, 5.7 Hz), 6.65 (1H, d, J=3.5 Hz), 7.03 (1H, dd, J=2.2, 8.7 Hz), 7.36 (1H, d, J=2.2 Hz), 7.41 (1H, d, J=2.2 Hz), 7.86 (1H, d, J=3.5 Hz), 8.06 (1H, d, J=5.7 Hz), 8.16-8.30 (2H, m), 8.59 (1H, s).

Example 89

N1-Cyclopropyl-5-(2-(piperidin-1-ylcarbonyl)amino)-4-pyridyl)oxy-1H-1-indolecarboxamide

[0295] N,N-Dimethylformamide (5 ml) and piperidine (0.42 ml, 4.2 mmol) were added to a mixture (467 mg) of

phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)-N-(phenoxycarbonyl)carbamate and phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate obtained in Example 68; and the reaction mixture was stirred overnight; the reaction mixture was partitioned between ethyl acetate and water; the organic layer was concentrated; the obtained solid was washed with a solvent mixture of hexane: diethyl ether = 1: 1 to yield the title compound as crystals (241 mg, 0.57 mmol).

MS Spectrum (ESI): 420 (M+1), 839 (2M+1).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.58-0.77 (4H, m), 1.34-1.55 (6H, m), 2.72-2.81 (1H, m), 3.27-3.40 (4H, m), 6.52 (1H, dd, J=2.6, 5.6 Hz), 6.64 (1H, d, J=3.6 Hz), 7.03 (1H, dd, J=2.6, 8.7 Hz), 7.30 (1H, d, J=2.6 Hz), 7.35 (1H, d, J=2.6 Hz), 7.87 (1H, d, J=3.6 Hz), 8.06 (1H, d, J=5.6 Hz), 8.23-8.30 (2H, m), 9.03 (1H, s).

Example 90

N4-(4-(1-(Cyclopentylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-4-morpholinecarboxamide

[0296] Phenyl N-(4-(1-cyclopentylaminocarbonyl-1H-indol-5-yloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (200 mg, 0.35 mmol) was dissolved in N,N-dimethylformamide (1.5 ml) and morpholine (0.15 ml, 1.73 mmol); and the reaction mixture was stirred at

room temperature overnight. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Fuji Silysia BW-300; ethyl acetate, ethyl acetate: methanol = 10: 1 in this order); the obtained colorless oil was crystallized by addition of diethyl ether; and the crystals were filtered off, washed with diethyl ether, and dried under aeration to yield the title compound as colorless crystals (140 mg, 0.31 mmol, 90%).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.48-1.64 (4H, m), 1.66-1.76 (2H, m), 1.88-1.98 (2H, m), 3.35 (4H, m), 3.51 (4H, m), 4.14 (1H, m), 6.56 (1H, d, J=6.0 Hz), 6.65 (1H, d, J=3.6 Hz), 7.02 (1H, d, J=8.8 Hz), 7.30 (1H, s), 7.35 (1H, s), 7.96 (1H, d, J=3.6 Hz), 8.00 (1H, d, J=6.8 Hz), 8.08 (1H, d, J=6.0 Hz), 8.25 (1H, d, J=8.8 Hz), 9.18 (1H, s).

[0297] The starting materials were synthesized as follows.

Production example 90-1

Phenyl N-cyclopentylcarbamate

Cyclopentylamine (9.9 ml, 100 mmol) was dissolved in tetrahydrofuran (400 ml); pyridine (8.9 ml,

110 mmol) was added thereto; and the reaction mixture was stirred. The reaction mixture was cooled with ice; phenyl chloroformate (13.8 ml, 110 mmol) was added dropwise for 5 minutes while stirring; and the reaction mixture was stirred at room temperature for 24.5 hours. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained crystals were suspended in a solvent mixture of hexane: ethyl acetate = 5: 1, filtered off, washed with hexane, and dried under aeration to yield the title compound as colorless crystals (16.6 g, 81 mmol, 81%).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.47 (4H, m), 1.63 (2H, m), 1.81 (2H, m), 3.81 (1H, m), 7.07 (2H, d, J=7.6 Hz), 7.17 (1H, t, J=7.6 Hz), 7.35 (2H, t, J=7.6 Hz), 7.75 (1H, d, J=6.8 Hz).

Production example 90-2

N1-Cyclopentyl-5-(2-aminopyridin-4-yloxy)-1H-1-indolecarboxamide

[0298] 4-(1H-5-Indolyloxy)-2-pyridinamine (2.50 g, 11.1 mmol, CAS No. 417722-11-3), which was described in WO 02/32872, was dissolved in N,N-dimethylformamide (30 ml); sodium hydride (0.530 g, 13.3 mmol) was added thereto at room temperature; and the reaction mixture

was stirred for 30 minutes. Phenyl N-cyclopentylcarbamate (2.50 g, 12.2 mmol) was added thereto at room temperature while stirring; and the reaction mixture was stirred for 30 minutes. Water was
5 added to the reaction mixture; and the precipitated crystals were filtered off, and washed with water. This crystals were dissolved in methanol, and purified by silica gel column chromatography (Fuji Silysia NH; hexane: ethyl acetate = 1: 1, ethyl acetate, ethyl
10 acetate: methanol = 98: 2 in this order). The obtained crystals were suspended in hexane: ethanol = 10: 1, filtered off, washed with hexane, and dried under aeration to yield the title compound as colorless crystals (2.08 g, 6.18 mmol, 55.7%).

15 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.56 (4H, m), 1.71 (2H, m), 1.92 (2H, m), 4.14 (1H, m), 5.74 (1H, d, J=2.0 Hz), 5.83 (2H, s), 6.12 (1H, dd, J=2.0, 5.6 Hz), 6.64 (1H, d, J=3.4 Hz), 7.00 (1H, dd, J=2.0, 8.8 Hz), 7.32 (1H, d, J=2.0 Hz), 7.75 (1H, d, J=5.6 Hz), 7.94 (1H, d, J=3.4 Hz),
20 7.97 (1H, d, J=6.4 Hz), 8.23 (1H, d, J=8.8 Hz).

Production example 90-3

Phenyl N-(4-(1-cyclopentylaminocarbonyl-1H-indol-5-yloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate
25 [0299] N1-Cyclopentyl-5-(2-aminopyridin-4-yloxy)-1H-1-

indolecarboxamide (1.55 g, 4.58 mmol) was dissolved in tetrahydrofuran (90 ml); triethylamine (1.43 ml, 10.31 mmol) and pyridine (0.56 ml, 6.88 mmol) were added thereto; and the reaction mixture was stirred. The reaction mixture was cooled with ice; phenyl chloroformate (1.44 ml, 11.45 mmol) was added dropwise; and the reaction mixture was stirred at room temperature for 2.5 hours. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Fuji Silysia BW-300; hexane: ethyl acetate = 1: 1, 1: 3 in this order) to yield the title compound as a colorless amorphous solid (2.516 g, 4.36 mmol, 95.2%).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.50-1.63 (4H, m), 1.66-1.74 (2H, m), 1.88-1.98 (2H, m), 4.15 (1H, m), 6.65 (1H, d, J=3.8 Hz), 6.95 (1H, dd, J=2.4, 5.6 Hz), 7.09 (1H, dd, J=2.4, 8.8 Hz), 7.16 (4H, d, J=7.6 Hz), 7.29 (2H, d, J=7.6 Hz), 7.42 (4H, d, J=7.6 Hz), 7.44 (1H, d, J=2.4 Hz), 7.51 (1H, d, J=2.4 Hz), 7.98 (1H, d, J=3.8 Hz), 8.01 (1H, d, J=6.8 Hz), 8.28 (1H, d, J=8.8 Hz), 8.42 (1H, d, J=5.6 Hz).

Example 91

5-(2-(((4-Hydroxypiperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-indole-1-carboxylic acid cyclopentylamide

[0300] Similarly to Example 90, the title compound was obtained as colorless crystals (129 mg, 0.278 mmol, 80.2%) from phenyl N-(4-(1-cyclopentylaminocarbonyl-1H-indol-5-yloxy)-2-pyridyl)-N-(phenoxy carbonyl) carbamate (200 mg, 0.346 mmol, Production example 90-3) and 4-hydroxypiperidine (175 mg, 1.73 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.23 (2H, m), 1.48-1.77 (8H, m), 1.92 (2H, m), 2.98 (2H, m), 3.59 (1H, m), 3.73 (2H, m), 4.15 (1H, m), 4.64 (1H, d, J=4.4 Hz), 6.53 (1H, dd, J=2.0, 5.6 Hz), 6.65 (1H, d, J=3.6 Hz), 7.02 (1H, dd, J=2.0, 9.0 Hz), 7.31 (1H, d, J=2.0 Hz), 7.35 (1H, d, J=2.0 Hz), 7.96 (1H, d, J=3.6 Hz), 7.99 (1H, d, J=6.8 Hz), 8.06 (1H, d, J=5.6 Hz), 8.24 (1H, d, J=9.0 Hz), 9.09 (1H, s).

Example 92

N1-Cyclopentyl-5-(2-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-

indolecarboxamide

[0301] Similarly to Example 90, the title compound was obtained as colorless crystals (83 mg, 0.161 mmol, 46.3%) from phenyl N-(4-(1-cyclopentylaminocarbonyl-1H-indol-5-yloxy)-2-pyridyl)-N-(phenoxy carbonyl) carbamate (200 mg, 0.346 mmol, Production example 90-3) and 4-(1-pyrrolidinyl)piperidine (268 mg, 1.73 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.18-1.30 (2H, m),
 1.50-1.80 (12H, m), 1.87-1.98 (2H, m), 2.08 (1H, m),
 2.43 (4H, m), 2.81 (2H, m), 3.91 (2H, m), 4.15 (1H, m),
 6.53 (1H, dd, J=2.0, 5.6 Hz), 6.65 (1H, d, J=3.6 Hz),
 5 7.02 (1H, dd, J=2.0, 9.0 Hz), 7.31 (1H, d, J=2.0 Hz),
 7.35 (1H, d, J=2.0 Hz), 7.96 (1H, d, J=3.6 Hz), 7.99
 (1H, d, J=6.8 Hz), 8.06 (1H, d, J=5.6 Hz), 8.25 (1H, d,
 J=9.0 Hz), 9.08 (1H, s).

10 Example 93

N1-(3-Methylbutyl)-5-(2-(((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide

[0302]N,N-dimethylformamide (30 ml), pyridine (0.52 ml,
 15 6.4 mmol) and triethylamine (1.35 ml, 9.69 mmol) were
 added to N1-(3-methylbutyl)-5-((2-amino-4-pyridyl)oxy)-
 1H-1-indolecarboxamide (1.45 g, 4.29 mmol); phenyl
 chloroformate (0.81 ml, 6.4 mmol) was added at 0 °C
 while stirring; and the reaction mixture was stirred at
 20 room temperature for 1 hour. The reaction mixture was
 partitioned between ethyl acetate and water; and the
 organic layer was concentrated and subjected to silica
 gel column chromatography to yield a mixture (2.0 g) of
 phenyl N-(4-(1-((3-methylbutyl)amino)carbonyl-1H-5-
 25 indolyl)oxy-2-pyridyl)carbamate and phenyl N-(4-(1-((3-
 methylbutyl)amino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-

N-(phenoxycarbonyl)carbamate. A portion of 0.4 g of the mixture was dissolved in N,N-dimethylformamide (4 ml); 4-(1-pyrrolidinyl)piperidine (0.43 g, 2.8 mmol) was added thereto; and the reaction mixture was stirred
5 for 2 hours. The reaction mixture was partitioned between ethyl acetate and water; the organic layer was concentrated; and the residue was purified by silica gel column chromatography (Fuji Silysia NH, ethyl acetate: methanol = 10: 1) to yield the title compound
10 as white crystals (275 mg, 0.53 mmol).

MS Spectrum (ESI): 519(M+1).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.91 (6H, d, J=7.6 Hz), 1.18-1.30 (3H, m), 1.47 (2H, q, J=7.6 Hz), 1.57-1.80 (6H, m), 2.03-2.22 (1H, m), 2.37-2.48 (4H, m),
15 2.76-2.85 (2H, m), 3.25-3.36 (2H, m), 3.88-3.97 (2H, m), 6.53 (1H, dd, J=2.4, 5.4 Hz), 6.66 (1H, d, J=3.6 Hz), 7.02 (1H, dd, J=2.4, 8.7 Hz), 7.31 (1H, d, J=2.4 Hz), 7.35 (1H, d, J=2.4 Hz), 7.90 (1H, d, J=3.6 Hz), 8.06 (1H, d, J=5.4 Hz), 8.16 (1H, t, J=5.4 Hz), 8.27 (1H, d,
20 J=8.7 Hz), 9.08 (1H, s).

The starting materials were synthesized as follows.

Production example 93-1

25 N1-(3-Methylbutyl)-5-((2-amino-4-pyridyl)oxy)-1H-1-indolecarboxamide

4-(1H-5-Indolyloxy)-2-pyridinamine (2.0 g, 8.9 mmol, CAS No. 417722-11-3) which was described in WO 02/32872 was dissolved in N,N-dimethylformamide (20 ml); and sodium hydride (426 mg, 10.7 mmol) was added thereto at room temperature while stirring. The reaction mixture was cooled with ice bath after 30 minutes; phenyl N-(3-methylbutyl)carbamate (2.02 g, 9.75 mmol) was added thereto; and the reaction mixture was stirred for 3 hours at room temperature. The reaction mixture was partitioned between ethyl acetate and water; the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, and concentrated; and the residue was purified by NH-silica gel column chromatography (hexane: ethyl acetate = 3:1) to yield the title compound as crystals (1.45 g, 4.3 mmol, 48%).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.89-0.93 (6H, m), 1.40-1.70 (3H, m), 3.25-3.40 (2H, m), 5.72-5.75 (1H, m), 5.83 (2H, s), 6.10-6.40 (1H, m), 6.64-6.68 (1H, m), 6.98-7.02 (1H, m), 7.30-7.34 (1H, m), 7.75 (1H, dd, J=1.5, 6.0 Hz), 7.86-7.90 (1H, m), 8.14 (1H, t, J=4.5 Hz), 8.25 (1H, d, J=9.0 Hz).

Production example 93-2

Phenyl N-(3-methylbutyl)carbamate

[0304] Phenyl chloroformate (14.8 ml, 0.117 mol) was dissolved in tetrahydrofuran (200 ml); triethylamine

(18.0 ml, 0.129 mol) and isoamylamine (15.0 ml, 0.129 mol) were added thereto at room temperature while stirring; and the reaction mixture was stirred overnight. The reaction mixture was partitioned
 5 between ethyl acetate and water; and the organic layer was concentrated and dried under reduced pressure to yield the title compound as crystals (14 g, 0.068 mol, 58%).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.89 (6H, d, J=7.9 Hz), 1.36 (2H, q, J=7.9 Hz), 1.55-1.69 (1H, m), 3.05
 10 (2H, q, J=7.9 Hz), 7.03-7.09 (2H, m), 7.14-7.19 (1H, m), 7.31-7.38 (2H, m), 7.68 (1H, t, J=4.8 Hz).

Example 94

15 N1-(3-Methylbutyl)-5-(2-((4-hydroxypiperidino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0305] N,N-dimethylformamide (2.5 ml) and 4-hydroxypiperidine (213 mg, 2.11 mmol) were added to a
 20 mixture (243 mg) of phenyl N-(4-(1-((3-methylbutyl)amino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-carbamate and phenyl N-(4-(1-((3-methylbutyl)amino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-N-(phenoxy-carbonyl)carbamate synthesized in Example 93;
 25 and the reaction mixture was stirred for 2 hours. The reaction mixture was partitioned between ethyl acetate

and water; the organic layer was concentrated; and the residue was purified by NH-silica gel column chromatography (ethyl acetate: methanol = 10: 1) to yield the title compound as white crystals (150 mg, 0.322 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.91 (6H, d, J=7.2 Hz), 1.18-1.30 (2H, m), 1.46 (2H, q, J=7.2 Hz), 1.60-1.70 (3H, m), 2.97 (2H, m), 3.25-3.35 (2H, m), 3.55-3.64 (1H, m), 3.69-3.80 (2H, m), 4.63 (1H, d, J=3.4 Hz), 6.53 (1H, dd, J=2.3, 5.8 Hz), 6.66 (1H, d, J=3.5 Hz), 7.02 (1H, dd, J=2.3, 8.6 Hz), 7.31 (1H, d, J=2.3 Hz), 7.35 (1H, d, J=2.3 Hz), 7.90 (1H, d, J=3.5 Hz), 8.06 (1H, d, J=5.8 Hz), 8.16 (1H, t, J=5.8 Hz), 8.26 (1H, t, J=8.6 Hz), 9.08 (1H, s).

Example 95

N4-(4-(1-((3-Methylbutyl)amino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-4-morpholinecarboxamide

[0306] N,N-dimethylformamide (5 ml) and morpholine (0.163 ml, 1.87 mmol) were added to a mixture (0.6 g) of phenyl N-(4-(1-((3-methylbutyl)amino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate and phenyl N-(4-(1-((3-methylbutyl)amino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-N-(phenoxy-carbonyl)carbamate synthesized in Example 93; and the reaction mixture was stirred for 2 hours. The reaction mixture was partitioned between ethyl acetate

and water; the organic layer was concentrated; and the residue was purified by silica gel column chromatography (Fuji Silysia NH, ethyl acetate: methanol = 10: 1) to yield the title compound as white crystals (0.202 g, 0.447 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.92 (6H, dd, J=1.7, 7.3 Hz), 1.47 (2H, q, J=7.3 Hz), 1.58-1.70 (1H, m), 3.25-3.60 (10H, m), 6.55-6.59 (1H, m), 6.65-6.70 (1H, m), 7.00-7.07 (1H, m), 7.32 (1H, s), 7.37 (1H, m), 7.90 (1H, m), 8.07 (1H, m), 8.17 (1H, t, J=5.2 Hz), 8.27 (1H, d, J=8.3 Hz), 9.18 (1H, s).

Example 96

N1-(1-Ethylpropyl)-5-(2-(((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide

[0307] Tetrahydrofuran (20 ml) and triethylamine (1.73 ml, 12.4 mmol) were added to N1-(1-ethylpropyl)-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (1.45 g, 4.29 mmol); phenyl chloroformate (1.15 ml, 9.1 mmol) was added thereto at 0 °C while stirring; and the reaction mixture was stirred at room temperature for 1 hour. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was concentrated and subjected to silica gel column chromatography to yield a mixture (1.8 g) of phenyl N-

(4-(1-((1-ethylpropyl)amino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate and phenyl N-(4-(1-((1-ethylpropyl)amino)carbonyl-1H-5-indolyl)-oxy-2-pyridyl)-N-(phenoxy-carbonyl)carbamate. A portion of
5 0.6 g of the mixture was dissolved in N,N-dimethylformamide (5 ml); 4-(1-pyrrolidinyl)piperidine (0.7 g, 4.7 mmol) and stirred for 2 hours; the reaction mixture was partitioned between ethyl acetate and water; the organic layer was concentrated; and the
10 residue was purified by silica gel column chromatography (Fuji Silysia NH, ethyl acetate: methanol = 10: 1) to yield as white crystals (202 mg, 0.391 mmol).

MS Spectrum (ESI): 519(M+1).

15 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.90 (6H, t, J=7.5 Hz), 1.20-1.30 (3H, m), 1.47-1.80 (9H, m), 2.03-2.12 (1H, m), 2.40-2.47 (4H, m), 2.77-2.86 (2H, m), 3.62-3.72 (1H, m), 3.88-3.95 (2H, m), 6.53 (1H, dd, J=2.4, 5.9 Hz), 6.66 (1H, d, J=3.5 Hz), 7.02 (1H, dd, J=2.4, 8.8 Hz),
20 7.11 (1H, d, J=2.4 Hz), 7.35 (1H, d, J=2.4 Hz), 7.78 (1H, d, J=8.8 Hz), 7.99 (1H, d, J=3.5 Hz), 8.06 (1H, d, J=5.9 Hz), 8.25 (1H, t, J=8.8 Hz), 9.08 (1H, s).

25 [0308] The starting materials were synthesized by following methods.

Production example 96-1N1-(1-Ethylpropyl)-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide

4-(1H-5-Indolyloxy)-2-pyridinamine (1.85 g, 8.2 mmol, CAS No. 417722-11-3) which was described in WO 02/32872 was dissolved in N,N-dimethylformamide (20 ml); and sodium hydride (394 mg, 9.84 mmol) was added thereto while stirring at room temperature. The reaction mixture was cooled with ice bath after 30 minutes; phenyl N-(1-ethylpropyl)carbamate (1.87 g, 9.03 mmol); and the reaction mixture was stirred at room temperature for 3 hours. The reaction mixture was partitioned between ethyl acetate and water; the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, concentrated; and the residue was purified by silica gel column chromatography (Fuji Silysia NH, hexane: ethyl acetate = 3: 1) to yield the title compound as crystals (1.95 g, 5.8 mmol, 71%).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.89 (6H, t, J=7.5 Hz), 1.44-1.63 (4H, m), 3.60-3.72 (1H, m), 5.73 (1H, d, J=2.6 Hz), 5.80 (2H, s), 6.12 (1H, dd, J=2.6, 6.0 Hz), 6.67 (1H, d, J=4.3 Hz), 7.00 (1H, dd, J=2.6, 8.6 Hz), 7.32 (1H, d, J=2.6 Hz), 7.75 (1H, d, J=6.0 Hz), 7.98 (1H, d, J=4.3 Hz), 8.23 (1H, d, J=8.6 Hz), 9.30 (1H, s).

Production example 96-2Phenyl N-(1-ethylpropyl)carbamate

[0309] 1-Ethylpropylamine (11.6 ml, 100 mmol) was dissolved in tetrahydrofuran (400 ml); pyridine (8.9 ml, 110 mmol) was added thereto at room temperature; and the reaction mixture was stirred. The reaction mixture was cooled with ice bath; phenyl chloroformate (13.8 ml, 110 mmol) was added dropwise; and the reaction mixture was stirred at room temperature for 24 hours. Water was added to the reaction mixture; the reaction mixture was partitioned between ethyl acetate and water; and the organic layer was washed with brine, dried over anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure. The obtained crystals were washed with diethyl ether: hexane = 1: 5 to yield the title compound as crystals (22.3 g, 147 mmol, 59.1%).
¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.87 (6H, t, J=7.5 Hz), 1.30-1.56 (4H, m), 3.20-3.34 (1H, m), 7.03-7.08 (2H, m), 7.14-7.19 (1H, m), 7.32-7.38 (2H, m), 7.51 (1H, d, J=8.7 Hz).

Example 97N1-(1-Ethylpropyl)-5-(2-((4-hydroxypiperidino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0310] N,N-Dimethylformamide (4 ml) and 4-

hydroxypiperidine (360 mg, 3.56 mmol) were added to a mixture (456 mg) of phenyl N-(4-(1-((1-ethylpropyl)amino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-carbamate and phenyl N-(4-(1-((1-ethylpropyl)amino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-N-(phenoxy-carbonyl)carbamate synthesized in Example 96; and the reaction mixture was stirred for 2 hours. The reaction mixture was partitioned between ethyl acetate and water; the organic layer was concentrated; and the residue was purified by silica gel column chromatography (Fuji Silysia NH, ethyl acetate: methanol = 10: 1) to yield the title compound as white crystals (1.37 mg, 0.294 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.90 (6H, t, J=7.5 Hz), 1.18-1.30 (3H, m), 1.45-1.70 (6H, m), 2.92-3.02 (2H, m), 3.55-3.80 (3H, m), 4.63 (1H, d, J=5.1 Hz), 6.53 (1H, m), 6.66 (1H, d, J=3.5 Hz), 7.02 (1H, dd, J=2.5, 8.8 Hz), 7.31 (1H, d, J=2.5 Hz), 7.36 (1H, d, J=2.5 Hz), 7.78 (1H, d, J=8.8 Hz), 7.98 (1H, d, J=3.5 Hz), 8.06 (1H, d, J=5.7 Hz), 8.24 (1H, t, J=8.8 Hz), 9.08 (1H, s).

Example 98

N4-(4-(1-((1-Ethylpropyl)amino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-4-morpholinecarboxamide

[0311] N,N-dimethylformamide (3 ml) and morpholine

(0.22 ml, 2.5 mmol) were added to a mixture (0.324 g) of phenyl N-(4-(1-((1-ethylpropyl)amino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate and phenyl N-(4-(1-((1-ethylpropyl)amino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-N-(phenoxy-carbonyl)carbamate synthesized in Example 96; and the reaction mixture was stirred for 2 hours. The reaction mixture was partitioned between ethyl acetate and water; the organic layer was concentrated; and the residue was purified by silica gel column chromatography (Fuji Silysia NH, ethyl acetate: methanol = 10: 1) to yield the title compound as white crystals (95 mg, 0.21 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.91 (6H, t, J=7.5 Hz), 1.45-1.65 (4H, m), 3.37-3.40 (4H, m), 3.48-3.58 (4H, m), 3.62-3.72 (1H, m), 6.56 (1H, dd, J=2.6, 5.8 Hz), 6.68 (1H, d, J=3.5 Hz), 7.02 (1H, dd, J=2.6, 8.8 Hz), 7.31 (1H, d, J=2.6 Hz), 7.36 (1H, d, J=2.6 Hz), 7.80 (1H, d, J= 9.1 Hz), 8.00 (1H, d, J=3.5 Hz), 8.08 (1H, d, J=5.8 Hz), 8.26 (1H, d, J=8.8 Hz), 9.18 (1H, s).

Example 99

N4-(4-(1-((1-Pentyl)amino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)-4-morpholinecarboxamide

[0312] Similarly to Example 90, the title compound was obtained as colorless crystals (131 mg, 0.29 mmol, 84%) from phenyl N-(4-(1-(1-pentylamino)carbonyl-1H-indol-5-

1
yloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (200 mg,
0.35 mmol) and morpholine (0.15 ml, 1.7 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.88 (3H, t, J=6.0
Hz), 1.31 (4H, m), 1.56 (2H, m), 3.26 (2H, m), 3.35 (4H,
5 m), 3.51 (4H, m), 6.56 (1H, d, J=5.6 Hz), 6.67 (1H, d,
J=3.0 Hz), 7.03 (1H, d, J=8.0 Hz), 7.31 (1H, s), 7.36
(1H, s), 7.91 (1H, d, J=3.0 Hz), 8.08 (1H, d, J=5.6 Hz),
8.20 (1H, t, J=5.6 Hz), 8.26 (1H, d, J=8.0 Hz), 9.18
(1H, s).

10

[0313] The starting materials were synthesized by
following procedures.

Production example 99-1

Phenyl N-(1-pentyl)carbamate

15

Similarly to Example 90-1, the title compound
was obtained as pale yellow crystals (20.5 g, 99 mmol,
99%) from 1-pentylamine (11.6 ml, 100 mmol), pyridine
(8.9 ml, 110 mmol) and phenyl chloroformate (13.8 ml,
110 mmol).

20

¹H-NMR Spectrum (CDCl₃) δ (ppm): 0.92 (3H, t, J=6.8 Hz),
1.36 (4H, m), 1.58 (2H, m), 3.26 (2H, q, J=6.8 Hz),
5.00 (1H, brs), 7.13 (2H, d, J=7.6 Hz), 7.19 (1H, t,
J=7.6 Hz), 7.35 (2H, t, J=7.6 Hz).

25

Production example 99-2

N1-(1-Pentyl)-5-(2-aminopyridin-4-yloxy)-1H-1-

indolecarboxamide

[0314] 4-(1H-5-Indolyloxy)-2-pyridinamine (5.0 g, 22 mmol, CAS No. 417722-11-3) which was described in WO 02/32872 was dissolved in N,N-dimethylformamide (60 ml); sodium hydride (1.06 g, 27 mmol) was added thereto at room temperature; and the reaction mixture was stirred for 30 minutes. Phenyl N-n-pentylcarbamate (5.06 g, 24 mmol) while stirring at room temperature; and the reaction mixture was stirred for 30 minutes. The reaction mixture was partitioned between water and ethyl acetate (insoluble portions were perfectly dissolved by adding a small amount of methanol); and the organic layer was washed with brine, dried over anhydrous sodium sulfate, concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Fuji Silysia NH; hexane: ethyl acetate = 1: 1, ethyl acetate, ethyl acetate: methanol = 95: 5 in this order). The obtained crystals were suspended in hexane: ethanol = 10: 1, filtered off, washed with hexane, and dried under aeration to yield the title compound as colorless crystals (1.55 g, 4.58 mmol, 21%).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.87 (3H, t, J=6.6 Hz), 1.31 (4H m), 1.56 (2H, m), 3.25 (2H, m), 5.74 (1H, d, J=2.8 Hz), 5.83 (2H, s), 6.12 (1H, dd, J=2.8, 5.8 Hz), 6.65 (1H, d, J=3.6 Hz), 7.00 (1H, dd, J=2.0, 8.8

Hz), 7.32 (1H, d, J=2.0 Hz), 7.75 (1H, d, J=5.8 Hz),
7.89 (1H, d, J=3.6 Hz), 8.17 (1H, t, J=5.4 Hz), 8.25
(1H, d, J=8.8 Hz).

5 Production example 99-3

Phenyl N-(4-((1-pentyl)aminocarbonyl-1H-indol-5-yloxy)-
2-pyridyl)-N-(phenoxycarbonyl)carbamate

[0315] Similarly to Example 90-3, the title compound
was obtained as a colorless amorphous solid (2.39 g,
10 4.13 mmol, 90.1%) from N1-(1-pentyl)-5-(2-aminopyridin-
4-yloxy)-1H-1-indolecarboxamide (1.55 g, 4.58 mmol),
triethylamine (1.43 ml, 10.31 mmol), pyridine (0.56 ml,
6.88 mmol), and phenyl chloroformate (1.44 ml, 11.45
mmol).

15 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.87 (3H, t, J=6.4
Hz), 1.31 (4H, m), 1.56 (2H, m), 3.27 (2H, m), 6.56 (1H,
d, J=3.6 Hz), 6.96 (1H, dd, J=2.4, 5.4 Hz), 7.09 (1H,
dd, J=2.4, 9.0 Hz), 7.16 (4H, d, J=7.6 Hz), 7.29 (2H, t,
J=7.6 Hz), 7.43 (5H, m), 7.51 (1H, d, J=2.4 Hz), 7.93
20 (1H, d, J=2.4 Hz), 8.21 (1H, t, J=5.6 Hz), 8.31 (1H, d,
J=9.0 Hz), 8.42 (1H, d, J=5.4 Hz).

Example 100

25 N1-(1-Pentyl)-5-(2-(((4-hydroxypiperidin-1-
yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-
indolecarboxamide

[0316] Similarly to Example 90, the title compound was obtained as colorless crystals (149 mg, 0.320 mmol, 92.6%) from phenyl N-(4-(1-(1-pentyl)aminocarbonyl-1H-indol-5-yloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (200 mg, 0.346 mmol, Production example 99-3) and 4-hydroxypiperidine (174 mg, 1.73 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.87 (3H, m), 1.15-1.40 (6H, m), 1.50-1.70 (4H, m), 2.98 (2H, m), 3.36 (2H, m), 3.59 (1H, m), 3.74 (2H, m), 4.64 (1H, d, J=4.0 Hz), 6.53 (1H, d, J=5.2 Hz), 6.70 (1H, d, J=3.6 Hz), 7.03 (1H, d, J=8.6 Hz), 7.31 (1H, s), 7.35 (1H, s), 7.91 (1H, d, J=3.6 Hz), 8.06 (1H, d, J=5.2 Hz), 8.19 (1H, m), 8.26 (1H, d, J=8.6 Hz), 9.09 (1H, s).

Example 101

N1-(1-Pentyl)-5-(2-((4-(pyrrolidin-1-yl)piperidin-1-ylcarbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide

[0317] Similarly to Example 90, the title compound was obtained as colorless crystals (124 mg, 0.239 mmol, 69.2%) from phenyl N-(4-(1-(1-pentyl)aminocarbonyl-1H-indol-5-yloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (200 mg, 0.346 mmol, Production example 99-3) and 4-(1-pyrrolidinyl)piperidine (267 mg, 1.73 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.87 (3H, t, J=6.8 Hz), 1.20-1.35 (6H, m), 1.52-1.67 (6H, m), 1.74 (2H, m),

2.08 (1H, m), 2.43 (2H, m), 2.81 (2H, t, J=7.6 Hz),
 3.23-3.29 (4H, m), 3.92 (2H, m), 6.53 (1H, dd, J=2.4,
 5.6 Hz), 6.67 (1H, d, J=3.8 Hz), 7.03 (1H, dd, J=2.4,
 9.2 Hz), 7.31 (1H, d, J=2.4 Hz), 7.35 (1H, d, J=2.4 Hz),
 5 7.91 (1H, t, J=3.8 Hz), 8.06 (1H, d, J=5.6 Hz), 8.19
 (1H, d, J=5.4 Hz), 8.26 (1H, d, J=9.2 Hz), 9.09 (1H, s).

Example 102

N1-Methyl-3-chloro-5-(2-(((3-
 10 (diethylamino)propyl)amino)carbonyl)amino-4-
pyridyl)oxy-1H-1-indolecarboxamide

[0318] Phenyl N-(4-(3-chloro-1-(methyamino)carbonyl-
 1H-5-indolyl)oxy-2-pyridyl)carbamate (160 mg), 3-
 (diethylamino)propylamine (120 mg), N,N-
 15 dimethylformamide (5 ml) were mixed together and
 stirred at room temperature for 10 minutes. After the
 addition of aqueous sodium hydrogencarbonate,
 extraction was performed with ethyl acetate. The
 purification by silica gel column chromatography (Fuji
 20 Silysia NH, ethyl acetate and sequentially ethyl
 acetate: methanol = 10: 1) to yield the title compound
 as a white solid (86 mg).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.90 (6H, t, J= 7.2
 Hz), 1.46-1.56 (2H, m), 2.32-2.46 (6H, m), 2.83 (3H, d,
 25 J=4.4 Hz), 3.08-3.15 (2H, m), 6.52 (1H, dd, J=5.6, 2.4
 Hz), 6.84 (1H, d, J=2.4 Hz), 7.16 (1H, dd, J=8.8, 2.4

Hz), 7.28 (1H, d, J=2.4 Hz), 8.02 (1H, d, J= 5.6 Hz), 8.09 (2H, s), 8.21 (1H, q, J=4.4 Hz), 8.33 (1H, d, J=8.8 Hz), 9.04 (1H, s).

5 [0319] The starting materials were synthesized as follows.

Production example 102-1

N1-Methyl-5-(2-amino-4-pyridyl)oxy-3-chloro-1H-indolecarboxamide

10 5-((2-amino-4-pyridyl)oxy)-3-chloro-1H-1-indole
(4.0 g, 15 mmol, CAS No. 417721-98-3) which was described in WO 02/32872 was dissolved in N,N-dimethylformamide (20 ml); sodium hydride (0.68 g, 60% in oil) and phenyl N-methylcarbamate (2.6 g, the
15 product of Production example 2-1) were added thereto; and the reaction mixture was stirred at room temperature for 1 hour. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was concentrated under reduced pressure.
20 The residue was purified by silica gel column chromatography (Fuji Silysia NH, hexane: ethyl acetate = 1: 2) to yield the title compound as a colorless amorphous solid(1.5 g).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.83 (3H, d, J=4.0
25 Hz), 5.78 (1H, d, J=2.0 Hz), 5.88 (2H, brs), 6.14 (1H, dd, J=2.0, 5.8 Hz), 7.14 (1H, dd, J=2.4, 9.0 Hz), 7.23

(1H, d, J=2.4 Hz), 7.78 (1H, d, J=5.8 Hz), 8.08 (1H, s),
8.19 (1H, m), 8.32 (1H, d, J=9.0 Hz).

Production example 102-2

5 Phenyl N-(4-(3-chloro-1-(methylamino)carbonyl-1H-5-
indolyl)oxy-2-pyridyl)carbamate

[0320] While a mixture of N1-methyl-5-(2-amino-4-pyridyl)oxy-3-chloro-1H-1-indolecarboxamide (850 mg, Production example 102-1), triethylamine (0.37 ml),
10 pyridine (320 mg) and N,N-dimethylformamide (10 ml) was cooled with ice and sodium chloride, phenyl chloroformate (630 mg) was added dropwise to the mixture. Aqueous solution of sodium hydrogencarbonate was added thereto after stirring for 20 minutes;
15 extraction was performed with ethyl acetate; and purification was performed by silica gel column chromatography (ethyl acetate). The crystals precipitated by adding ethyl acetate to the residue were filtered off to yield the title compound as white
20 crystals (160 mg).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.80 (3H, d, J=4.4 Hz), 6.70 (1H, dd, J=5.6, 2.4 Hz), 7.10-7.25 (4H, m), 7.26-7.40 (4H, m), 8.07 (1H, s), 8.18 (2H, m), 8.31 (1H, d, J=8.8 Hz), 10.77 (1H, s).

25

Example 103

N1-Methyl-3-chloro-5-(2-((4-(pyrrolidin-1-yl)piperidino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0321] A mixture of N1-methyl-5-(2-amino-4-pyridyl)oxy-3-chloro-1H-1-indolecarboxamide (278 mg, Production example 102-1), triethylamine (0.37 ml), tetrahydrofuran (5 ml) was ice-cooled and stirred; phenyl chloroformate (0.33 ml) was added dropwise to the mixture; and the reaction mixture was further stirred for 10 minutes. Water was added thereto; extraction was performed with ethyl acetate; and purification by silica gel column chromatography was performed to yield a 373 mg of residue. A portion of 245 mg of the residue was dissolved in N,N-dimethylformamide (2 ml); 4-(1-pyrrolidinyl)piperidine (345 mg) was added thereto; and the reaction mixture was stirred at room temperature for 30 minutes. Extraction was performed with ethyl acetate after the addition of water; and the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Fuji Silysia NH) to yield the title compound (154 mg).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.19-1.30 (2H, m), 1.58-1.68 (4H, m), 1.70-1.78 (2H, m), 2.03-2.13 (1H, m),

2.36-2.46 (4H, m), 2.77-2.87 (5H, m), 3.88-3.97 (2H, m),
6.55 (1H, d, J=5.6 Hz), 7.16 (1H, dd. J=9.2, 2.4 Hz),
7.27 (1H, d, J=2.4 Hz), 7.32 (1H, s), 8.08 (1H, d,
J=5.6 Hz), 8.10 (1H, s), 8.19-8.22 (1H, m), 8.33 (1H, d,
5 J=9.2 Hz), 9.13 (1H, brs).

Example 104

N1-Methyl-3-chloro-5-(2-((4-
hydroxypiperidino)carbonyl)amino-4-pyridyl)oxy-1H-1-
10 indolecarboxamide

[0322] A mixture of N1-methyl-5-(2-amino-4-pyridyl)oxy-
3-chloro-1H-1-indolecarboxamide (480 mg, Production
example 102-1), triethylamine (0.63 ml),
tetrahydrofuran (15 ml) was ice-cooled and stirred;
15 phenyl chloroformate (710 mg) was added dropwise to the
mixture; and the reaction mixture was further stirred
for 10 minutes. Extraction was performed with ethyl
acetate after addition of water; and purification was
performed by silica gel column chromatography (hexane:
20 ethyl acetate = 1: 1). The obtained residue was
dissolved in N,N-dimethylformamide (5 ml); 4-
hydroxypiperidine (450 mg) was added thereto; and the
reaction mixture was stirred at room temperature
overnight. Water was added to the reaction mixture;
25 extraction was performed with ethyl acetate; and
purification was performed by silica gel column

chromatography (Fuji Silysia NH, ethyl acetate: methanol = 40: 1) to yield the title compound as colorless powder (78 mg).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.20-1.30 (2H, m),
 5 1.61-1.79 (2H, m), 2.82 (3H, d, J=4.4 Hz), 2.94-3.03
 (2H, m), 3.56-3.63 (1H, m), 3.70-3.78 (2H, m), 4.64 (1H,
 d, J=4.0 Hz), 6.55 (1H, dd, J= 5.6, 2.4 Hz), 7.16 (1H,
 dd, J= 8.8, 2.4 Hz), 7.27 (1H, d, J=2.4 Hz), 7.32 (1H,
 d, J=2.4 Hz), 8.08 (1H, d, J= 5.6 Hz), 8.09 (1H, s),
 10 8.21 (1H, q, J=4.4 Hz), 8.32 (1H, d, J=8.8 Hz), 9.13
 (1H, s).

Example 105

N1-Methyl-3-chloro-5-(2-(((3-(4-
 15 hydroxypiperidino)propyl)amino)carbonyl)amino-4-
pyridyl)oxy-1H-1-indolecarboxamide

[0323] Similarly to Example 103, the title compound was obtained as white crystals from 1-(3-aminopropyl)-4-hydroxypiperidine.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.25-1.38 (2H, m),
 20 1.48-1.58 (2H, m), 1.62-1.70 (2H, m), 1.86-1.97 (2H, m),
 2.18-2.25 (2H, m), 2.60-2.68 (2H, m), 2.83 (3H, d,
 J=3.6 Hz), 3.02-3.13 (2H, m), 3.34-3.42 (1H, m), 4.49
 (1H, d, J=4.0 Hz), 6.52 (1H, dd, J=6.0, 2.4 Hz), 6.84-
 25 6.86 (1H, m), 7.17 (1H, dd, J=8.8, 2.4 Hz), 7.28 (1H, d,
 J=2.4 Hz), 8.01-8.05 (2H, m), 8.10 (1H, s), 8.19-8.24

(1H, m), 8.33 (1H, d, J=8.8 Hz), 9.04 (1H, brs).

[0324] The starting materials were synthesized as follows.

5 Production example 105-1

2-(3-(4-Hydroxypiperidino)propyl)isoindolin-1,3-dione

 N-(3-bromopropyl)phthalimide (26.8 g), 4-hydroxypiperidine (15.0 g) and potassium carbonate (27.6 g) were added to N,N-dimethylformamide; and the
10 reaction mixture was stirred at room temperature overnight. After the addition of water, extraction was performed with ethyl acetate; the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure
15 to yield the title compound (13.9 g).

¹H-NMR Spectrum (CDCl₃) δ (ppm): 1.40-2.05 (6H, m), 2.10-2.60 (4H, m), 2.70-2.90 (2H, m), 3.60-3.85 (3H, m), 7.70-7.75 (2H, m), 7.82-7.87 (2H, m).

20 Production example 105-2

Benzyl N-(3-(4-hydroxypiperidino)propyl)carbamate

 [0325] Ethanol (100 ml) and hydrazine hydrate (1.5 g) were added to 2-(3-(4-hydroxypiperidino)propyl)isoindolin-1,3-dione (4.5 g);
25 the reaction mixture was heated to reflux for 2.5 hours; and the produced crystals were filtered off. N-

Methylmorpholine (2.5 ml) and N-(benzyloxycarbonyloxy)succinimide (5.2 g) were added to the filtrate; and the reaction mixture was stirred at room temperature overnight. Aqueous solution of sodium hydrogencarbonate was added to the reaction mixture; extraction was performed with ethyl acetate; and the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography to yield the title compound (2.96 g).

¹H-NMR Spectrum (CDCl₃) δ (ppm): 1.52-2.10 (6H, m), 2.10-2.60 (4H, m), 2.78-2.90 (2H, m), 3.24-3.33 (2H, m), 3.53-3.86 (1H, m), 5.09 (2H, s), 5.88-5.96 (1H, m), 7.28-7.38 (5H, m).

Production example 105-3

1-(3-Aminopropyl)-4-hydroxypiperidine

[0326] Ethanol (200 ml) and palladium carbon (2.5 g) were added to benzyl N-(3-(4-hydroxypiperidino)propyl)carbamate (2.96 g); and the reaction mixture was stirred vigorously under hydrogen atmosphere overnight. Palladium carbon was removed by filtration, and the filtrate was concentrated to yield the title compound (1.5 g).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.25-1.38 (2H, m),

1.41-1.49 (2H, m), 1.61-1.69 (2H, m), 1.84-1.95 (2H, m),
2.18-2.25 (2H, m), 2.49-2.57 (2H, m), 2.58-2.69 (2H, m),
3.30-3.42 (1H, m).

5 Example 106

N1-Methyl-3-chloro-5-(2-((4-(2-hydroxyethyl)piperazin-
1-yl)carbonyl)amino-4-pyridyl)oxy-1H-1-
indolecarboxamide

10 [0327] Similarly to Example 104, the title compound was
obtained from 4-(2-hydroxyethyl)piperazine.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.30-2.48 (6H, m),
2.82 (3H, d, J=4.4 Hz), 3.30-3.40 (4H, m), 3.46 (2H, q,
J=5.6 Hz), 4.38 (1H, t, J=5.6 Hz), 6.57 (1H, dd, J=5.6,
2.4 Hz), 7.16 (1H, dd, J=8.8, 2.4 Hz), 7.29 (1H, d,
15 J=2.4 Hz), 7.32 (1H, d, J=2.4 Hz), 8.07-8.13 (2H, m),
8.21 (1H, q, J=4.4 Hz), 8.32 (1H, d, J=8.8 Hz), 9.15
(1H, s).

Example 107

20 N4-(4-(3-Chloro-1-(methylamino)carbonyl-1H-5-
indolyl)oxy-2-pyridyl)-4-morpholinecarboxamide

[0328] Similarly to Example 104, the title compound was
obtained from morpholine.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.82 (3H, d, J=4.4
25 Hz), 3.33-3.40 (4H, m), 3.49-3.56 (4H, m), 6.58 (1H, dd,
J=5.6, 2.4 Hz), 7.16 (1H, dd, J=8.8, 2.4 Hz), 7.27 (1H,

d, J= 2.4 Hz), 7.32 (1H, d, J=2.4 Hz), 8.06-8.13 (2H, m), 8.21 (1H, q, J=4.4 Hz), 8.32 (1H, d, J=8.8 Hz), 9.22 (1H, s).

5 Example 108

N1-Methyl-3-chloro-5-(2-((4-ethylpiperazin-1-yl)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide
[0329] Similarly to Example 103, the title compound was obtained from N-ethylpiperazine.

10 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.96 (3H, t, J=7.2 Hz), 2.24-2.32 (6H, m), 2.82 (3H, d, J=4.0 Hz), 3.34-3.39 (4H, m), 6.57 (1H, dd, J=6.0, 2.4 Hz), 7.17 (1H, dd, J=9.2, 2.4 Hz), 7.27 (1H, d, J=2.4 Hz), 7.32 (1H, d, J=2.4 Hz), 8.07-8.10 (2H, m), 8.18-8.25 (1H, m), 8.33
15 (1H, d, J=9.2 Hz), 9.17 (1H, brs).

Example 109

N1-Ethyl-3-chloro-5-(2-((4-hydroxypiperidino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

20 [0330] Similarly to Example 104, the title compound was obtained as a colorless amorphous solid from N1-ethyl-5-(2-amino-4-pyridyl)oxy-3-chloro-1H-1-indolecarboxamide.

25 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.15 (2H, m), 1.61 (3H, t, J=7.2 Hz), 1.60-1.70 (2H, m), 2.94-3.02 (2H, m),

3.26-3.36 (2H, m), 3.56-3.63 (1H, m), 3.70-3.78 (2H, m),
4.64 (1H, d, J= 4.4 Hz), 6.55 (1H, dd, J=5.6, 2.4 Hz),
7.16 (1H, dd, J= 8.8, 2.4 Hz), 7.27 (1H, d, J=2.4 Hz),
7.32 (1H, d, J=2.4 Hz), 8.08 (1H, d, J= 5.6 Hz), 8.13
5 (1H, s), 8.22-8.27 (1H, m), 8.32 (1H, d, J=8.8 Hz),
9.12 (1H, s).

[0331]The starting material was synthesized as follows.

Production example 109-1

10 N1-Ethyl-5-(2-amino-4-pyridyl)oxy-3-chloro-1H-1-
indolecarboxamide

Phenyl N-ethylcarbamate was added to a solution
of 5-(2-amino-4-pyridyl)oxy-3-chloro-1H-1-indole (1.35
g, CAS No. 417721-98-3) which was described in WO
15 02/32872, sodium hydride (210 mg) and N,N-
dimethylformamide; and the reaction mixture was stirred
for 1 hour. An aqueous solution of ammonium chloride
was added to the reaction mixture; extraction was
performed with ethyl acetate; and purification by
20 silica gel column chromatography (Fuji Silysia NH,
hexane: ethyl acetate = 1: 2) to yield the title
compound as a colorless oil (1.07 g).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.15 (3H, t, J=7.2
Hz), 3.25-3.35 (2H, m), 5.76 (1H, d, J=2.4 Hz), 5.87
25 (2H, s), 6.14 (1H, dd, J=5.6, 2.4 Hz), 7.13 (1H, dd,
J=8.8, 2.4 Hz), 7.23 (1H, d, J=2.4 Hz), 7.77 (1H, d,

J=5.6 Hz), 8.11 (1H, s), 8.20-8.25 (1H, m), 8.31 (1H, d, J=8.8 Hz).

Example 110

5 N1-Ethyl-3-chloro-5-(2-(((3-(4-hydroxypiperidino)propyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0332] Similarly to Example 103, the title compound was obtained as white crystals from N1-ethyl-5-(2-amino-4-pyridyl)oxy-3-chloro-1H-1-indolecarboxamide and 1-(3-aminopropyl)-4-hydroxypiperidine.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.16 (3H, t, J=7.2 Hz), 1.26-1.38 (2H, m), 1.48-1.57 (2H, m), 1.63-1.70 (2H, m), 1.86-1.97 (2H, m), 2.18-2.25 (2H, m), 2.60-2.68 (2H, m), 3.05-3.13 (2H, m), 3.26-3.34 (2H, m), 3.34-3.42 (1H, m), 4.49 (1H, d, J=4.0 Hz), 6.52 (1H, dd, J=6.0, 2.4 Hz), 6.84-6.86 (1H, m), 7.16 (1H, dd, J=8.8, 2.4 Hz), 7.28 (1H, d, J=2.4 Hz), 7.98-8.05 (2H, m), 8.14 (1H, s), 8.22-8.28 (1H, m), 8.33 (1H, d, J=8.8 Hz), 9.03 (1H, brs).

Example 111

25 N1-Ethyl-3-chloro-5-(2-(((3-(diethylamino)propyl)amino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0333] Similarly to Example 104, the title compound was

obtained from N1-ethyl-5-(2-amino-4-pyridyl)oxy-3-chloro-1H-1-indolecarboxamide and 3-(diethylamino)propylamine.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.90 (6H, t, J=7.2 Hz), 1.16 (3H, t, J=7.2 Hz), 1.46-1.54 (2H, m), 2.33-2.44 (6H, m), 3.07-3.14 (2H, m), 3.26-3.34 (2H, m), 6.52 (1H, dd, J=5.6, 2.4 Hz), 6.83 (1H, s), 7.16 (1H, dd, J=8.8, 2.4 Hz), 7.28 (1H, d, J=2.4 Hz), 8.02 (1H, d, J=5.6 Hz), 8.04-8.13 (1H, brs), 8.14 (1H, s), 8.23-8.27 (1H, m), 8.33 (1H, d, J=8.8 Hz), 9.04 (1H, s).

Example 112

N1,3-Dimethyl-5-(2-((4-hydroxypiperidino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0334] Similarly to Example 104, the title compound was obtained as a colorless amorphous solid from N1,3-dimethyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.17-1.30 (2H, m), 1.61-1.70 (2H, m), 2.19 (3H, s), 2.80 (3H, d, J=4.0 Hz), 2.94-3.03 (2H, m), 3.56-3.64 (1H, m), 3.70-3.78 (2H, m), 4.64 (1H, d, J=4.0 Hz), 6.52 (1H, dd, J=5.6, 2.4 Hz), 7.02 (1H, dd, J=8.8, 2.4 Hz), 7.29-7.33 (2H, m), 7.66 (1H, s), 8.00 (1H, q, J=4.0 Hz), 8.05 (1H, d, J=5.6 Hz), 8.25 (1H, d, J=8.8 Hz), 9.08 (1H, s).

[0335] The starting materials were synthesized as follows.

Production example 112-1

5 4-(3-Methyl-1H-5-indolyl)oxy-2-pyridinamine

 A mixture of 5-hydroxy-3-methylindole (4.7 g),
2-amino-4-chloropyridine (4.1 g), sodium hydride (1.3
g), and dimethyl sulfoxide (40 ml) was stirred at 160
°C for 15 hours. Water was added thereto; extraction
10 was performed with ethyl acetate; and purification was
performed by silica gel column chromatography (ethyl
acetate, sequentially, ethyl acetate: methanol= 20: 1).
The solvent was distilled off to yield the title
compound as a brown solid (1.6 g).

15 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.29 (3H, s), 5.70
(1H, d, J=2.4 Hz), 5.77 (2H, s), 6.10 (1H, dd, J=5.6,
2.4 Hz), 6.80 (1H, dd, J=8.8, 2.4 Hz), 7.15 (1H, s),
7.17 (1H, d, J=2.4 Hz), 7.35 (1H, d, J=8.8 Hz), 7.72
(1H, d, J=5.6 Hz), 10.83 (1H, s).

20

Production example 112-2

N1,3-Dimethyl-5-(2-amino-4-pyridyl)oxy-1H-1-
indolecarboxamide

 [0336] Phenyl N-methylcarbamate (350 mg, Production
25 example 2-1) was added to a solution of 4-(3-methyl-1H-
5-indolyl)oxy-2-pyridinamine (500 mg), sodium hydride

(93 mg) and N,N-dimethylformamide (5 ml) at room temperature; and the reaction mixture was stirred for 2 hours and 45 minutes. Water was added to the reaction mixture; extraction was performed with ethyl acetate; and purification was performed by NH-silica gel column chromatography (Fuji Silysia, hexane: ethyl acetate = 1: 2, sequentially, ethyl acetate) to yield the title compound as a pale yellow amorphous solid (365 mg).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.19 (3H, s), 2.80 (3H, d, J=4.0 Hz), 5.73 (1H, d, J=2.4 Hz), 5.83 (2H, s), 6.12 (1H, dd, J=5.6, 2.4 Hz), 7.00 (1H, dd, J=8.8, 2.4 Hz), 7.27 (1H, d, J= 2.4 Hz), 7.64 (1H, s), 7.75 (1H, d, J=5.6 Hz), 7.98 (1H, q, J=4.0 Hz), 8.24 (1H, d, J=8.8 Hz).

Example 113

N1,3-Dimethyl-5-(2-((4-(pyrrolidin-1-yl)piperidino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0337] Similarly to Example 104, the title compound was obtained as a colorless amorphous solid from N1,3-dimethyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide and 4-(1-pyrrolidinyl)piperidine.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.17-1.79 (2H, m), 1.60-1.67 (4H, m), 1.70-1.79 (2H, m), 2.03-2.13 (1H, m), 2.19 (3H, s), 2.40-2.57 (4H, m), 2.77-2.86 (5H, m),

3.88-3.96 (2H, m), 6.52 (1H, dd, J= 5.6, 2.4 Hz), 7.02 (1H, dd, J=8.8, 2.4 Hz), 7.28-7.85 (2H, m), 7.66 (1H, s), 8.00 (1H, q, J=4.0 Hz), 8.05 (1H, d, J=5.6 Hz), 8.25 (1H, d, J=8.8 Hz), 9.08 (1H, s).

5

Example 114

N1-Cyclopropyl-5-(2-((4-hydroxypiperidino)carbonyl)amino-4-pyridyl)oxy-3-methyl-1H-1-indolecarboxamide

10 [0338] Similarly to Example 104, the title compound was obtained as a colorless amorphous solid from N1-cyclopropyl-5-(2-amino-4-pyridyl)oxy-3-methyl-1H-1-indolecarboxamide.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.56-0.60 (2H, m),
15 2.67-2.73 (2H, m), 1.19-1.29 (2H, m), 1.61-1.70 (2H, m), 2.18 (3H, s), 2.72-2.78 (1H, m), 2.94-3.03 (2H, m), 3.56-3.63 (1H, m), 3.70-3.77 (2H, m), 4.64 (1H, d, J=4.0 Hz), 6.51 (1H, dd, J=5.6, 2.4 Hz), 7.02 (1H, dd, J=8.8, 2.4 Hz), 7.28-7.32 (2H, m), 7.65 (1H, s), 8.05
20 (1H, d, J=5.6 Hz), 8.11 (1H, d, J=2.4 Hz), 8.24 (1H, d, J=8.8 Hz), 9.08 (1H, s).

[0339]The starting material was synthesized as follows.

Production example 114-1

25 N1-Cyclopropyl-5-(2-amino-4-pyridyl)oxy-3-methyl-1H-1-indolecarboxamide

Similarly to Production example 112-2, the title compound was obtained as a colorless amorphous solid from phenyl N-cyclopropylcarbamate.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.55-0.60 (2H, m),
5 0.68-0.73 (2H, m), 2.18 (3H, s), 2.70-2.79 (1H, m),
5.73 (1H, d, J=2.4 Hz), 5.83 (2H, s), 6.12 (1H, dd,
J=5.6, 2.4 Hz), 7.00 (1H, dd, J=8.8, 2.4 Hz), 7.26 (1H,
d, J=2.4 Hz), 7.63 (1H, s), 7.75 (1H, d, J=5.6 Hz),
8.09 (1H, d, J=2.4 Hz), 8.23 (1H, d, J=8.8 Hz).

10

Example 115

N1-Cyclopropyl-5-(2-((4-(2-hydroxyethyl)piperazin-1-yl)carbonyl)amino-4-pyridyl)oxy-3-methyl-1H-1-indolecarboxamide

15 [0340] Similarly to Example 104, the title compound was obtained as a colorless amorphous solid from N1-cyclopropyl-5-(2-amino-4-pyridyl)oxy-3-methyl-1H-1-indolecarboxamide and 1-(2-hydroxyethyl)piperazine.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.57-0.60 (2H, m),
20 0.67-0.74 (2H, m), 2.18 (3H, s), 2.30-2.37 (6H, m),
2.70-2.78 (1H, m), 3.30-3.38 (4H, m), 3.46 (2H, q, J=6.4 Hz),
4.38 (1H, t, J=6.4 Hz), 6.53 (1H, dd, J=5.6, 2.4 Hz),
7.02 (1H, dd, J=8.8, 2.4 Hz), 7.28-7.32 (2H, m),
7.65 (1H, s), 8.06 (1H, d, J=5.6 Hz), 8.11 (1H, d,
25 J=2.4 Hz), 8.24 (1H, d, J=8.8 Hz), 9.10 (1H, s).

Example 116N1-Methyl-5-(2-((methylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0341] Similarly to Example 5, the title compound was
5 obtained as white crystals (19.5 mg, 0.058 mmol, 58%)
from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (52 mg, 0.1 mmol) synthesized in Production example 5-2 and 40% methylamine in methanol.

10 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.64 (3H, d, J=4.4 Hz), 2.83 (3H, d, J=4.4 Hz), 6.50 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 6.82 (1H, d, J=2.4 Hz), 7.04 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, d, J=2.4 Hz), 7.87 (1H, d, J=3.6 Hz), 7.95 (1H, m), 8.02 (1H, d, J=5.6 Hz),
15 8.16 (1H, m), 8.28 (1H, d, J=8.8 Hz), 9.07 (1H, s).

Example 117N1-Methyl-5-(2-((ethylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

20 [0342] Similarly to Example 5, the title compound was
obtained as white crystals (15.0 mg, 0.042 mmol, 42%)
from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (52
25 mg, 0.1 mmol) synthesized in Production example 5-2 and 2.0 M ethylamine in tetrahydrofuran.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.02 (3H, t, J=7.2 Hz), 2.83 (3H, d, J=4.0 Hz), 3.10 (2H, m), 6.50 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 6.86 (1H, d, J=2.4 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, d, J=2.4 Hz), 7.87 (1H, d, J=3.6 Hz), 7.96 (1H, m), 8.02 (1H, d, J=5.6 Hz), 8.17 (1H, m), 8.28 (1H, d, J=8.8 Hz), 8.99 (1H, s).

Example 118

10 N1-Methyl-5-(2-((cyclopropylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0343] Similarly to Example 5, the title compound was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxy carbonyl) carbamate (52 mg, 0.1 mmol) synthesized in Production example 5-2 and cyclopropylamine.

15 ¹H-NMR Spectrum (CDCl₃) δ (ppm): 0.58-0.62 (2H, m), 0.71-0.79 (2H, m), 2.70 (1H, m), 3.07 (3H, d, J=4.8 Hz), 5.64 (1H, m), 6.26 (1H, m), 6.41 (1H, m), 6.58 (1H, d, J=3.6 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.20-7.30 (2H, m), 7.42-7.53 (2H, m), 7.90 (1H, d, J=5.6 Hz), 8.19 (1H, d, J=8.8 Hz).

25 Example 119

N1-Methyl-5-(2-((diethylamino)carbonyl)amino-4-

pyridyl)oxy-1H-1-indolecarboxamide

[0344] Similarly to Example 5, the title compound was obtained as white crystals (24.7 mg, 0.065 mmol, 65%) from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (52 mg, 0.1 mmol) synthesized in Production example 5-2 and diethylamine.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.01 (6H, t, J=7.2 Hz), 2.82 (3H, d, J=4.4 Hz), 3.20-3.50 (4H, m), 6.54 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 7.04 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, d, J=2.4 Hz), 7.40 (1H, d, J=2.4 Hz), 7.78 (1H, d, J=3.6 Hz), 8.06 (1H, d, J=5.6 Hz), 8.16 (1H, m), 8.28 (1H, d, J=8.8 Hz), 8.59 (1H, s).

Example 120N1-Methyl-5-(2-((1-propylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0345] Similarly to Example 5, the title compound (28.0 mg, 0.076 mmol, 76%) was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (52 mg, 0.1 mmol) synthesized in Production example 5-2 and 1-propylamine.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.83 (3H, t, J=7.2 Hz), 1.40 (2H, m), 2.83 (3H, d, J=4.4 Hz), 3.04 (2H, m),

6.49 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz),
6.86 (1H, s), 7.04 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, d,
J=2.4 Hz), 7.87 (1H, d, J=3.6 Hz), 8.01-8.03 (2H, m),
8.16 (1H, m), 8.28 (1H, d, J=8.8 Hz), 9.00 (1H, s).

5 [0346]

Example 121

N1-Methyl-5-(2-((2-propylamino)carbonyl)amino-4-
pyridyl)oxy-1H-1-indolecarboxamide

Similarly to Example 5, the title compound (20.7
10 mg, 0.056 mmol, 56%) was obtained as white crystals
from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-
indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (52
mg, 0.1 mmol) synthesized in Production example 5-2 and
2-propylamine.

15 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.06 (6H, d, J=6.8
Hz), 2.83 (3H, d, J=4.4 Hz), 3.74 (1H, m), 6.49 (1H, dd,
J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 6.89 (1H, d,
J=2.4 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, d,
J=2.4 Hz), 7.81 (1H, m), 7.87 (1H, d, J=3.6 Hz), 8.02
20 (1H, d, J=5.6 Hz), 8.16 (1H, m), 8.28 (1H, d, J=8.8 Hz),
8.90 (1H, s).

Example 122

N1-Methyl-5-(2-((cyclopentylamino)carbonyl)amino-4-
25 pyridyl)oxy-1H-1-indolecarboxamide

[0347] Similarly to Example 5, the title compound (30.7

mg, 0.078 mmol, 78%) was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (52 mg, 0.1 mmol) synthesized in Production example 5-2 and cyclopentylamine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.10-1.90 (8H, m), 2.83 (3H, d, J=4.4 Hz), 3.89 (1H, m), 6.50 (1H, d, J=2.4, 5.6 Hz), 6.68 (1H, d, J=3.6 Hz), 6.90 (1H, d, J=2.4 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, d, J=2.4 Hz), 7.87 (1H, m), 7.93 (1H, d, J=3.6 Hz), 8.00 (1H, d, J=5.6 Hz), 8.15 (1H, m), 8.28 (1H, d, J=8.8 Hz), 8.89 (1H, s).

Example 123

N1-Methyl-5-(2-((cyclohexylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0348] Similarly to Example 5, the title compound (32.5 mg, 0.080 mmol, 80%) was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (52 mg, 0.1 mmol) synthesized in Production example 5-2 and cyclohexylamine.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 1.00-2.00 (8H, m), 2.83 (3H, d, J=4.4 Hz), 3.40-3.60 (2H, m), 3.75 (1H, m), 6.11 (1H, s), 6.43 (1H, m), 6.60 (1H, d, J=3.6 Hz), 6.95 (1H, m), 7.04 (1H, m), 7.20-7.30 (2H, m), 7.44 (1H,

d, J=3.6 Hz), 7.95 (1H, d, J=5.6 Hz), 8.20 (1H, d, J=8.8 Hz), 9.20 (1H, m).

Example 124

5 N1-Methyl-5-(2-((2-propenylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0349] Similarly to Example 5, the title compound (18.5 mg, 0.051 mmol, 51%) was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxy-carbonyl)carbamate (52 mg, 0.1 mmol) synthesized in Production example 5-2 and allylamine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.85 (3H, d, J=4.8 Hz), 3.75 (2H, m), 5.06 (1H, dd, J=1.6, 10.4 Hz), 5.14 (1H, dd, J=1.6, 17.2 Hz), 5.87 (1H, m), 6.54 (1H, dd, J=2.4, 5.6 Hz), 6.69 (1H, d, J=3.6 Hz), 6.87 (1H, d, J=2.4 Hz), 7.06 (1H, dd, J=2.4, 8.8 Hz), 7.39 (1H, d, J=2.4 Hz), 7.89 (1H, d, J=3.6 Hz), 8.05 (1H, d, J=5.6 Hz), 8.16-8.19 (2H, m), 8.30 (1H, d, J=8.8 Hz), 9.13 (1H, s).

Example 125

25 N1-Methyl-5-(2-((2-propynylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0350] Similarly to Example 5, the title compound (16.8 mg, 0.046 mmol, 46%) was obtained as white crystals

from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (52 mg, 0.1 mmol) synthesized in Production example 5-2 and propargylamine.

5 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.85 (3H, d, J=4.4 Hz), 3.12 (1H, m), 3.92 (2H, m), 6.56 (1H, dd, J=2.4, 5.6 Hz), 6.70 (1H, d, J=3.6 Hz), 6.87 (1H, d, J=2.4 Hz), 7.06 (1H, dd, J=2.4, 8.8 Hz), 7.39 (1H, d, J=2.4 Hz), 7.89 (1H, d, J=3.6 Hz), 8.07 (1H, d, J=5.6 Hz), 8.18
10 (1H, m), 8.29-8.31 (2H, m), 9.21 (1H, s).

Example 126

N1-Methyl-5-(2-((benzylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

15 [0351] Similarly to Example 5, the title compound (26.1 mg, 0.063 mmol, 63%) was obtained as white crystals from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (52 mg, 0.1 mmol) synthesized in Production example 5-2 and
20 benzylamine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.85 (3H, d, J=4.4 Hz), 4.34 (2H, d, J=5.6 Hz), 6.53 (1H, dd, J=2.4, 5.6 Hz), 6.69 (1H, d, J=3.6 Hz), 6.90 (1H, d, J=2.4 Hz), 7.06 (1H, dd, J=2.4, 8.8 Hz), 7.20-7.35 (5H, m), 7.39
25 (1H, d, J=2.4 Hz), 7.89 (1H, d, J=3.6 Hz), 8.04 (1H, d, J=5.6 Hz), 8.17 (1H, m), 8.30 (1H, d, J=8.8 Hz), 8.51

(1H, m), 9.17 (1H, s).

Example 127

N1-Methyl-5-(2-((furfurylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0352] Similarly to Example 5, the title compound (9.1 mg, 0.022 mmol, 22%) was obtained as white powder from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (52 mg, 0.1 mmol) synthesized in Production example 5-2 and furfurylamine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.85 (3H, d, J=4.4 Hz), 4.32 (2H, d, J=5.2 Hz), 6.24 (1H, s), 6.38 (1H, s), 6.54 (1H, m), 6.69 (1H, d, J=3.6 Hz), 6.90 (1H, s), 7.06 (1H, m), 7.39 (1H, s), 7.57 (1H, s), 7.89 (1H, d, J=2.4 Hz), 8.05 (1H, d, J=5.6 Hz), 8.18 (1H, m), 8.31 (1H, d, J=8.8 Hz), 8.38 (1H, m), 9.15 (1H, s).

Example 128

N1-Methyl-5-(2-((thiophen-2-ylmethylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0353] Similarly to Example 5, the title compound (22.6 mg, 0.054 mmol, 54%) was obtained as white powder from phenyl N-(4-(1-(methylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (52 mg, 0.1

mmol) synthesized in Production example 5-2 and 2-thiophenemethylamine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.85 (3H, d, J=4.4 Hz), 4.50 (2H, d, J=5.6 Hz), 6.54 (1H, dd, J=2.4, 5.6 Hz), 6.69 (1H, d, J=3.6 Hz), 6.88-6.98 (3H, m), 7.06 (1H, dd, J=2.4, 8.8 Hz), 7.35-7.39 (2H, m), 7.89 (1H, d, J=3.6 Hz), 8.04 (1H, d, J=5.6 Hz), 8.18 (1H, m), 8.30 (1H, d, J=8.8 Hz), 8.55 (1H, m), 9.18 (1H, s).

10 Example 129

N1-Ethyl-5-(2-((ethylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0354] Similarly to Example 27, the title compound (23.1 mg, 0.063 mmol, 63%) was obtained as white crystals from phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate (42 mg, 0.1 mmol) synthesized in Production example 27-2 and 2.0 M ethylamine in tetrahydrofuran.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.04 (3H, t, J=7.2 Hz), 1.18 (3H, t, J=7.2 Hz), 3.12 (2H, m), 3.31 (2H, m), 6.52 (1H, dd, J=2.4, 5.6 Hz), 6.69 (1H, d, J=3.6 Hz), 6.88 (1H, d, J=2.4 Hz), 7.05 (1H, dd, J=2.4, 8.8 Hz), 7.38 (1H, d, J=2.4 Hz), 7.92 (1H, d, J=3.6 Hz), 7.97 (1H, m), 8.04 (1H, d, J=5.6 Hz), 8.23 (1H, m), 8.30 (1H, d, J=8.8 Hz), 9.01 (1H, s).

Example 130N1-Ethyl-5-(2-((diethylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0355] Similarly to Example 27, the title compound
5 (25.8 mg, 0.065 mmol, 65%) was obtained as white
crystals from phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-
indolyl)oxy-2-pyridyl)carbamate (42 mg, 0.1 mmol)
synthesized in Production example 27-2 and diethylamine.
¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.03 (6H, t, J=7.2
10 Hz), 1.19 (3H, t, J=7.2 Hz), 3.20-3.40 (6H, m), 6.55
(1H, dd, J=2.4, 5.6 Hz), 6.69 (1H, d, J=3.6 Hz), 7.05
(1H, dd, J=2.4, 8.8 Hz), 7.38 (1H, s), 7.43 (1H, d,
J=2.4 Hz), 7.92 (1H, d, J=3.6 Hz), 8.08 (1H, d, J=5.6
Hz), 8.23 (1H, m), 8.30 (1H, d, J=8.8 Hz), 8.62 (1H, s).

15

Example 131N1-Dimethyl-5-(2-((methylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0356] Similarly to Example 28, the title compound
20 (17.5 mg, 0.05 mmol, 35%) was obtained as white powder
from N1-dimethyl-5-(2-amino-4-pyridyl)oxy-1H-1-
indolecarboxamide (42 mg, 0.14 mmol) and 40%
methylamine in methanol.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.67 (3H, d, J=4.4
25 Hz), 3.05 (6H, s), 6.53 (1H, dd, J=2.4, 5.6 Hz), 6.67
(1H, d, J=3.6 Hz), 6.83 (1H, d, J=2.4 Hz), 7.04 (1H, dd,

J=2.4, 8.8 Hz), 7.40 (1H, d, J=2.4 Hz), 7.68-7.71 (2H, m), 8.00-8.05 (2H, m), 9.10 (1H, s).

5 [0357] The starting materials were synthesized by the following methods.

Production example 131-1

Phenyl N,N-dimethylcarbamate

10 Similarly to Production example 2-1, the title compound (6.27 g, 0.038 mol, 38%) was obtained as a colorless oil from 2.0 M diethylamine in tetrahydrofuran (50 ml, 0.1 mol), phenyl chloroformate (13.8 ml, 0.11 mol) and pyridine (8.9 ml, 0.11 mol).

¹H-NMR Spectrum (CDCl₃) δ(ppm): 3.02 (3H, s), 3.11 (3H, s), 7.09-7.39 (5H, m).

15

Production example 131-2

N1-Dimethyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide

20 [0358] Similarly to Production example 1-3, the title compound (128 mg, 0.43 mmol, 43%) was obtained as white crystals from 4-(1H-5-indolyloxy)-2-pyridinamine (225 mg, 1.0 mmol) and phenyl N,N-dimethylcarbamate.

25 ¹H-NMR Spectrum (CDCl₃) δ(ppm): 3.13 (6H, s), 4.36 (2H, brs), 5.92 (1H, d, J=2.4 Hz), 6.31 (1H, dd, J=2.4, 5.6 Hz), 6.59 (1H, d, J=3.6 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.30 (1H, d, J=2.4 Hz), 7.37 (1H, d, J=3.6 Hz),

7.70 (1H, d, J=8.8 Hz), 7.91 (1H, d, J=5.6 Hz).

Example 132

N1-Dimethyl-5-(2-((ethylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

5 [0359] Similarly to Example 28, the title compound (21.4 mg, 0.058 mmol, 41%) was obtained as white powder from N1-dimethyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (42 mg, 0.14 mmol) and 2.0 M ethylamine in tetrahydrofuran.

10 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.05 (3H, t, J=7.2 Hz), 3.05 (6H, s), 3.13 (2H, m), 6.52 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 6.87 (1H, d, J=2.4 Hz), 7.04 (1H, dd, J=2.4, 8.8 Hz), 7.40 (1H, d, J=2.4 Hz),
15 7.68-7.71 (2H, m), 8.00-8.05 (2H, m), 9.02 (1H, s).

Example 133

N1-Dimethyl-5-(2-((dimethylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

20 [0360] Similarly to Example 28, the title compound (15.1 mg, 0.041 mmol, 29%) was obtained as white powder from N1-dimethyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (42 mg, 0.14 mmol) and 2.0 M dimethylamine in tetrahydrofuran.

25 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.85 (6H, s), 3.03 (6H, s), 6.54 (1H, d, J=5.6 Hz), 6.65 (1H, d, J=3.6 Hz),

7.02 (1H, d, J=8.8 Hz), 7.30-7.50 (2H, m), 7.60-7.69 (2H, m), 8.06 (1H, d, J=5.6 Hz), 8.81 (1H, s).

Example 134

5 N1-Benzyl-5-(2-((methylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0361] Similarly to Example 28, the title compound (12.5 mg, 0.030 mmol, 24%) was obtained as white powder from N1-benzyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (45 mg, 0.13 mmol) and 40% methylamine in methanol.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.66 (3H, d, J=4.4 Hz), 4.51 (2H, d, J=5.6 Hz), 6.52 (1H, dd, J=2.4, 5.6 Hz), 6.72 (1H, d, J=3.6 Hz), 6.84 (1H, d, J=2.4 Hz), 7.06 (1H, dd, J=2.4, 8.8 Hz), 7.20-7.44 (6H, m), 7.96 (1H, m), 8.00-8.05 (2H, m), 8.31 (1H, d, J=8.8 Hz), 8.83 (1H, t, J=5.6 Hz), 9.09 (1H, s).

[0362] The starting material was synthesized by the following methods.

Production example 134-1

N1-Benzyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide

Similarly to Production example 1-3, the title compound (45 mg, 0.13 mmol, 13%) was obtained as white powder from 4-(1H-5-indolyloxy)-2-pyridinamine (225 mg,

1.0 mmol) and phenyl N-benzylcarbamate.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 4.38 (2H, brs), 4.68 (2H, d, J=4.0 Hz), 5.82 (1H, m), 5.92 (1H, m), 6.30 (1H, m), 6.61 (1H, d, J=3.6 Hz), 7.07 (1H, dd, J=2.4, 8.8 Hz), 7.26-7.47 (7H, m), 7.91 (1H, d, J=5.6 Hz), 8.19 (1H, d, J=8.8 Hz).

Example 135

5-(2-((Methylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indole-1-carboxylic acid pyrrolidin-1-ylamide

[0363] Similarly to Example 28, the title compound (21.1 mg, 0.056 mmol, 27%) was obtained as white powder from 5-(2-amino-4-pyridyl)oxy-1H-1-indole-1-carboxylic acid pyrrolidin-1-ylamide (67 mg, 0.21 mmol) and 40% methylamine in methanol.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.80-2.00 (4H, m), 2.67 (3H, d, J=4.4 Hz), 3.40-3.60 (4H, m), 6.53 (1H, dd, J=2.4, 5.6 Hz), 6.66 (1H, d, J=3.6 Hz), 6.85 (1H, d, J=2.4 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.39 (1H, d, J=2.4 Hz), 7.80 (1H, d, J=3.6 Hz), 7.83 (1H, d, J=8.8 Hz), 8.00 (1H, m), 8.04 (1H, d, J=5.6 Hz), 9.10 (1H, s).

[0364] The starting materials were synthesized by the following methods.

Production example 135-1

Phenyl pyrrolidin-1-ylcarboxylate

Similarly to Production example 2-1, the title compound (2.68 g, 0.014 mol, 14%) was obtained as white crystals from pyrrolidine (8.3 ml, 0.1 mol), phenyl chloroformate (13.8 ml, 0.11 mol) and pyridine (8.9 ml, 0.11 mol).

¹H-NMR Spectrum (CDCl₃) δ(ppm): 1.90-1.99 (4H, m), 3.46-3.59 (4H, m), 7.20-7.37 (5H, m).

Production example 135-2

5-(2-Amino-4-pyridyl)oxy-1H-1-indole-1-carboxylic acid pyrrolidin-1-ylamide

[0365] Similarly to Production example 1-3, the title compound (146 mg, 0.45 mmol, 60%) was obtained as white powder from 4-(1H-5-indolyloxy)-2-pyridinamine (170 mg, 0.76 mmol) and phenyl pyrrolidin-1-ylcarboxylate.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 1.96-2.02 (4H, m), 3.60-3.67 (4H, m), 4.35 (2H, brs), 5.91 (1H, d, J=2.4 Hz), 6.31 (1H, dd, J=2.4, 5.6 Hz), 6.57 (1H, d, J=3.6 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.29 (1H, d, J=2.4 Hz), 7.43 (1H, d, J=3.6 Hz), 7.88 (1H, d, J=8.8 Hz), 7.91 (1H, d, J=5.6 Hz).

Example 136

5-(2-((Pyrrolidin-1-ylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indole-1-carboxylic acid pyrrolidin-1-ylamide

[0366] Similarly to Example 28, the title compound (6.2 mg, 0.015 mmol, 9.2%) was obtained as white powder from 5-(2-amino-4-pyridyl)oxy-1H-1-indole-1-carboxylic acid pyrrolidin-1-ylamide (52 mg, 0.16 mmol) and pyrrolidine.

5 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.70-1.90 (8H, m), 3.20-3.40 (4H, m), 3.50-3.70 (4H, m), 6.56 (1H, dd, J=2.4, 5.6 Hz), 6.66 (1H, d, J=3.6 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.38 (1H, d, J=2.4 Hz), 7.45 (1H, d, J=2.4 Hz), 7.80 (1H, d, J=3.6 Hz), 7.84 (1H, d, J=8.8
10 Hz), 8.08 (1H, d, J=5.6 Hz), 8.61 (1H, s).

Example 137

N1-(2-Propynyl)-5-(2-((ethylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

15 [0367] Similarly to Example 28, the title compound (16.5 mg, 0.044 mmol, 25%) was obtained as white crystals from N1-(2-propynyl)-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (54 mg, 0.18 mmol) and 2.0 M ethylamine in tetrahydrofuran.

20 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.04 (3H, t, J=7.2 Hz), 3.12 (2H, m), 3.23 (1H, m), 4.10 (2H, m), 6.52 (1H, dd, J=2.4, 5.6 Hz), 6.53 (1H, d, J=3.6 Hz), 6.88 (1H, d, J=2.4 Hz), 7.08 (1H, dd, J=2.4, 8.8 Hz), 7.40 (1H, d, J=2.4 Hz), 7.92 (1H, d, J=3.6 Hz), 8.00 (1H, m), 8.04
25 (1H, d, J=5.6 Hz), 8.31 (1H, d, J=8.8 Hz), 8.73 (1H, m), 9.02 (1H, s).

[0368] The starting materials were synthesized by the following methods.

Production example 137-1

5 Phenyl N-(2-propynyl)carbamate

Similarly to Production example 2-1, the title compound (7.64 g, 0.044 mol, 87%) was obtained as white crystals from 2-propynylamine (3.43 ml, 0.05 mol), phenyl chloroformate (6.9 ml, 0.055 mol) and pyridine
10 (4.45 ml, 0.055 mol).

¹H-NMR Spectrum (CDCl₃) δ(ppm): 2.30 (1H, t, J=2.8 Hz), 4.05-4.15 (2H, m), 5.22 (1H, brs), 7.10-7.40 (5H, m).

Production example 137-2

15 N1-(2-Propynyl)-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0369] Similarly to Production example 1-3, the title compound (169 mg, 0.55 mmol, 28%) was obtained as white crystals from 4-(1H-5-indolyloxy)-2-pyridinamine (450
20 mg, 2.0 mmol) and phenyl N-(2-propynyl)carbamate.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 2.35 (1H, m), 4.20-4.40 (4H, m), 5.72 (1H, brs), 5.92 (1H, d, J=2.4 Hz), 6.30 (1H, dd, J=2.4, 5.6 Hz), 6.63 (1H, d, J=3.6 Hz), 7.08 (1H, dd, J=2.4, 8.8 Hz), 7.30 (1H, d, J=2.4 Hz), 7.46
25 (1H, d, J=3.6 Hz), 7.92 (1H, d, J=5.6 Hz), 8.20 (1H, d, J=8.8 Hz).

Example 138N1-(2-Propynyl)-5-(2-((diethylamino)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

5 [0370] Similarly to Example 28, the title compound (27.9 mg, 0.069 mmol, 39%) was obtained as white crystals from N1-(2-propynyl)-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (54 mg, 0.18 mmol) and N,N-diethylamine.

10 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.03 (6H, t, J=7.2 Hz), 3.23 (1H, m), 3.25-3.40 (4H, m), 4.01 (2H, m), 6.56 (1H, dd, J=2.4, 5.6 Hz), 6.72 (1H, d, J=3.6 Hz), 7.08 (1H, dd, J=2.4, 8.8 Hz), 7.39 (1H, d, J=2.4 Hz), 7.43 (1H, d, J=2.4 Hz), 7.92 (1H, d, J=3.6 Hz), 8.08
15 (1H, d, J=5.6 Hz), 8.31 (1H, d, J=8.8 Hz), 8.63 (1H, s), 8.73 (1H, m).

Example 139N1-(2-Propynyl)-5-(2-((pyrrolidin-1-yl)carbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

20 [0371] Similarly to Example 28, the title compound (25.1 mg, 0.062 mmol, 35%) was obtained as white crystals from N1-(2-propynyl)-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (54 mg, 0.18 mmol) and
25 pyrrolidine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.70-1.90 (4H, m),

3.22 (1H, m), 3.25-3.40 (4H, m), 4.10 (2H, m), 6.56 (1H, dd, J=2.4, 5.6 Hz), 6.71 (1H, d, J=3.6 Hz), 7.07 (1H, dd, J=2.4, 8.8 Hz), 7.39 (1H, d, J=2.4 Hz), 7.44 (1H, d, J=2.4 Hz), 7.92 (1H, d, J=5.6 Hz), 8.08 (1H, d, J=5.6 Hz), 8.30 (1H, d, J=8.8 Hz), 8.62 (1H, s), 8.72 (1H, m).

Example 140

N1-Methyl-5-(6-((morpholin-4-yl)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0372] Similarly to Example 28, the title compound (12.5 mg, 0.032 mmol, 12%) was obtained as pale yellow powder from N1-methyl-5-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (73 mg, 0.26 mmol) and morpholine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.83 (3H, d, J=3.6 Hz), 3.43-3.45 (4H, m), 3.55-3.58 (4H, m), 6.63 (1H, d, J=3.6 Hz), 7.26 (1H, s), 7.32 (1H, d, J=8.8 Hz), 7.77 (1H, d, J=3.6 Hz), 7.90 (1H, s), 8.05 (1H, m), 8.14 (1H, d, J=8.8 Hz), 8.29 (1H, s), 9.21 (1H, s), 9.34 (1H, s).

[0373] The starting materials were synthesized by the following methods.

Production example 140-1

6-Chloro-4-(1H-5-indolylamino)pyrimidine

4,6-Dichloropyrimidine (5.89 g, 40 mmol), 5-

aminoindole (6.27 g, 47 mmol) and N,N-diisopropylethylamine (20.6 ml, 0.12 mol) were dissolved in N-methylpyrrolidone (80 ml), and the reaction mixture was stirred at 50°C for 2.5 hours. The reaction mixture was partitioned between ethyl acetate and water; the aqueous layer was subjected to re-extraction with ethyl acetate; and the combined organic layer was washed with brine, and dried over anhydrous sodium sulfate. The solvent was distilled off under reduced pressure; a small amount of ethyl acetate was added to the residue to crystallize; and the crystals were filtered off, washed with diethyl ether, and dried under aeration to yield the title compound (3.70 g, 15 mmol, 38%) as white crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 6.42 (1H, m), 6.62 (1H, brs), 7.11 (1H, d, J=8.0 Hz), 7.35-7.40 (2H, m), 7.72 (1H, brs), 8.38 (1H, s), 9.68 (1H, s), 11.11 (1H, s).

Production example 140-2

6-Amino-4-(1H-5-indolylamino)pyrimidine

[0374] A 7N ammonia in methanol (60 ml) was added to 6-chloro-4-(1H-5-indolylamino)pyrimidine (2.455 g, 10 mmol); and the reaction mixture was heated in a sealed tube at 130°C for 90 hours. The solvent was distilled off under reduced pressure; the residue was purified by

silica gel column chromatography (eluent; ethyl acetate: tetrahydrofuran = 1: 1); diethyl ether was added to crystallize; and the crystals were filtered off, washed with diethyl ether, and dried under aeration to yield the title compound (1.563 g, 6.9 mmol, 69%) as pale brown crystals.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 4.50 (2H, brs), 5.66 (1H, m), 6.55 (1H, m), 6.68 (1H, brs), 7.07 (1H, dd, J=2.4, 8.8 Hz), 7.25-7.28 (1H, m), 7.40 (1H, d, J=8.8 Hz), 7.52 (1H, d, J=2.4 Hz), 8.19 (1H, s), 8.29 (1H, brs).

Production example 140-3

N1-Methyl-5-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0375] Similarly to Production example 2-3, the title compound (295 mg, 1.05 mmol, 52%) was obtained as white crystals from 6-amino-4-(1H-5-indolylamino)pyrimidine (450.5 mg, 2.0 mmol) and phenyl N-methylcarbamate.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 3.09 (3H, d, J=4.0 Hz), 4.56 (2H, brs), 5.52 (1H, m), 5.73 (1H, m), 6.61 (1H, d, J=3.6 Hz), 6.66 (1H, brs), 7.19 (1H, dd, J=2.4, 8.8 Hz), 7.43 (1H, d, J=3.6 Hz), 7.48 (1H, d, J=2.4 Hz), 8.13 (1H, d, J=2.4 Hz), 8.21 (1H, s).

Example 141

N1-Methyl-5-(6-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0376] Similarly to Example 28, the title compound
 5 (20.6 mg, 0.045 mmol, 17%) was obtained as white crystals from N1-methyl-5-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (73 mg, 0.26 mmol) and 4-(pyrrolidin-1-yl)piperidine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.20-1.36 (2H, m),
 10 1.60-1.70 (4H, m), 1.70-1.85 (2H, m), 2.40-2.60 (5H, m), 2.83 (3H, d, J=4.4 Hz), 2.85-2.95 (2H, m), 3.95-4.05 (2H, m), 6.63 (1H, d, J=3.6 Hz), 7.24 (1H, s), 7.31 (1H, dd, J=2.4, 8.8 Hz), 7.77 (1H, d, J=3.6 Hz), 7.90 (1H, s), 8.05 (1H, m), 8.14 (1H, d, J=8.8 Hz), 8.28 (1H, s),
 15 9.14 (1H, s), 9.31 (1H, s).

Example 142

N1-Ethyl-5-(6-((ethylamino)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0377] Sodium hydride (69 mg, 1.73 mmol) was suspended
 20 in N,N-dimethylformamide (3 ml); 6-amino-4-(1H-5-indolylamino)pyrimidine (311 mg, 1.38 mmol) was added thereto at room temperature under nitrogen stream; the reaction mixture was stirred for 30minutes; phenyl N-ethylcarbamate (286 mg, 1.73 mmol) was added thereto;
 25 and the reaction mixture was stirred overnight. The

reaction mixture was partitioned between a solvent mixture of ethyl acetate-tetrahydrofuran (1: 1) and a saturated aqueous solution of sodium hydrogencarbonate; the organic layer was washed with water and brine, and dried over anhydrous sodium sulfate; the solvent was distilled off; the residue was purified by silica gel column chromatography (eluent; ethyl acetate: tetrahydrofuran=1: 1); eluted fractions were concentrated; ethyl acetate was added to the residue to crystallize; and the crystals were filtered off, and dried under aeration to yield the title compound (14.3 mg, 0.039 mmol, 2.8%) as white crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.07 (3H, t, J=7.2 Hz), 1.18 (3H, t, J=7.2 Hz), 3.11-3.40 (4H, m), 6.64 (1H, d, J=3.6 Hz), 6.87 (1H, s), 7.29 (1H, dd, J=2.4, 8.8 Hz), 7.62 (1H, m), 7.80 (1H, d, J=3.6 Hz), 7.86 (1H, s), 8.09-8.17 (2H, m), 8.27 (1H, s), 9.06 (1H, s), 9.35 (1H, s).

Furthermore, the eluted fractions obtained in the above chromatography were concentrated; the residue was purified again by silica gel column chromatography (eluent; ethyl acetate: methanol = 95: 5); eluted fractions were concentrated; ethyl acetate was added to the residue to crystallize; and the crystals were filtered off, and dried under aeration to yield N1-ethyl-5-(6-aminopyrimidin-4-yl)amino-1H-1-

indolecarboxamide (210 mg, 0.71 mmol, 51%) as white crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.18 (3H, t, J=7.2 Hz), 3.20-3.40 (2H, m), 5.72 (1H, m), 6.24 (2H, brs),
5 6.61 (1H, d, J=3.6 Hz), 7.21 (1H, dd, J=2.4, 8.8 Hz),
7.76 (1H, s), 7.79 (1H, d, J=3.6 Hz), 8.01 (1H, s),
8.07-8.14 (2H, m), 8.74 (1H, s).

Example 143

10 N1-Ethyl-5-(6-((diethylamino)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0378] Similarly to Example 28, the title compound (24.5 mg, 0.062 mmol, 26%) was obtained as white crystals from N1-ethyl-5-(6-aminopyrimidin-4-yl)amino-
15 1H-1-indolecarboxamide(70 mg, 0.24 mmol) and diethylamine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.07 (6H, t, J=7.2 Hz), 1.18 (3H, t, J=7.2 Hz), 3.20-3.50 (6H, m), 6.63 (1H, d, J=3.6 Hz), 7.31-7.33 (2H, m), 7.80 (1H, d, J=3.6 Hz),
20 7.91 (1H, s), 8.09-8.15 (2H, m), 8.28 (1H, s), 8.66 (1H, s), 9.33 (1H, s).

Example 144

25 N1-Ethyl-5-(6-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0379] Similarly to Example 28, the title compound (43.3 mg, 0.091 mmol, 39%) was obtained as white crystals from N1-ethyl-5-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (70 mg, 0.24 mmol) and 4-(pyrrolidin-1-yl)piperidine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.18 (3H, t, J=7.2 Hz), 1.20-1.36 (2H, m), 1.60-1.70 (4H, m), 1.70-1.85 (2H, m), 2.40-2.60 (5H, m), 2.85-2.95 (2H, m), 3.20-3.50 (2H, m), 3.95-4.05 (2H, m), 6.63 (1H, d, J=3.6 Hz), 7.24 (1H, s), 7.31 (1H, d, J=8.0 Hz), 7.80 (1H, d, J=3.6 Hz), 7.90 (1H, s), 8.10-8.15 (2H, m), 8.28 (1H, s), 9.14 (1H, s), 9.31 (1H, s).

Example 145

N1-Ethyl-5-(6-((2-(N,N-diethylamino)ethylamino)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0380] Similarly to Example 28, the title compound (43.0 mg, 0.098 mmol, 42%) was obtained as white crystals from N1-ethyl-5-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (70 mg, 0.24 mmol) and 2-(N,N-diethylamino)ethylamine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.96 (6H, t, J=7.2 Hz), 1.18 (3H, t, J=7.2 Hz), 2.30-2.60 (6H, m), 3.10-3.40 (4H, m), 6.64 (1H, d, J=3.6 Hz), 6.87 (1H, s), 7.29 (1H, d, J=8.8 Hz), 7.71 (1H, m), 7.80 (1H, d,

J=3.6 Hz), 7.88 (1H, s), 8.09-8.20 (2H, m), 8.25 (1H, s), 9.21 (1H, s), 9.34 (1H, s).

Example 146

5 N1-Phenyl-5-(6-((diethylamino)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0381] Similarly to Example 28, the title compound (27.5 mg, 0.062 mmol, 27%) was obtained as white crystals from N1-phenyl-5-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (80 mg, 0.23 mmol) and diethylamine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.08 (6H, t, J=7.2 Hz), 3.20-3.40 (4H, m), 6.74 (1H, d, J=3.6 Hz), 7.13 (1H, dd, J=2.4, 8.8 Hz), 7.33-7.42 (4H, m), 7.64-7.67 (2H, m), 7.98-8.03 (2H, m), 8.13 (1H, d, J=8.8 Hz), 8.31 (1H, s), 8.69 (1H, s), 9.39 (1H, s), 10.00 (1H, s).

[0382] The starting material was synthesized by the following method.

20 Production example 146-1

N1-Phenyl-5-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

Similarly to Production example 2-3, the title compound (160 mg, 0.46 mmol, 35%) was obtained as pale brown powder from 6-amino-4-(1H-5-indolylamino)pyrimidine (300 mg, 1.33 mmol) and phenyl

isocyanate.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 4.61 (2H, brs), 5.76 (1H, m), 6.68 (1H, d, J=3.6 Hz), 6.77 (1H, s), 7.22-7.25 (2H, m), 7.35-7.45 (3H, m), 7.50-7.60 (4H, m), 8.16 (1H, d, J=8.8 Hz), 8.22 (1H, s).

Example 147

N1-Phenyl-5-(6-((3-(N,N-diethylamino)propylamino)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0383] Similarly to Example 28, the title compound (56.2 mg, 0.11 mmol, 48%) was obtained as white powder from N1-phenyl-5-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (80 mg, 0.23 mmol) and 3-(N,N-diethylamino)propylamine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.80-1.00 (6H, m), 1.40-1.65 (2H, m), 2.20-2.60 (6H, m), 3.00-3.40 (2H, m), 6.70-6.88 (2H, m), 7.10-7.17 (1H, m), 7.30-7.49 (3H, m), 7.60-7.80 (3H, m), 7.90-8.40 (4H, m), 9.10-9.40 (2H, m), 10.00-10.14 (1H, m).

Example 148

N1-Cyclopropyl-5-(6-((4-(piperidin-1-yl)piperidin-1-yl)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0384] Similarly to Production example 2-3, a crude

product of N1-cyclopropyl-5-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (132 mg) was obtained as white powder from 6-amino-4-(1H-5-indolylamino)pyrimidine (300 mg, 1.33 mmol) and phenyl N-cyclopropylcarbamate.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.60-0.63 (2H, m), 0.70-0.74 (2H, m), 2.76 (1H, m), 5.73 (1H, s), 6.24 (2H, brs), 6.59 (1H, d, J=3.6 Hz), 7.02 (1H, dd, J=2.4, 8.8 Hz), 7.74-7.76 (2H, m), 8.01 (1H, s), 8.12 (1H, d, J=8.8 Hz), 8.15 (1H, d, J=2.4 Hz), 8.75 (1H, s).

[0385] Similarly to Example 28, the title compound (20.6 mg, 0.041 mmol, 3.1% in 2 processes) was obtained as white crystals from the above crude product.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.59-0.63 (2H, m), 0.70-0.76 (2H, m), 1.20-1.60 (8H, m), 1.60-1.80 (2H, m), 2.30-2.80 (8H, m), 4.05-4.20 (2H, m), 6.61 (1H, d, J=3.6 Hz), 7.24 (1H, s), 7.32 (1H, dd, J=2.4, 8.8 Hz), 7.76 (1H, d, J=3.6 Hz), 7.90 (1H, s), 8.13 (1H, d, J=8.8 Hz), 8.17 (1H, d, J=2.4 Hz), 8.28 (1H, s), 9.15 (1H, s), 9.32 (1H, s).

Example 149

N1-Dimethyl-5-(6-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0386] Similarly to Example 28, the title compound (19.2 mg, 0.040 mmol, 21%) was obtained as white powder from N1-dimethyl-5-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (56 mg, 0.19 mmol) and 4-(pyrrolidin-1-yl)piperidine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.20-1.36 (2H, m), 1.60-1.70 (4H, m), 1.70-1.85 (2H, m), 2.40-2.60 (5H, m), 2.85-2.95 (2H, m), 3.00 (6H, s), 3.95-4.05 (2H, m), 6.60 (1H, d, J=3.2 Hz), 7.22 (1H, d, J=1.2 Hz), 7.30 (1H, dd, J=2.0, 8.8 Hz), 7.50-7.55 (2H, m), 7.88 (1H, brs), 8.26 (1H, d, J=1.2 Hz), 9.13 (1H, s), 9.29 (1H, s).

[0387] The starting material was synthesized by the following method.

Production example 149-1

N1-Dimethyl-5-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

Similarly to Production example 2-3, the title compound (101 mg, 0.34 mmol, 34%) was obtained as white crystals from 6-amino-4-(1H-5-indolylamino)pyrimidine (225.3 mg, 1.0 mmol) and phenyl N,N-dimethylcarbamate.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 3.02 (6H, s), 5.71 (1H, s), 6.23 (2H, brs), 6.60 (1H, d, J=3.6 Hz), 7.22 (1H, dd, J=2.0, 8.8 Hz), 7.50-7.55 (2H, m), 7.74 (1H, d, J=2.0 Hz), 8.00 (1H, s), 8.73 (1H, s).

Example 150

N1-Dimethyl-5-(6-((3-diethylaminopropyl)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

5 [0388] Similarly to Example 28, the title compound (55.3 mg, 0.12 mmol, 66%) was obtained as white crystals from N1-dimethyl-5-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (55 mg, 0.19 mmol) and 10 3-diethylaminopropylamine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.92 (6H, t, J=7.2 Hz), 1.50-1.55 (2H, m), 2.30-2.45 (6H, m), 3.00 (6H, s), 3.10-3.15 (2H, m), 6.60 (1H, dd, J=0.8, 3.6 Hz), 6.82 (1H, brs), 7.28 (1H, dd, J=2.0, 8.8 Hz), 7.50-7.55 (2H, 15 m), 7.71 (1H, m), 7.84 (1H, brs), 8.23 (1H, d, J=0.8 Hz), 9.08 (1H, s), 9.32 (1H, s).

Example 151

5-(6-((4-(Pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino-4-pyrimidyl)amino-1H-1-indole-1-carboxylic acid pyrrolidin-1-ylamide

20 [0389] Similarly to Example 28, the title compound (13.1 mg, 0.026 mmol, 14%) was obtained as white crystals from 5-(6-amino-4-pyrimidyl)amino-1H-1-indole-1-carboxylic acid pyrrolidin-1-ylamide (61 mg, 0.19 25 mmol) and 4-(pyrrolidin-1-yl)piperidine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.20-1.36 (2H, m),
1.60-1.70 (4H, m), 1.70-1.90 (6H, m), 2.40-2.60 (5H, m),
2.85-2.95 (2H, m), 3.40-3.60 (4H, m), 3.95-4.05 (2H, m),
6.59 (1H, d, J=3.2 Hz), 7.22 (1H, brs), 7.28 (1H, dd,
5 J=2.0, 8.8 Hz), 7.60-7.70 (2H, m), 7.87 (1H, m), 8.26
(1H, d, J=1.2 Hz), 9.12 (1H, s), 9.28 (1H, s).

[0390] The starting material was synthesized by the
following method.

10 Production example 151-1

5-(6-Amino-4-pyrimidyl)amino-1H-1-indole-1-carboxylic
acid pyrrolidin-1-ylamide

Similarly to Production example 2-3, the title
compound (122 mg, 0.38 mmol, 38%) was obtained as white
15 crystals from 6-amino-4-(1H-5-indolylamino)pyrimidine
(225.3 mg, 1.0 mmol) and phenyl pyrrolidin-1-
ylcarboxylate.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.80-1.95 (4H, m),
3.50-3.60 (4H, m), 5.71 (1H, s), 6.23 (2H, brs), 6.59
20 (1H, d, J=3.6 Hz), 7.20 (1H, dd, J=2.0, 8.8 Hz), 7.64-
7.69 (2H, m), 7.74 (1H, d, J=2.0 Hz), 8.00 (1H, s),
8.73 (1H, s).

Example 152

25 5-(6-((Morpholin-4-yl)carbonyl)amino-4-pyrimidyl)amino-
1H-1-indole-1-carboxylic acid pyrrolidin-1-ylamide

[0391] Similarly to Example 28, the title compound (30.3 mg, 0.070 mmol, 37%) was obtained as white powder from 5-(6-amino-4-pyrimidyl)amino-1H-1-indole-1-carboxylic acid pyrrolidin-1-ylamide (61 mg, 0.19 mmol) and morpholine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.80-1.90 (4H, m), 3.40-3.50 (4H, m), 3.50-3.60 (8H, m), 6.59 (1H, d, J=2.8 Hz), 7.24 (1H, d, J=2.0 Hz), 7.28 (1H, dd, J=2.0, 8.8 Hz), 7.63-7.69 (2H, m), 7.88 (1H, brs), 8.27 (1H, d, J=2.8 Hz), 9.19 (1H, s), 9.31 (1H, s).

Example 153

N1-(2-Propyl)-5-(6-((2-propylamino)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0392] Sodium hydride (48 mg, 1.2 mmol) was suspended in N,N-dimethylformamide (2.5 ml); 6-amino-4-(1H-5-indolylamino)pyrimidine (225.3 mg, 1.0 mmol) was added thereto at room temperature under nitrogen stream; the reaction mixture was stirred for 30minutes; phenyl N-(2-propyl)carbamate (215 mg, 1.2 mmol) was added thereto; and the reaction mixture was stirred overnight. The reaction mixture was partitioned between a solvent mixture of ethyl acetate-tetrahydrofuran (1: 1) and a saturated aqueous solution of sodium hydrogencarbonate; and the organic layer was washed with water and brine,

and dried over anhydrous sodium sulfate. After the solvent was distilled off, the residue was purified by silica gel column chromatography (eluent; ethyl acetate); eluted fractions were concentrated; ethyl acetate was added to the residue to crystallize; and the crystals were filtered off, and dried under aeration to yield the title compound (31.3 mg, 0.079 mmol, 7.9%) as white crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.09 (6H, d, J=6.8 Hz), 1.20 (6H, d, J=6.8 Hz), 3.75 (1H, m), 4.00 (1H, m), 6.60 (1H, d, J=3.6 Hz), 6.89 (1H, s), 7.27 (1H, dd, J=2.0, 8.8 Hz), 7.45 (1H, m), 7.80-7.90 (3H, m), 8.11 (1H, d, J=8.8 Hz), 8.24 (1H, s), 8.91 (1H, s), 9.31 (1H, s).

The above chromatography was further performed by eluting ethyl acetate: methanol = 95: 5; the eluted fractions were concentrated; ethyl acetate-hexane (1: 5) was added to the residue to crystallize; and the crystals were filtered off, and dried under aeration to yield N1-(2-propyl)-5-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (77.8 mg, 0.25 mmol, 25%) as white crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.29 (6H, d, J=6.8 Hz), 3.98 (1H, m), 5.70 (1H, s), 6.21 (2H, brs), 6.57 (1H, d, J=2.8 Hz), 7.18 (1H, d, J=8.8 Hz), 7.72 (1H, s), 7.79-7.82 (2H, m), 7.98 (1H, s), 8.08 (1H, d, J=8.8 Hz),

8.72 (1H, s).

Example 154

N1-(2-Propyl)-5-(6-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0393] Similarly to Example 28, the title compound (36.3 mg, 0.074 mmol, 60%) was obtained as white powder from N1-(2-propyl)-5-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (38 mg, 0.12 mmol) and 4-(pyrrolidin-1-yl)piperidine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.20 (6H, d, J=6.8 Hz), 1.20-1.36 (2H, m), 1.60-1.70 (4H, m), 1.70-1.85 (2H, m), 2.40-2.60 (5H, m), 2.85-2.95 (2H, m), 3.90-4.10 (3H, m), 6.60 (1H, d, J=3.6 Hz), 7.22 (1H, s), 7.29 (1H, d, J=8.0 Hz), 7.80-8.00 (3H, m), 8.10 (1H, d, J=8.0 Hz), 8.26 (1H, s), 9.13 (1H, s), 9.29 (1H, s).

Example 155

N1-(2-Propyl)-5-(6-((3-diethylaminopropyl)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0394] Similarly to Example 28, the title compound (27.3 mg, 0.059 mmol, 48%) was obtained as white crystals from N1-(2-propyl)-5-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (38 mg, 0.12 mmol) and

3-diethylaminopropylamine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.91 (6H, t, J=6.8 Hz), 1.20 (6H, d, J=6.8 Hz), 1.40-1.60 (2H, m), 2.20-2.50 (6H, m), 3.10-3.20 (2H, m), 4.00 (1H, m), 6.60 (1H, d, J=3.6 Hz), 6.82 (1H, s), 7.26 (1H, d, J=8.8 Hz), 7.70 (1H, m), 7.80-7.85 (3H, m), 8.11 (1H, d, J=8.8 Hz), 8.24 (1H, s), 9.08 (1H, s), 9.32 (1H, s).

Example 156

N1-Methyl-4-(6-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0395] Similarly to Example 28, the title compound (21.0 mg, 0.045 mmol, 31%) was obtained as white powder from N1-methyl-4-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (41 mg, 0.15 mmol) and 4-(pyrrolidin-1-yl)piperidine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.20-1.40 (2H, m), 1.60-1.70 (4H, m), 1.70-1.85 (2H, m), 2.40-2.60 (5H, m), 2.81 (3H, d, J=4.4 Hz), 2.85-2.95 (2H, m), 3.90-4.10 (2H, m), 6.85 (1H, m), 7.18 (1H, t, J=8.0 Hz), 7.35 (1H, d, J=4.0 Hz), 7.55 (1H, d, J=8.0 Hz), 7.64 (1H, d, J=8.0 Hz), 7.70 (1H, d, J=4.0 Hz), 7.93 (1H, d, J=8.0 Hz), 8.08 (1H, m), 8.26 (1H, s), 9.19 (1H, s).

[0396] The starting materials were synthesized by the

following methods.

Production example 156-1

6-Chloro-4-(1H-4-indolylamino)pyrimidine

4,6-Dichloropyrimidine (1.01 g, 6.6 mmol), 4-
5 aminoindole (900 mg, 6.6 mmol) and N,N-
diisopropylethylamine (3.14 ml, 18 mmol) were dissolved
in N,N-dimethylformamide (20 ml), and the reaction
mixture was stirred at 80°C for 6 hours. The reaction
mixture was partitioned between ethyl acetate and
10 water; the aqueous layer was subjected to re-extraction
with ethyl acetate, and the combined organic layer was
washed with brine, and dried over anhydrous sodium
sulfate. The solvent was distilled off under reduced
pressure; and a small amount of methanol was added to
15 the residue to crystallize; and the crystals were
filtered off, washed with methanol and ethyl acetate,
and dried under aeration to yield the title compound
(599 mg, 2.5 mmol, 37%) as pale brown crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 6.49 (1H, brs), 6.75
20 (1H, brs), 7.10 (1H, m), 7.25 (1H, d, J=8.0 Hz), 7.33-
7.40 (2H, m), 8.42 (1H, s), 9.71 (1H, brs), 11.24 (1H,
brs).

Production example 156-2

25 6-Amino-4-(1H-4-indolylamino)pyrimidine

[0397] A 7N methanol solution of ammonia (50 ml) and

tetrahydrofuran (20 ml) were added to 6-chloro-4-(1H-4-indolylamino)pyrimidine (599 mg, 2.5 mmol) and the reaction mixture was heated in a sealed tube at 130 °C for 137 hours. The solvent was distilled off under reduced pressure; the residue was purified by silica gel column chromatography (eluent; ethyl acetate: tetrahydrofuran = 1: 1); diethyl ether was added to the residue to crystallize; the crystals were filtered off, washed with diethyl ether, and dried under aeration to yield the title compound (454 mg, 2.0 mmol, 82%) as pale brown crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 5.74 (1H, s), 6.20 (2H, brs), 6.50 (1H, m), 7.00 (1H, t, J=8.0 Hz), 7.09 (1H, d, J=8.0 Hz), 7.15-7.30 (2H, m), 7.98 (1H, s), 8.55 (1H, s), 11.06 (1H, brs).

Production example 156-3

N1-Methyl-4-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0398] Similarly to Production example 2-3, the title compound (124.7 mg, 0.44 mmol, 44%) was obtained as pale brown crystals from 6-amino-4-(1H-5-indolylamino)pyrimidine (225.3 mg, 1.0 mmol) and phenyl N-methylcarbamate.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.81 (3H, d, J=4.0 Hz), 5.75 (1H, s), 6.27 (2H, brs), 6.76 (1H, d, J=4.0

Hz), 7.17 (1H, t, J=8.0 Hz), 7.43 (1H, d, J=8.0 Hz),
7.70 (1H, d, J=4.0 Hz), 7.92 (1H, d, J=8.0 Hz), 8.00
(1H, s), 8.06 (1H, m), 8.70 (1H, s).

5 Example 157

N1-Methyl-4-(6-((4-(piperidin-1-yl)piperidin-1-yl)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

10 [0399] Similarly to Example 28, the title compound (7.7 mg, 0.016 mmol, 11%) was obtained as white powder from N1-methyl-4-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (41 mg, 0.15 mmol) and 4-(piperidin-1-yl)piperidine.

15 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.20-1.60 (8H, m), 1.60-1.80 (2H, m), 2.30-2.80 (7H, m), 2.81 (3H, d, J=4.4 Hz), 4.05-4.20 (2H, m), 6.85 (1H, m), 7.18 (1H, t, J=8.0 Hz), 7.35 (1H, d, J=4.0 Hz), 7.55 (1H, d, J=8.0 Hz), 7.65-7.70 (2H, m), 7.92 (1H, d, J=8.0 Hz), 8.06 (1H, m), 8.26 (1H, s), 9.18 (1H, s).

20

Example 158

N1-Methyl-4-(6-((3-diethylaminopropylamino)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

25 [0400] Similarly to Example 28, the title compound (23.3 mg, 0.053 mmol, 37%) was obtained as white

crystals from N1-methyl-4-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (41 mg, 0.15 mmol) and 3-diethylaminopropylamine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.92 (6H, t, J=6.8 Hz), 1.40-1.60 (2H, m), 2.20-2.50 (6H, m), 2.82 (3H, d, J=4.4 Hz), 3.10-3.20 (2H, m), 6.80 (1H, m), 6.93 (1H, d, J=6.8 Hz), 7.19 (1H, t, J= 8.0 Hz), 7.55 (1H, d, J=8.0 Hz), 7.60-7.70 (2H, m), 7.95 (1H, d, J=8.0 Hz), 8.08 (1H, m), 8.23 (1H, s), 9.11 (1H, s), 9.26 (1H, s).

Example 159

N1-(4-Fluorophenyl)-4-(6-((3-diethylaminopropylamino)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0401] Similarly to Example 28, the title compound (28.6 mg, 0.055 mmol, 40%) was obtained as white powder from N1-(4-fluorophenyl)-4-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (50 mg, 0.14 mmol) and 3-diethylaminopropylamine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.93 (6H, t, J=6.8 Hz), 1.40-1.60 (2H, m), 2.30-2.50 (6H, m), 3.10-3.15 (2H, m), 6.90 (1H, d, J=3.6 Hz), 6.97 (1H, m), 7.18-7.26 (3H, m), 7.60-7.70 (4H, m), 7.90-8.00 (2H, m), 8.25 (1H, s), 9.13 (1H, s), 9.32 (1H, s), 10.09 (1H, brs).

[0402] The starting material was synthesized by the following method.

Production example 159-1

N1-(4-Fluorophenyl)-4-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

Similarly to Production example 2-3, the title compound (109 mg, 0.30 mmol, 30%) was obtained as pale yellow powder from 6-amino-4-(1H-4-indolylamino)pyrimidine (225.3 mg, 1.0 mmol) and phenyl N-(4-fluorophenyl)carbamate.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 5.78 (1H, s), 6.29 (2H, brs), 6.86 (1H, d, J=3.6 Hz), 7.15-7.30 (3H, m), 7.51 (1H, d, J=8.0 Hz), 7.60-7.70 (2H, m), 7.89 (1H, d, J=8.0 Hz), 7.92 (1H, d, J=3.6 Hz), 8.01 (1H, s), 8.76 (1H, s), 10.07 (1H, s).

Example 160

N1-(4-Fluorophenyl)-4-(6-((2-diethylaminoethylamino)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0403] Similarly to Example 28, the title compound (36.1 mg, 0.072 mmol, 52%) was obtained as white crystals from N1-(4-fluorophenyl)-4-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (50 mg, 0.14 mmol) and 2-diethylaminoethylamine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.95 (6H, t, J=6.8

Hz), 2.30-2.50 (6H, m), 3.10-3.20 (2H, m), 6.91 (1H, d, J=3.6 Hz), 6.99 (1H, m), 7.18-7.26 (3H, m), 7.60-7.75 (4H, m), 7.90-8.00 (2H, m), 8.25 (1H, s), 9.24 (1H, s), 9.31 (1H, s), 10.09 (1H, brs).

5

Example 161

N1-Methyl-6-(6-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

10 [0404] Similarly to Example 28, the title compound (35.6 mg, 0.077 mmol, 43%) was obtained as white crystals from N1-methyl-6-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (51 mg, 0.18 mmol) and 4-(pyrrolidin-1-yl)piperidine.

15 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.20-1.40 (2H, m), 1.60-1.70 (4H, m), 1.70-1.85 (2H, m), 2.40-2.60 (5H, m), 2.81 (3H, d, J=4.0 Hz), 2.85-2.95 (2H, m), 3.90-4.10 (2H, m), 6.57 (1H, d, J=3.6 Hz), 7.23 (1H, s), 7.39-7.50 (2H, m), 7.68 (1H, d, J=3.6 Hz), 8.02 (1H, m),
20 8.26 (1H, s), 8.51 (1H, s), 9.13 (1H, s), 9.40 (1H, s).

[0405] The starting materials were synthesized by the following methods.

Production example 161-1

25 6-Chloro-4-(1H-6-indolylamino)pyrimidine

The title compound (1.229 g, 5.0 mmol, 46%) was

obtained as pale yellow crystals from 4,6-dichloropyrimidine (1.69 g, 11 mmol), 6-aminoindole and N,N-diisopropylethylamine by the method similar to the synthesis of 6-chloro-4-(1H-5-indolylamino)pyrimidine.

5 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 6.39 (1H, s), 6.74 (1H, s), 7.02 (1H, dd, J=2.8, 8.8 Hz), 7.30 (1H, t, J=2.8 Hz), 7.50 (1H, d, J=8.8 Hz), 7.83 (1H, m), 8.43 (1H, s), 9.78 (1H, s), 11.07 (1H, brs).

10 Production example 161-2

6-Amino-4-(1H-6-indolylamino)pyrimidine

[0406] A 7N ammonia in methanol solution (75 ml) was added to 6-chloro-4-(1H-6-indolylamino)pyrimidine (1.229 g, 5.0 mmol); and the reaction mixture was
15 heated in a sealed tube at 130°C for 6 days. The solvent was distilled off under reduced pressure; the residue was purified by silica gel column chromatography (eluent; ethyl acetate: tetrahydrofuran = 1: 1); diethyl ether was added to the residue to
20 crystallize; the crystals were filtered off, and washed with diethyl ether, and dried under aeration to yield the title compound (883 mg, 3.9 mmol, 78%) as pale yellow crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 5.71 (1H, s), 6.19 (2H, brs), 6.32 (1H, s), 6.92 (1H, dd, J=1.2, 8.0 Hz),
25 7.20 (1H, m), 7.40 (1H, d, J=8.0 Hz), 7.65 (1H, s),

7.98 (1H, s), 8.65 (1H, s), 10.91 (1H, s).

Production example 161-3

N1-Methyl-6-(6-aminopyrimidin-4-yl)amino-1H-1-
5 indolecarboxamide

[0407] Similarly to Production example 2-3, the title compound (105 mg, 0.37 mmol, 48%) was obtained as pale brown crystals from 6-amino-4-(1H-6-indolylamino)pyrimidine (175 mg, 0.78 mmol) and phenyl
10 N-methylcarbamate.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.80 (3H, d, J=4.4 Hz), 5.73 (1H, s), 6.24 (2H, brs), 6.56 (1H, d, J=2.8 Hz), 7.33 (1H, d, J=8.4 Hz), 7.44 (1H, d, J=8.4 Hz), 7.66 (1H, d, J=2.8 Hz), 7.90-8.10 (2H, m), 8.33 (1H, s),
15 8.84 (1H, s).

[0408]

Example 162

N1-Methyl-6-(6-((3-
diethylaminopropylamino)carbonyl)aminopyrimidin-4-
20 yl)amino-1H-1-indolecarboxamide

Similarly to Example 28, the title compound (37.3 mg, 0.085 mmol, 47%) was obtained as white crystals from N1-methyl-6-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (51 mg, 0.18 mmol) and 3-
25 diethylaminopropylamine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.92 (6H, t, J=6.8

Hz), 1.40-1.60 (2H, m), 2.20-2.40 (6H, m), 2.81 (3H, d, J=4.0 Hz), 3.05-3.20 (2H, m), 6.58 (1H, d, J=4.0 Hz), 6.83 (1H, s), 7.38 (1H, d, J=8.0 Hz), 7.46 (1H, d, J=8.0 Hz), 7.60-7.80 (2H, m), 8.03 (1H, m), 8.23 (1H, s), 8.46 (1H, s), 9.10 (1H, s), 9.43 (1H, s).

Example 163

N1-(2-Propyl)-6-(6-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0409] Similarly to Example 28, the title compound (40.4 mg, 0.082 mmol, 54%) was obtained as white crystals from N1-(2-propyl)-6-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (47 mg, 0.15 mmol) and 4-(pyrrolidin-1-yl)piperidine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.20 (6H, d, J=6.8 Hz), 1.20-1.36 (2H, m), 1.60-1.70 (4H, m), 1.70-1.85 (2H, m), 2.40-2.60 (5H, m), 2.80-3.00 (2H, m), 3.90-4.10 (3H, m), 6.56 (1H, d, J=3.6 Hz), 7.24 (1H, s), 7.30-7.50 (2H, m), 7.74 (1H, d, J=3.6 Hz), 7.81 (1H, d, J=8.8 Hz), 8.26 (1H, s), 8.50 (1H, s), 9.13 (1H, s), 9.39 (1H, s).

[0410] The starting material was synthesized by the following method.

Production example 163-1

N1-(2-Propyl)-6-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

Similarly to Production example 2-3, the title compound (95.3 mg, 0.31 mmol, 40%) was obtained as pale brown crystals from 6-amino-4-(1H-6-indolylamino)pyrimidine (175 mg, 0.78 mmol) and phenyl N-(2-propyl)carbamate.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.20 (6H, d, J=6.4 Hz), 4.00 (1H, m), 5.73 (1H, s), 6.24 (2H, brs), 6.55 (1H, d, J=3.6 Hz), 7.33 (1H, dd, J=2.0, 8.0 Hz), 7.44 (1H, d, J=8.0 Hz), 7.73 (1H, d, J=3.6 Hz), 7.80 (1H, d, J=8.0 Hz), 7.98 (1H, s), 8.33 (1H, s), 8.84 (1H, s).

Example 164

N1-(2-Propyl)-6-(6-((3-diethylaminopropylamino)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0411] Similarly to Example 28, the title compound (40.1 mg, 0.086 mmol, 57%) was obtained as white crystals from N1-(2-propyl)-6-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (47 mg, 0.15 mmol) and 3-diethylaminopropylamine.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.92 (6H, t, J=6.8 Hz), 1.20 (6H, d, J=6.4 Hz), 1.40-1.60 (2H, m), 2.20-2.50 (6H, m), 3.05-3.20 (2H, m), 4.00 (1H, m), 6.57 (1H, d, J=3.6 Hz), 6.84 (1H, s), 7.36 (1H, dd, J=8.0 Hz),

7.46 (1H, d, J=8.0 Hz), 7.12 (1H, m), 7.76 (1H, d, J=3.6 Hz), 7.82 (1H, d, J=8.0 Hz), 8.23 (1H, s), 8.44 (1H, s), 9.10 (1H, s), 9.43 (1H, s).

5 Example 165

N1-(4-Fluorophenyl)-6-(6-((3-diethylaminopropylamino)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

10 [0412] Similarly to Example 28, the title compound (22.9 mg, 0.044 mmol, 24%) was obtained as white crystals from N1-(4-fluorophenyl)-6-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (67 mg, 0.19 mmol) and 3-diethylaminopropylamine.

15 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.92 (6H, t, J=6.8 Hz), 1.40-1.60 (2H, m), 2.20-2.40 (6H, m), 3.05-3.20 (2H, m), 6.68 (1H, d, J=3.6 Hz), 6.87 (1H, s), 7.24 (2H, t, J=8.8 Hz), 7.43 (1H, dd, J=2.0, 8.4 Hz), 7.52 (1H, d, J=8.4 Hz), 7.60-7.80 (4H, m), 7.89 (1H, d, J=3.6 Hz), 8.23 (1H, s), 8.48 (1H, s), 9.11 (1H, s), 9.45 (1H, s).

20

[0413] The starting material was synthesized by the following method.

Production example 165-1

25 N1-(4-Fluorophenyl)-6-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

Similarly to Production example 2-3, the title

compound (137 mg, 0.38 mmol, 49%) was obtained as pale brown crystals from 6-amino-4-(1H-6-indolylamino)pyrimidine (175 mg, 0.78 mmol) and phenyl N-(4-fluorophenyl)carbamate.

5 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 5.75 (1H, s), 6.26 (2H, brs), 6.66 (1H, d, J=3.6 Hz), 7.22 (2H, t, J=8.8 Hz), 7.39 (1H, dd, J=2.0, 8.4 Hz), 7.49 (1H, d, J=8.4 Hz), 7.60-7.70 (2H, m), 7.87 (1H, d, J=3.6 Hz), 7.99 (1H, s), 8.36 (1H, s), 8.91 (1H, s), 10.01 (1H, s).

10

Example 166

N1-(4-Fluorophenyl)-6-(6-((2-diethylaminoethylamino)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

15 [0414] Similarly to Example 28, the title compound (11.1 mg, 0.022 mmol, 12%) was obtained as white crystals from N1-(4-fluorophenyl)-6-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (67 mg, 0.19 mmol) and 2-diethylaminoethylamine.

20 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.80-1.00 (6H, m), 2.20-2.50 (6H, m), 3.00-3.20 (2H, m), 6.74 (1H, s), 6.84 (1H, d, J=3.6 Hz), 7.03 (1H, d, J=8.0 Hz), 7.20 (2H, t, J=8.8 Hz), 7.50-7.70 (3H, m), 7.70 (1H, d, J=8.0 Hz), 8.00 (1H, s), 8.12 (1H, d, J=3.6 Hz), 8.37
25 (1H, s), 9.23 (1H, s), 9.41 (1H, s), 10.12 (1H, s).

Example 167N1-Dimethyl-6-(6-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

5 [0415] Similarly to Example 28, the title compound (16.1 mg, 0.034 mmol, 17%) was obtained as pale yellow powder from N1-dimethyl-6-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (58 mg, 0.20 mmol) and 4-(pyrrolidin-1-yl)piperidine.

10 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.20-1.36 (2H, m), 1.60-1.70 (4H, m), 1.70-1.85 (2H, m), 2.40-2.60 (5H, m), 2.80-3.00 (2H, m), 3.02 (6H, s), 3.90-4.10 (2H, m), 6.55 (1H, s), 7.26 (1H, s), 7.30 (1H, d, J=8.0 Hz), 7.40-7.50 (2H, m), 8.01 (1H, s), 8.29 (1H, s), 9.16 (1H, s), 9.41 (1H, s).

[0416] The starting material was synthesized by the following method.

Production example 167-1

20 N1-Dimethyl-6-(6-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

Similarly to Production example 2-3, the title compound (58.3 mg, 0.20 mmol, 25%) was obtained as pale brown crystals from 6-amino-4-(1H-6-indolylamino)pyrimidine (175 mg, 0.78 mmol) and phenyl N,N-dimethylcarbamate.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 3.01 (6H, s), 5.72 (1H, s), 6.26 (2H, brs), 6.54 (1H, d, J=3.6 Hz), 7.24 (1H, dd, J=2.0, 8.0 Hz), 7.40-7.50 (2H, m), 7.84 (1H, s), 8.01 (1H, s), 8.86 (1H, s).

5

Example 168

N1-Diethyl-5-(2-((pyrrolidin-1-ylamino)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

10 [0417] Similarly to Example 28, the title compound (91.0 mg, 0.22 mmol, 38%) was obtained as pale yellow powder from N1-diethyl-5-(2-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (186 mg, 0.57 mmol) and pyrrolidine.

15 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.15 (6H, t, J=6.8 Hz), 1.60-1.90 (4H, m), 3.20-3.50 (8H, m), 6.28 (1H, d, J=6.0 Hz), 6.52 (1H, d, J=3.6 Hz), 7.40-7.50 (3H, m), 7.96 (1H, d, J=6.0 Hz), 8.24 (1H, brs), 8.67 (1H, s), 9.35 (1H, s).

20

[0418] The starting materials were synthesized by the following methods.

Production example 168-1

2-Amino-4-(1H-5-indolylamino)pyrimidine

25 A mixture of 2-amino-4,6-dichloropyrimidine (1.64 g, 10 mmol), 5-aminoindole (1.32 g, 10 mmol),

diisopropylethylamine (5.23 ml, 30 mmol) and N,N-dimethylformamide (30 ml) was heated at 60°C and stirred overnight under nitrogen atmosphere. The reaction mixture was partitioned between ethyl acetate and water after cooled to room temperature; and the organic layer was washed with water and brine, and dried over anhydrous sodium sulfate. The solvent was distilled off; the obtained residue was dissolved in tetrahydrofuran (100 ml); 10% Palladium on carbon (50% wet, 1.0 g) was added thereto under nitrogen atmosphere; and the reaction mixture was stirred for 4 days under hydrogen atmosphere at atmospheric pressure. The reaction system was purged with nitrogen; the catalyst was filtered off; the filtrate was concentrated under reduced pressure; the residue was purified by NH silica gel column chromatography (eluent; ethyl acetate: methanol = 95: 5); diethyl ether was added to the residue to crystallize; and the crystals were filtered off, and dried under aeration to yield the title compound (852 mg, 3.8 mmol, 38%) as pale brown crystals.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 5.89 (1H, d, J=5.6 Hz), 5.99 (2H, brs), 6.34 (1H, s), 7.12 (1H, d, J=8.4 Hz), 7.20-7.40 (2H, m), 7.70 (1H, d, J=5.6 Hz), 7.79 (1H, s), 8.73 (1H, s), 10.95 (1H, s).

Production example 168-2N1-Diethyl-5-(2-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

5 [0419] Similarly to Production example 2-3, the title compound (1.22 g, 3.8 mmol, quantitative) was obtained as pale brown powder from 2-amino-4-(1H-5-indolylamino)pyrimidine (852 mg, 3.8 mmol) and diethylcarbamy l chloride.

10 ¹H-NMR Spectrum (CDCl₃) δ(ppm): 1.26 (6H, t, J=7.2 Hz), 3.49 (4H, q, J=7.2 Hz), 4.78 (2H, brs), 6.03 (1H, d, J=5.6 Hz), 6.57 (1H, d, J=3.6 Hz), 6.66 (1H, s), 7.16 (1H, dd, J=2.0, 8.8 Hz), 7.31 (1H, d, J=3.6 Hz), 7.53 (1H, d, J=2.0 Hz), 7.65 (1H, d, J=8.8 Hz), 7.88 (1H, d, J=5.6 Hz).

15

Example 169N1-Diethyl-5-(5-iodo-2-((pyrrolidin-1-ylamino)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

20 [0420] Similarly to Example 28, the title compound (37.3 mg, 0.068 mmol, 26%) was obtained as white powder from N1-diethyl-5-(2-amino-5-iodopyrimidin-4-yl)amino-1H-1-indolecarboxamide (117 mg, 0.26 mmol) and pyrrolidine.

25 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.15 (6H, t, J=6.8 Hz), 1.60-1.80 (4H, m), 3.30-3.50 (8H, m), 6.54 (1H, s),

7.30-7.60 (3H, m), 8.09 (1H, s), 8.15 (1H, s), 8.33 (1H, s), 8.82 (1H, s).

[0421] The starting material was synthesized by the following method.

Production example 169-1

N1-Diethyl-5-(2-amino-5-iodopyrimidin-4-yl)amino-1H-1-indolecarboxamide

N1-Diethyl-5-(2-aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide (1.06 g, 3.27 mmol) was dissolved in N,N-dimethylformamide (10 ml) under nitrogen atmosphere; N-iodosuccinimide (920 mg, 4.08 mmol) was added thereto while cooling with an ice water bath; and the reaction mixture was stirred at room temperature overnight. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was washed with water and brine, and dried over anhydrous sodium sulfate. The solvent was distilled off; and the residue was purified by silica gel column chromatography (eluent; ethyl acetate) to yield the title compound (1.00 g, 2.33 mmol, 68%) as yellow powder.

¹H-NMR Spectrum (CDCl₃) δ(ppm): 1.26 (6H, t, J=7.2 Hz), 3.49 (4H, q, J=7.2 Hz), 4.84 (2H, brs), 6.58 (1H, d, J=3.6 Hz), 6.95 (1H, s), 7.27-7.40 (2H, m), 7.63 (1H, d, J=8.8 Hz), 7.82 (1H, s), 8.16 (1H, s).

Example 170

N1-Diethyl-5-(5-cyano-2-((pyrrolidin-1-ylamino)carbonyl)aminopyrimidin-4-yl)amino-1H-1-indolecarboxamide

[0422] Similarly to Example 28, the title compound (35.3 mg, 0.079 mmol, 28%) was obtained as white crystals from N1-diethyl-5-(2-amino-5-cyanopyrimidin-4-yl)amino-1H-1-indolecarboxamide (100 mg, 0.29 mmol) and pyrrolidine.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.15 (6H, t, J=6.8 Hz), 1.60-1.80 (4H, m), 3.20-3.50 (8H, m), 6.56 (1H, s), 7.40-7.60 (3H, m), 8.03 (1H, s), 8.49 (1H, s), 9.43 (1H, s), 9.50 (1H, s).

[0423] The starting material was synthesized as follows.

Production example 170-1

N1-Diethyl-5-(2-amino-5-cyanopyrimidin-4-yl)amino-1H-1-indolecarboxamide

N1-Diethyl-5-(2-amino-5-iodopyrimidin-4-yl)amino-1H-1-indolecarboxamide (882 mg, 1.96 mmol) was dissolved in N,N-dimethylformamide (10 ml) under nitrogen atmosphere; zinc cyanide (253 mg, 2.15 mmol) and tetrakis(triphenylphosphine)palladium (226 mg, 0.2 mmol) was added thereto; the reaction mixture was stirred at 100 °C for 2 hours. The reaction mixture

was partitioned between ethyl acetate and water; and the organic layer was washed with water and brine, and dried over anhydrous sodium sulfate. The solvent was distilled off; and the residue was purified by silica gel column chromatography (eluent; ethyl acetate) to yield the title compound (493 mg, 1.41 mmol, 72%) as white crystals.

¹H-NMR Spectrum (CDCl₃) δ (ppm): 1.26 (6H, t, J=7.2 Hz), 3.49 (4H, q, J=7.2 Hz), 5.26 (2H, brs), 6.59 (1H, d, J=3.6 Hz), 7.05 (1H, s), 7.27-7.35 (2H, m), 7.66 (1H, d, J=8.8 Hz), 7.78 (1H, m), 8.27 (1H, s).

Example 171

5-(2-(3-Ethylureido)pyridin-4-yloxy)indole-1-carboxylic acid (2-diethylaminoethyl)amide

[0424] Similarly to Production example 5-1, a crude product of 5-(2-aminopyridin-4-yloxy)indole-1-carboxylic acid (2-diethylaminoethyl)amide (81 mg) was obtained as a pale yellow oil from 4-(1H-5-indolyloxy)-2-pyridinamine (225 mg, 1.00 mmol, WO 02/32872), sodium hydride (80 mg, 2.00 mmol, 60% in oil), and phenyl N-(2-diethylaminoethyl)carbamate (314 mg, 1.50 mmol). Similarly to Production example 5-2, a mixture of phenyl (4-(1-(2-diethylaminoethyl)carbamoyl-1H-indol-5-yloxy)pyridin-2-yl)-N-(phenoxycarbonyl)carbamate and phenyl (4-(1-(2-diethylaminoethyl)carbamoyl-1H-indol-5-

xyloxy)pyridin-2-yl)carbamate (32 mg) was obtained as a pale yellow oil from the crude product obtained above, phenyl chloroformate (0.041 ml, 0.33 mmol) and triethylamine (0.049 ml, 0.35 mmol). Similarly to
5 Example 5, the title compound (11 mg, 0.025 mmol, 12%) was obtained as pale yellow crystals from the mixture obtained above, ethylamine hydrochloride (30 mg, 0.26 mmol) and triethylamine (0.5 ml).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.97 (6H, t, J=7.0 Hz), 1.02 (3H, t, J=7.0 Hz), 2.44-2.60 (8H, m), 3.10 (2H, m), 6.50 (1H, dd, J=1.6, 6.0 Hz), 6.68 (1H, d, J=3.6 Hz), 6.86 (1H, d, J=1.6 Hz), 7.03 (1H, dd, J=2.0, 8.8 Hz), 7.36 (1H, d, J=2.0 Hz), 7.89 (1H, d, J=3.6 Hz), 7.97 (1H, m), 8.02 (1H, d, J=6.0 Hz), 8.17 (1H, m),
10 8.28 (1H, d, J=8.8 Hz), 9.00 (1H, s).
15

ESI-MS: 439.30 (M+H).

[0425]The starting material was synthesized as follows.

Production example 171-1

20 Phenyl N-(2-diethylaminoethyl)carbamate

Similarly to Production example 2-1, a crude product was obtained from 2-diethylaminoethylamine (7.3 ml, 50 mmol) and phenyl chloroformate (6.9 ml, 55 mmol). The crude product was purified by silica gel column chromatography (Fuji Silysia NH, ethyl acetate), and
25 further purified by silica gel column chromatography

(Fuji Silysia NH; hexane: ethyl acetate = 3: 1, 1: 1, ethyl acetate in this order) to yield the title compound (1.3 g, 6.4 mmol, 13%) as a colorless oil.

¹H-NMR Spectrum (CDCl₃) δ (ppm): 1.04 (6H, t, J=7.2 Hz),
5 2.52-2.62 (6H, m), 3.31 (2H, q, J=5.6 Hz), 5.62 (1H, brs), 7.13 (2H, d, J=7.6 Hz), 7.18 (1H, t, J=7.6 Hz),
7.35 (2H, t, J=7.6 Hz).

Example 172

10 5-(2-(3,3-Diethylureido)pyridin-4-yloxy)indole-1-carboxylic acid (2-ethoxyethyl)amide

[0426] Similarly to Production example 5-2, a mixture of phenyl (4-(1-(2-ethoxyethyl)carbamoyl-1H-indol-5-yloxy)pyridin-2-yl)-N-(phenoxycarbonyl)carbamate and
15 phenyl (4-(1-(2-ethoxyethyl)carbamoyl-1H-indol-5-yloxy)pyridin-2-yl)carbamate (3.42 g) was obtained as a pale yellow oil from 5-(2-aminopyridin-4-yloxy)indole-1-carboxylic acid (2-ethoxyethyl)amide (1.86 g, 5.46 mmol), phenyl chloroformate (1.51 ml, 12.0 mmol) and
20 triethylamine (1.90 ml, 13.7 mmol). Similarly to Example 5, the title compound was obtained as pale pink crystals (84 mg, 0.19 mmol) from this intermediates (174 mg) and diethylamine (0.16 ml, 1.5 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.01 (6H, t, J=7.2 Hz),
25 1.11 (3H, t, J=7.2 Hz), 3.26-3.31 (4H, m), 3.42-3.50 (4H, m), 3.53 (2H, m), 6.54 (1H, dd, J=2.4, 5.6

Hz), 6.68 (1H, d, J=3.6 Hz), 7.04 (1H, dd, J=2.4, 9.0 Hz), 7.36 (1H, d, J=2.4 Hz), 7.41 (1H, d, J=2.4 Hz), 7.93 (1H, d, J=3.6 Hz), 8.06 (1H, d, J=5.6 Hz), 8.28 (1H, d, J=9.0 Hz), 8.31 (1H, m), 8.60 (1H, s).

5 ESI-MS: 440.47 (M+H).

[0427] The starting materials were synthesized as follows.

Production example 172-1

10 Phenyl N-(2-ethoxyethyl)carbamate

Similarly to Example 5, a crude product was obtained from 2-ethoxyethylamine (5.2 ml, 50 mmol), phenyl chloroformate (6.9 ml, 55 mmol), and pyridine (4.5 ml, 55 mmol). The obtained crude product was
15 purified by silica gel column chromatography (Fuji Silysia BW-300, hexane: ethyl acetate = 85: 15 to 50: 50) to yield the title compound (8.38 g, 40.4 mmol, 80.9%) as a pale yellow oil.

¹H-NMR Spectrum (CDCl₃) δ (ppm): 1.23 (3H, t, J=7.0 Hz),
20 3.44-3.48 (2H, m), 3.52-3.58 (4H, m), 5.41 (1H, brs), 7.13 (2H, d, J=7.6 Hz), 7.19 (1H, t, J=7.6 Hz), 7.35 (2H, t, J=7.6 Hz).

Production example 172-2

25 5-(2-Aminopyridin-4-yloxy)indole-1-carboxylic acid (2-ethoxyethyl)amide

[0428] Similarly to Production example 5-1, the title compound (1.86 g, 5.46 mmol, 61.5%) was obtained as a pale brown oil from 4-(1H-5-indolyloxy)-2-pyridinamine (2.00 g, 8.88 mmol, WO 02/32872), sodium hydride (462 mg, 11.5 mmol, 60% in oil), and phenyl N-(2-ethoxyethyl)carbamate (2.23 g, 10.7 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.11 (3H, t, J=7.2 Hz), 3.42 (2H, m), 3.47 (2H, q, J=7.2 Hz), 3.53 (2H, t, J=6.0 Hz), 5.74 (1H, d, J=2.0 Hz), 5.84 (2H, m), 6.12 (1H, dd, J=2.0, 6.0 Hz), 6.67 (1H, d, J=3.8 Hz), 7.01 (1H, dd, J=2.0, 8.8 Hz), 7.33 (1H, d, J=2.0 Hz), 7.75 (1H, d, J=6.0 Hz), 7.91 (1H, d, J=6.0 Hz), 8.26 (1H, d, J=8.8 Hz), 8.28 (1H, m).

Example 173

5-(2-(3-Ethylureido)pyridin-4-yloxy)indole-1-carboxylic acid (2-ethoxyethyl)amide

[0429] Similarly to Example 5, the title compound was obtained as colorless crystals (84 mg, 0.204 mmol) from a mixture (174 mg) of phenyl (4-(1-(2-ethoxyethyl)carbamoyl-1H-indol-5-yloxy)pyridin-2-yl)-N-(phenoxy-carbonyl)carbamate and phenyl (4-(1-(2-ethoxyethyl)carbamoyl-1H-indol-5-yloxy)pyridin-2-yl)carbamate, which was obtained as an intermediate in Example 172, ethylamine hydrochloride (122 mg, 1.50 mmol) and triethylamine (0.5 ml).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.02 (3H, t, J=7.0 Hz), 1.12 (3H, t, J=7.0 Hz), 3.10 (2H, m), 3.40-3.49 (4H, m), 3.53 (2H, t, J=5.8 Hz), 6.50 (1H, dd, J=2.4, 5.8 Hz), 6.68 (1H, d, J=3.6 Hz), 6.86 (1H, d, J=2.4 Hz), 7.04 (1H, dd, J=2.4, 8.8 Hz), 7.37 (1H, d, J=2.4 Hz), 7.93 (1H, d, J=3.6 Hz), 7.96 (1H, m), 8.02 (1H, d, J=5.8 Hz), 8.28 (1H, d, J=8.8 Hz), 8.31 (1H, m), 9.00 (1H, s).

ESI-MS: 412.18 (M+H).

Example 174

5-(2-(3-Ethylureido)pyridin-4-yloxy)indole-1-carboxylic acid (3-ethoxypropyl)amide

[0430] Similarly to Production example 5-2, a mixture of phenyl (4-(1-(3-ethoxypropyl)carbamoyl-1H-indol-5-yloxy)pyridin-2-yl)-N-(phenoxycarbonyl)carbamate and phenyl (4-(1-(3-ethoxypropyl)carbamoyl-1H-indol-5-yloxy)pyridin-2-yl)carbamate was obtained as a pale brown oil (720 mg) from 5-(2-Aminopyridin-4-yloxy)indole-1-carboxylic acid (3-ethoxypropyl)amide (900 mg, 2.54 mmol), phenyl chloroformate (0.669 ml, 5.33 mmol) and triethylamine (0.885 ml, 6.35 mmol). Similarly to Example 5, the title compound was obtained as pale pink crystals (41 mg, 0.096 mmol) from this intermediate (100 mg) ethylamine hydrochloride (64 mg, 0.841 mmol) and triethylamine (0.5 ml).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.02 (3H, t, J=7.2 Hz), 1.10 (3H, t, J=7.2 Hz), 1.79 (2H, m), 3.10 (2H, m), 3.34 (2H, m), 3.42 (4H, m), 6.50 (1H, dd, J=2.4, 5.8 Hz), 6.67 (1H, d, J=3.8 Hz), 6.86 (1H, d, J=2.4 Hz),
5 7.03 (1H, dd, J=2.4, 9.2 Hz), 7.36 (1H, d, J=2.4 Hz), 7.90 (1H, d, J=3.8 Hz), 7.95 (1H, m), 8.02 (1H, d, J=5.8 Hz), 8.20 (1H, m), 8.28 (1H, m), 8.99 (1H, s).
ESI-MS: 426.39 (M+H).

10 [0431]The starting material was synthesized as follows.
Production example 174-1
5-(2-Aminopyridin-4-yloxy)indol-1-carboxylic acid (3-ethoxypropyl)amide

Similarly to Production example 5-1, the title
15 compound (900 mg, 2.54 mmol, 57.2%) was obtained as a pale brown oil from 4-(1H-5-indolyloxy)-2-pyridinamine (1.00 g, 4.44 mmol, WO 02/32872), sodium hydride (213 mg, 5.33 mmol, 60% in oil), and phenyl N-(3-ethoxypropyl)carbamate (1.19 g, 5.33 mmol, WO 02/32872).

20 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.07-1.13 (3H, m), 1.81 (2H, m), 3.33-3.47 (6H, m), 5.76 (1H, d, J=2.4 Hz), 5.85 (2H, s), 6.14 (1H, dd, J=2.4, 6.0 Hz), 6.68 (1H, d, J=3.6 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.34 (1H, d, J=2.4 Hz), 7.77 (1H, d, J=6.0 Hz), 7.90 (1H, d, J=3.6
25 Hz), 8.20 (1H, m), 8.27 (1H, d, J=8.8 Hz).

Example 1755-(2-(3-Ethylureido)pyridin-4-yloxy)indole-1-carboxylic acid (3-methylsulphanylpropyl)amide

[0432] Similarly to Production example 5-1, a crude product of 5-(2-aminopyridin-4-yloxy)indole-1-carboxylic acid (3-methylsulfanylpropyl)amide (105 mg) was obtained as a pale yellow oil from 4-(1H-5-indolyloxy)-2-pyridinamine (125 mg, 0.555 mmol, WO 02/32872), sodium hydride (28 mg, 0.694 mmol, 60% in oil), and phenyl N-(3-methylsulfanylpropyl)carbamate (156 mg, 0.694 mmol, WO 02/32872). Similarly to Production example 5-2, a mixture of phenyl (4-(1-(3-methylsulfanylpropyl)carbamoyl-1H-indol-5-yloxy)pyridin-2-yl)-N-(phenoxycarbonyl)carbamate and phenyl (4-(1-(3-methylsulfanylpropyl)carbamoyl-1H-indol-5-yloxy)pyridin-2-yl)carbamate was obtained as a pale yellow oil from the crude product obtained above, phenyl chloroformate (0.10 ml, 0.76 mmol) and triethylamine (0.12 ml, 0.83 mmol). Similarly to Example 5, the title compound (17 mg, 0.040 mmol) was obtained as colorless crystals from the mixture obtained above, ethylamine hydrochloride (141 mg, 1.73 mmol) and triethylamine (0.5 ml).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.02 (3H, t, J=7.2 Hz), 1.83 (2H, m), 2.05 (3H, s), 2.52 (2H, t, J=7.6 Hz), 3.10 (2H, m), 3.35 (2H, m), 6.50 (1H, d, J=5.6 Hz),

6.68 (1H, d, J=3.4 Hz), 6.86 (1H, s), 7.03 (1H, d, J=9.2 Hz), 7.36 (1H, s), 7.91 (1H, d, J=3.4 Hz), 7.94 (1H, m), 8.02 (1H, d, J=5.6 Hz), 8.24 (1H, m), 8.27 (1H, d, J=9.2 Hz), 8.99 (1H, s).

5

Example 176

5-(2-(3,3-Diethylureido)pyridin-4-yloxy)indole-1-carboxylic acid thiazol-2-ylamide

[0433] Similarly to Production example 5-2, a mixture
10 (267 mg) of phenyl (4-(1-(thiazol-2-yl)carbamoyl-1H-indol-5-yloxy)pyridin-2-yl)-N-(phenoxycarbonyl)carbamate and phenyl (4-(1-(thiazol-2-yl)carbamoyl-1H-indol-5-yloxy)pyridin-2-yl)carbamate was obtained as a pale yellow oil from 5-(2-aminopyridin-4-yloxy)indole-1-carboxylic acid thiazol-2-ylamide (145 mg, 0.413 mmol), phenyl chloroformate (0.110 ml, 0.909 mmol), and triethylamine (0.140 ml, 1.03 mmol). Similarly to Example 5, the title compound (74 mg, 0.16 mmol) was obtained as pale pink crystals
15 from the intermediate obtained above (131 mg) and diethylamine (0.120 ml, 1.11 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.01 (6H, t, J=6.8 Hz), 3.28 (4H, m), 6.56 (1H, dd, J=2.0, 5.6 Hz), 6.66 (1H, m), 7.06 (2H, m), 7.37 (1H, d, J=2.0 Hz), 7.43 (1H, s), 7.47 (1H, d, J=4.4 Hz), 8.05 (1H, m), 8.07 (1H, d, J=5.6 Hz), 8.60 (2H, m).
25

ESI-MS: 451.15 (M+H).

[0434]The starting material was synthesized as follows.

Production example 176-1

5 5-(2-Aminopyridin-4-yloxy)indole-1-carboxylic acid
 thiazol-2-ylamide

 Similarly to Production example 5-1, the title
compound (145 mg, 0.413 mmol, 57.2%) was obtained as a
pale brown oil from 4-(1H-5-indolyloxy)-2-pyridinamine
10 (225 mg, 1.00 mmol, WO 02/32872), sodium hydride (120
mg, 3.00 mmol, 60% in oil) and phenyl N-(thiazol-2-
yl)carbamate (286 mg, 1.30 mmol, WO 02/32872).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 5.77 (1H, d, J=2.4
Hz), 5.87 (2H, brs), 6.15 (1H, dd, J=2.4, 5.6 Hz), 6.65
15 (1H, d, J=3.6 Hz), 7.03 (1H, dd, J=2.4, 9.0 Hz), 7.07
 (1H, d, J=4.6 Hz), 7.34 (1H, d, J=2.4 Hz), 7.46 (1H, d,
J=4.6 Hz), 7.77 (1H, d, J=5.6 Hz), 8.04 (1H, d, J=3.6
Hz), 8.58 (1H, d, J=9.0 Hz).

20 Example 177

5-(2-(3-Ethylureido)pyridin-4-yloxy)indole-1-carboxylic
acid thiazol-2-ylamide

 [0435] Similarly to Example 5, the title compound (71
mg, 0.168 mmol) was obtained as colorless crystals from
25 a mixture (135 mg) of phenyl (4-(1-(thiazol-2-
yl)carbamoyl-1H-indol-5-yloxy)pyridin-2-yl)-N-

(phenoxycarbonyl)carbamate and phenyl (4-(1-(thiazol-2-yl)carbamoyl-1H-indol-5-yloxy)pyridin-2-yl)carbamate obtained in Example 176, ethylamine hydrochloride (91 mg, 1.1 mmol), and triethylamine (0.5 ml).

5 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.07 (3H, t, J=7.2 Hz), 3.07-3.14 (2H, m), 6.51 (1H, dd, J=2.0, 6.0 Hz), 6.61 (1H, s), 7.01 (2H, m), 7.35 (1H, s), 7.41 (1H, m), 8.01-8.06 (3H, m), 8.05 (1H, m), 8.62 (1H, d, J=9.2 Hz), 9.00 (1H, s).

10 ESI-MS: 423.23 (M+H).

Example 178

1-Ethyl-3-(4-(1-((4-methylpiperazin-1-yl)carbonyl)-1H-indol-5-yloxy)pyridin-2-yl)urea

15 [0436] Similarly to Production example 5-2, a mixture (1.09 g) of phenyl (4-(1-((4-methylpiperazin-1-yl)carbonyl)-1H-indol-5-yloxy)pyridin-2-yl)-(N-phenoxycarbonyl)carbamate and phenyl (4-(1-((4-methylpiperazin-1-yl)carbonyl)-1H-indol-5-yloxy)pyridin-2-yl)carbamate was obtained as a colorless amorphous solid from (5-(2-aminopyridin-4-yloxy)indol-1-yl)-(4-methylpiperazin-1-yl)methanone (0.66 g, 1.9 mmol), phenyl chloroformate (0.52 ml, 4.2 mmol), and triethylamine (0.66 ml, 4.8 mmol).

20

25 Similarly to Example 5, the title compound (41 mg, 0.097 mmol) was obtained as colorless crystals from the

intermediate obtained above (177 mg), ethylamine hydrochloride (0.122 g, 1.50 mmol) and triethylamine (0.5 ml).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.03 (3H, t, J=6.0 Hz), 2.21 (3H, s), 2.39 (4H, m), 3.12 (2H, m), 3.51 (4H, m), 6.51 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.4 Hz), 6.86 (1H, d, J=2.4 Hz), 7.04 (1H, dd, J=2.4, 8.8 Hz), 7.39 (1H, d, J=2.4 Hz), 7.61 (1H, d, J=3.4 Hz), 7.70 (1H, d, J=8.8 Hz), 8.02 (1H, m), 8.03 (1H, d, J=8.8 Hz), 9.00 (1H, s).

ESI-MS: 423.27(M+H).

[0437] The starting materials were synthesized as follows.

15 Production example 178-1

Phenyl (4-methylpiperazin-1-yl)carboxylate

Similarly to Production example 2-1, crystals were obtained from 1-methylpiperazine (5.5 ml, 50 mmol), phenyl chloroformate (6.9 ml, 55 mmol), and pyridine (4.5ml, 55 mmol). The obtained crystals were suspended in diethylether: hexane = 2: 1, filtered off, washed with hexane, and dried to yield the title compound (9.7 g, 44 mmol, 88%) as an oil with pale orange color.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.20 (3H, s), 2.34 (4H, m), 3.40 (2H, m), 3.56 (2H, m), 7.09 (2H, d, J=7.6 Hz), 7.20 (1H, t, J=7.6 Hz), 7.36 (2H, t, J=7.6 Hz).

Production example 178-2(5-(2-Aminopyridin-4-yloxy)indol-1-yl)-(4-methylpiperazin-1-yl)methanone

- 5 [0438] Similarly to Production example 5-1, a crude product was obtained from 4-(1H-5-indolyloxy)-2-pyridinamine (2.00 g, 8.88 mmol, WO 02/32872), sodium hydride (462 mg, 11.5 mmol, 60% in oil) and phenyl (4-methylpiperazin-1-yl)carboxylate (2.35 g, 10.7 mmol).
- 10 The obtained crude product was purified by silica gel column chromatography (Fuji Silysia NH; hexane: ethyl acetate = 3: 7, ethyl acetate, ethyl acetate: methanol = 9: 1 in this order) to yield the title compound (0.66 g, 1.9 mmol, 21%) as a colorless amorphous solid.
- 15 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.20 (3H, s), 2.39 (4H, m), 3.51 (4H, m), 5.75 (1H, d, J=2.0 Hz), 5.84 (2H, m), 6.13 (1H, dd, J=2.0, 6.0 Hz), 6.66 (1H, d, J=3.2 Hz), 7.02 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, d, J=2.4 Hz), 7.59 (1H, d, J=3.2 Hz), 7.68 (1H, d, J=8.8 Hz),
- 20 7.76 (1H, d, J=6.0 Hz).

Example 1791-Ethyl-3-(4-(1-(morpholin-4-ylcarbonyl)-1H-indol-5-yloxy)pyridin-2-yl)urea

- 25 [0439] Similarly to Production example 5-2, a crude product was obtained from 5-(2-aminopyridin-4-

5 yloxy)indol-1-yl)-(morpholin-4-yl)methanone (0.60 g,
1.8 mmol), phenyl chloroformate (0.49 ml, 3.9 mmol),
and triethylamine (0.62 ml, 4.4 mmol). The obtained
crude product was filtrated by silica gel filtration
5 (Fuji Silysia BW-300, ethyl acetate) and concentrated
under reduced pressure to yield a mixture (1.11 g) of
phenyl (4-(1-(morpholin-4-ylcarbonyl)-1H-indol-5-
yloxy)pyridin-2-yl)-N-(phenoxycarbonyl)carbamate and
phenyl (4-(1-(morpholine-4-ylcarbonyl)-1H-indol-5-
10 yloxy)pyridin-2-yl)carbamate as a pale yellow oil.
Similarly to Example 5, the title compound (73 mg,
0.178 mmol) was obtained as colorless crystals from the
intermediate obtained above (173 mg), ethylamine
hydrochloride (122 mg, 1.50 mmol) and triethylamine
15 (0.5 ml).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.03 (3H, t, J=7.2
Hz), 3.07-3.14 (2H, m), 3.52 (4H, m), 3.68 (4H, m),
6.50 (1H, dd, J=2.4, 6.0 Hz), 6.68 (1H, d, J=3.2 Hz),
6.87 (1H, d, J=2.4 Hz), 7.05 (1H, dd, J=2.4, 8.8 Hz),
20 7.40 (1H, d, J=2.4 Hz), 7.64 (1H, d, J=3.2 Hz), 7.73
(1H, d, J=8.8 Hz), 8.00 (1H, m), 8.03 (1H, d, J=6.0 Hz),
9.01 (1H, s).

ESI-MS: 410.57 (M+H).

[0440]

25 The starting materials were synthesized as
follows.

Production example 179-1Phenyl (morpholin-4-yl)carboxylate

Similarly to Production example 2-1, a crude product was obtained from morpholine (4.4 ml, 50 mmol), phenyl chloroformate (6.9 ml, 55 mmol), and pyridine (4.5 ml, 55 mmol). The obtained crude product was purified by silica gel column chromatography (Fuji Silysia BW-300; hexane: ethyl acetate= 85: 15, 60: 40 in this order) to yield the title compound (8.9 g, 43 mmol, 86%) as colorless crystals.

¹H-NMR Spectrum (CDCl₃) δ (ppm): 3.57 (2H, brs), 3.68 (2H, brs), 3.75 (4H, m), 7.11 (2H, d, J=7.6 Hz), 7.21 (1H, t, J=7.6 Hz), 7.37 (2H, t, J=7.6 Hz).

[0441]

Production example 179-2(5-(2-Aminopyridin-4-yloxy)indol-1-yl)-(morpholin-4-yl)methanone

Similarly to Production example 5-1, a crude product was obtained from 4-(1H-5-indolyloxy)-2-pyridinamine (2.00 g, 8.88 mmol, WO 02/32872), sodium hydride (462 mg, 11.5 mmol, 60% in oil) and phenyl (morpholin-4-yl)carboxylate (2.21 g, 10.7 mmol). The obtained crude product was purified by silica gel column chromatography (Fuji Silysia NH, hexane: ethyl acetate = 2: 3 or ethyl acetate), and further purified by silica gel column chromatography (Fuji Silysia BW-

300, hexane: ethyl acetate = 2: 3, ethyl acetate, or ethyl acetate: methanol = 9: 1) to yield the title compound (0.60 g, 1.8 mmol, 20%) as colorless crystals.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 3.52 (4H, m), 3.68 (4H, m), 5.77 (1H, d, J=2.4 Hz), 5.83 (2H, brs), 6.13 (1H, dd, J=2.4, 5.6 Hz), 6.67 (1H, d, J=3.2 Hz), 7.02 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, d, J=2.4 Hz), 7.61 (1H, d, J=3.2 Hz), 7.71 (1H, d, J=8.8 Hz), 7.76 (1H, d, J=5.6 Hz).

Example 180

1,1-Diethyl-3-(4-(1-(morpholin-4-ylcarbonyl)-1H-indol-5-yloxy)pyridin-2-yl)urea

[0442] Similarly to Example 5, the title compound (85 mg, 0.194 mmol) was obtained as colorless crystals from a mixture (173 mg) of phenyl (4-(1-(morpholin-4-ylcarbonyl)-1H-indol-5-yloxy)pyridin-2-yl)-N-(phenoxy carbonyl) carbamate and phenyl (4-(1-(morpholin-4-ylcarbonyl)-1H-indol-5-yloxy)pyridin-2-yl) carbamate synthesized as intermediate in Example 179, and diethylamine (0.16 ml, 1.50 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.01 (6H, t, J=6.8 Hz), 3.30 (4H, m), 3.53 (4H, m), 3.68 (4H, m), 6.54 (1H, d, J=6.0 Hz), 6.68 (1H, d, J=3.4 Hz), 7.05 (1H, dd, J=2.0, 8.8 Hz), 7.39 (1H, d, J=2.0 Hz), 7.43 (1H, s), 7.64 (1H, d, J=3.4 Hz), 7.73 (1H, d, J=8.8 Hz), 8.07

(1H, d, J=6.0 Hz), 8.62 (1H, s).

ESI-MS: 438.25 (M+H).

Example 181

5 5-(2-(3-Ethylureido)pyridin-4-yloxy)indole-1-carboxylic
 acid piperidin-4-ylamide

[0443] Similarly to Production example 5-1, a crude
product of t-butyl 4-((5-(2-aminopyridin-4-
yloxy)indole-1-carbonyl)amino)piperidine-1-carboxylate
10 was obtained from 4-(1H-5-indolyloxy)-2-pyridinamine
(144 mg, 0.639 mmol, WO 02/32872), sodium hydride (29
mg, 0.735 mmol, 60% in oil), and t-butyl (4-
phenoxycarbonylaminopiperidin-1-yl)carboxylate (215 mg,
0.671 mmol). A reaction similar to Production example
15 5-2 was performed using the entire amount of this crude
product, phenyl chloroformate (0.20 ml, 1.6 mmol) and
triethylamine (0.22 ml); and the solvent was distilled
off under reduced pressure after the reaction was
completed. A reaction similar to Example 5 was
20 performed using the entire amount of the residue,
ethylamine hydrochloride (260 mg, 3.92 mmol) and
triethylamine (0.5 ml); the organic layer was
partitioned between ethyl acetate and water, washed
with brine, and dried over anhydrous sodium sulfate;
25 and the solvent was distilled off under reduced
pressure. The residue was dissolved in

trifluoroacetate (3.0 ml); the reaction mixture was stirred at room temperature for 15 minutes, and concentrated; and the residue was partitioned between ethyl acetate and water. The organic layer was washed with brine and dried over anhydrous sodium sulfate; the solvent was distilled off under reduced pressure; the residue was purified by silica gel column chromatography (Fuji Silysia NH, ethyl acetate: methanol = 98: 2 to 75: 25). The obtained crystals were suspended in diethyl ether and filtered off, washed with diethyl ether, and dried to yield the title compound (43 mg, 0.10 mmol, 16%) as colorless crystals.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.02 (3H, t, J=7.2 Hz), 1.37-1.49 (2H, m), 1.80 (2H, m), 2.48 (2H, m), 2.95 (2H, m), 3.10 (2H, m), 3.71 (1H, m), 6.50 (1H, dd, J=2.4, 6.0 Hz), 6.66 (1H, d, J=3.4 Hz), 6.86 (1H, d, J=2.4 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, d, J=2.4 Hz), 7.90-8.01 (3H, m), 8.02 (1H, d, J=6.0 Hz), 8.26 (1H, d, J=8.8 Hz), 8.99 (1H, s).

ESI-MS: 423.26 (M+H).

[0444]The starting material was synthesized as follows.

Production example 181-1

t-Butyl (4-phenoxyaminopiperidin-1-yl)carboxylate

A reaction similar to Production example 2-1

using t-Butyl 4-aminopiperidin-1-ylcarboxylate (328 mg, 1.64 mmol), phenyl chloroformate (0.226 ml, 1.80 mmol) and pyridine (0.146 ml, 1.80 mmol); and the obtained crystals were suspended in hexane: ethyl acetate = 4: 1, filtered off, and the filtrate was purified by silica gel column chromatography (Fuji Silysia BW-300, hexane: ethyl acetate = 4: 1 to 1: 1). The purified crystals were then suspended in hexane: ethyl acetate = 4: 1 and filtered off. The title compound (215 mg, 0.617 mmol, 40.9%) was obtained as colorless crystals, together with the previously obtained crystals.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.22-1.34 (2H, m), 1.38 (9H, s), 1.77 (2H, m), 2.83 (2H, m), 3.51 (1H, m), 3.84 (2H, m), 7.08 (2H, d, J=7.6 Hz), 7.18 (1H, t, J=7.6 Hz), 7.35 (2H, t, J=7.6 Hz), 7.78 (1H, d, J=8.0 Hz).

ESI-MS: 343.15 (M+Na).

Example 182

5-(2-(3-Ethylureido)pyridin-4-yloxy)indole-1-carboxylic acid (1-methylpiperidin-4-yl)amide

[0445] 5-(2-(3-Ethylureido)pyridin-4-yloxy)indole-1-carboxylic acid piperidin-4-ylamide (36 mg, 0.085 mmol, Example 181) was dissolved in tetrahydrofuran (2.0 ml) and methanol (1.0 ml); and a 37% aqueous formaldehyde solution (0.036 ml, 0.43 mmol) and acetic acid (0.0098

ml, 0.17 mmol) were added thereto. While stirring at room temperature, sodium triacetoxyborohydride (27 mg, 0.13 mmol) was added; and the reaction mixture was stirred for 30 minutes. The reaction mixture was partitioned between ethyl acetate and a saturated aqueous solution of sodium hydrogencarbonate; and the organic layer was washed with brine and dried over anhydrous sodium sulfate. The solvent was distilled off under reduced pressure; the obtained crystals were suspended in diethyl ether, filtered off, washed with diethyl ether, and dried to yield the title compound (25 mg, 0.057 mmol, 67%) as colorless crystals.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.02 (3H, t, J=7.2 Hz), 1.54-1.68 (2H, m), 1.83 (2H, m), 1.96 (2H, m), 2.16 (3H, s), 2.78 (2H, m), 3.10 (2H, m), 3.64 (1H, m), 6.50 (1H, dd, J=2.4, 6.0 Hz), 6.66 (1H, d, J=3.6 Hz), 6.86 (1H, d, J=2.4 Hz), 7.03 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, d, J=2.4 Hz), 7.95 (1H, d, J=3.6 Hz), 7.97 (2H, m), 8.02 (1H, d, J=6.0 Hz), 8.25 (1H, d, J=8.8 Hz), 8.99 (1H, m).

ESI-MS: 437.37 (M+H).

Example 183

5-(2-(N-Methyl-(4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)indole-1-carboxylic acid methanamide

[0446] 5-(2-(Methylamino)pyridin-4-yloxy)indole-1-carboxylic acid methylamide (70 mg, 0.24 mmol) was dissolved in tetrahydrofuran (7.0 ml); triethylamine (0.039 ml) and 4-nitrophenylchloroformate (57 mg, 0.28 mmol) were added thereto one by one; and the reaction mixture was stirred at room temperature for 2 hours. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was washed with a saturated aqueous solution of sodium hydrogencarbonate and brine, then dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was dissolved in N,N-dimethylformamide (2.0 ml); 4-(pyrrolidin-1-yl)piperidine (43 mg, 0.28 mmol) was added thereto; and the reaction mixture was stirred at room temperature for 24 hours. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Fuji Silysia NH, hexane-ethyl acetate-methanol system); the obtained oil was solidified with hexane; the obtained solid was then suspended in hexane, filtered off, washed with hexane, and dried to yield the title compound (51 mg, 0.11 mmol, 45%) as pale yellow crystals.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.13-1.30 (2H, m),

1.65 (6H, m), 2.02 (1H, m), 2.42 (4H, m), 2.72 (2H, m),
2.83 (3H, d, J=4.0 Hz), 3.08 (3H, s), 3.53 (2H, m),
6.23 (1H, s), 6.51 (1H, d, J=6.0 Hz), 6.66 (1H, d,
J=3.4 Hz), 7.04 (1H, d, J=9.0 Hz), 7.36 (1H, s), 7.87
5 (1H, d, J=3.4 Hz), 8.11 (1H, d, J=6.0 Hz), 8.15 (1H, m),
8.29 (1H, d, J=9.0 Hz).

ESI-MS: 477.38 (M+H).

[0447]The starting material was synthesized as follows.

10 Production example 183-1

5-(2-(Methylamino)pyridin-4-yloxy)indole-1-carboxylic
acid methanamide

N1-Methyl-5-(2-aminopyridin-4-yl)oxy-1H-1-
indolecarboxamide (5.00 g, 17.7 mmol, Production
15 example 5-1) was dissolved in ethanol (170 ml) and N,N-
dimethylformamide (40 ml); 1H-benzotriazole-1-methanol
(2.64 g, 17.7 mmol) was added thereto; and the reaction
mixture was heated to reflux for 2 hours. After
allowing to be cooled to room temperature, sodium
20 borohydride (1.49 g, 35.4 mmol) was added to the
reaction mixture; the reaction mixture was stirred at
room temperature for 2 hours. The reaction mixture was
partitioned between ethyl acetate and water; and the
organic layer was washed with brine, dried over
25 anhydrous sodium sulfate, and concentrated under
reduced pressure. The residue was purified by silica

gel column chromatography (Fuji Silysia BW-300, hexane-ethyl acetate-methanol system). The obtained crystals were suspended in acetone: diethyl ether = 1: 3, filtered off, washed with hexane, and dried to yield
5 the title compound (1.05 g, 3.55 mmol, 20.1%) as pale yellow crystals.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.66 (3H, d, J=4.8 Hz), 2.82 (3H, d, J=4.0 Hz), 5.76 (1H, d, J=2.0 Hz), 6.10 (1H, dd, J=2.0, 6.0 Hz), 6.36 (1H, m), 6.65 (1H, d, J=4.0 Hz),
10 7.00 (1H, dd, J=2.4, 8.8 Hz), 7.31 (1H, d, J=2.4 Hz), 7.83 (2H, m), 8.13 (1H, m), 8.26 (1H, d, J=8.8 Hz).

Example 184

15 5-(2-(1-Methylureido)pyridin-4-yloxy)indole-1-carboxylic acid methylamide

[0448] 4-Nitrophenyl N-methyl-(4-(1-methylcarbamoyl-indol-5-yloxy)pyridin-2-yl)carbamate (105 mg, 0.228 mmol) was dissolved in N,N-dimethylformamide (2.5 ml);
20 aqueous ammonia (0.5 ml, 28.0%) was added thereto; the reaction mixture was stirred at room temperature for 10.5 hours. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was washed with brine, dried over anhydrous sodium
25 sulfate, and concentrated under reduced pressure. The obtained crystals were suspended in ethanol: diethyl

ether = 1: 1 (6 ml), filtered off, washed with diethyl ether, and dried to yield the title compound (37 mg, 0.11 mmol, 48%) as pale yellow crystals.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.83 (3H, d, J=4.0 Hz), 3.19 (3H, s), 6.51 (1H, d, J=5.6 Hz), 6.67 (1H, d, J=3.6 Hz), 6.84 (1H, s), 7.07 (1H, d, J=9.0 Hz), 7.39 (1H, s), 7.87 (1H, d, J=3.6 Hz), 8.14 (2H, m), 8.29 (1H, d, J=9.0 Hz).

ESI-MS: 340.07 (M+H).

10

[0449]The starting material was synthesized as follows.

Production example 184-1

4-Nitrophenyl N-methyl-(4-(1-methylcarbamoyl-indol-5-yloxy)pyridin-2-yl)carbamate

15

5-(2-(Methylamino)pyridin-4-yloxy)indole-1-carboxylic acid methylamide (200 mg, 0.675 mmol) synthesized in Production example 183-1 was dissolved in tetrahydrofuran (20 ml); triethylamine (0.100 ml, 0.742 mmol) and 4-nitrophenylchloroformate (150 mg, 0.742 mmol) was added thereto one by one; and the reaction mixture was stirred at room temperature for 2 hours. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was washed with a saturated aqueous solution of sodium hydrogencarbonate and with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure.

25

The residue was purified by silica gel column chromatography (Fuji Silysia BW-300, hexane-ethyl acetate system) to yield the title compound (210 mg, 0.455 mmol, 67.4%) as a pale yellow oil.

5 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.82 (3H, d, J=4.4 Hz), 3.46 (3H, s), 6.62 (1H, d, J=3.6 Hz), 6.83 (1H, dd, J=2.0, 5.6 Hz), 7.03 (1H, dd, J=2.0, 8.4 Hz), 7.24 (1H, d, J=2.0 Hz), 7.37 (1H, d, J=2.0 Hz), 7.41 (2H, d, J=9.2 Hz), 7.85 (1H, d, J=3.6 Hz), 8.14 (1H, m), 8.22
10 (2H, d, J=9.2 Hz), 8.23 (1H, d, J=8.4 Hz), 8.32 (1H, d, J=5.6 Hz).

Example 185

15 5-(2-(3,3-Diethyl-1-methylureido)pyridin-4-yloxy)indole-1-carboxylic acid methylamide

[0450] Similarly to Example 184, the title compound (14 mg, 0.035 mmol, 16%) was obtained as a colorless amorphous solid from 4-nitrophenyl N-methyl-(4-(1-methylcarbamoyl-indol-5-yloxy)pyridin-2-yl)carbamate
20 (105 mg, 0.228 mmol, Production example 184-1) and diethylamine (0.028 ml, 0.27 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.80 (6H, t, J=6.4 Hz), 2.83 (3H, d, J=3.2 Hz), 3.03 (3H, s), 3.07 (4H, m), 6.11 (1H, s), 6.49 (1H, m), 6.66 (1H, s), 7.02 (1H, d, J=9.0 Hz), 7.35 (1H, s), 7.86 (1H, s), 8.09 (1H, d, J=5.6 Hz), 8.15 (1H, m), 8.28 (1H, d, J=9.0 Hz).

25

ESI-MS: 396.18 (M+H).

Example 186

5-(2-(3-Ethyl-1-methylureido)pyridin-4-yloxy)indole-1-
5 carboxylic acid methylamide

[0451] Similarly to Production example 27-2, phenyl N-methyl-(4-(1-methylcarbamoyl-indol-5-yloxy)pyridin-2-yl)carbamate (324 mg, 0.778 mmol) was obtained as a colorless amorphous solid from 5-(2-(methylamino)pyridin-4-yloxy)indole-1-carboxylic acid methylamide (500 mg, 1.69 mmol), phenyl chloroformate (0.23 ml, 1.9 mmol) and triethylamine (0.26 ml, 1.9 mmol). This intermediate (125 mg, 0.300 mmol) was dissolved in N,N-dimethylformamide (2.5 ml)-triethylamine (0.5 ml); ethylamine hydrochloride (122 mg, 1.50 mmol) was added thereto; and the reaction mixture was stirred at room temperature overnight, and then stirred at 80 °C for 1.5 hours. Ethylamine hydrochloride (122 mg, 1.50 mmol) was added thereto; the reaction mixture was stirred at 80 °C for 2 hours; ethylamine hydrochloride (122 mg, 1.50 mmol) and triethylamine (0.5 ml) were further added thereto; and the reaction mixture was stirred at 80 °C for 0.5 hours, and then stirred at room temperature for 2 days. The reaction mixture was partitioned between ethyl acetate (100 ml) and water (50 ml); and the organic layer was

washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Fuji Silysia BW-300, hexane: ethyl acetate = 3: 2, then ethyl acetate); the obtained crystals were suspended in diethyl ether (10 ml) -hexane (50 ml), filtered off, and dried to yield the title compound (41 mg, 0.11 mmol) as colorless crystals.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.04 (3H, t, J=7.2 Hz), 2.83 (3H, d, J=4.4 Hz), 3.15 (2H, m), 3.19 (3H, s), 6.51 (1H, dd, J=2.4, 6.6 Hz), 6.67 (1H, d, J=4.0 Hz), 6.77 (1H, d, J=2.4 Hz), 7.07 (1H, dd, J=2.4, 8.8 Hz), 7.39 (1H, d, J=2.4 Hz), 7.87 (1H, d, J=4.0 Hz), 8.15 (1H, d, J=6.0 Hz), 8.16 (1H, m), 8.29 (1H, d, J=8.8 Hz), 9.27 (1H, m).

ESI-MS: 368.13 (M+H).

Example 187

6-(2-(3-Ethylureido)pyridin-4-yloxy)indole-1-carboxylic acid methylamide

[0452] Similarly to Example 5, the title compound (54 mg, 0.15 mmol, 80%) was obtained as colorless crystals from phenyl (4-(1-methylcarbamoyl-1H-indol-6-yloxy)pyridin-2-yl)-N-(phenoxycarbonyl)carbamate (100 mg, 0.19 mmol), ethylamine hydrochloride (78 mg, 0.96 mmol) and triethylamine (0.5 ml).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.03 (3H, t, J=7.2 Hz), 2.79 (3H, d, J=4.4 Hz), 3.07-3.14 (2H, m), 6.52 (1H, dd, J=2.4, 5.8 Hz), 6.71 (1H, d, J=3.6 Hz), 6.88 (1H, d, J=2.4 Hz), 6.99 (1H, dd, J=2.4, 8.4 Hz), 7.65 (1H, d, J=8.4 Hz), 7.84 (1H, d, J=3.6 Hz), 7.96 (2H, m), 8.04 (1H, d, J=5.8 Hz), 8.16 (1H, m), 9.02 (1H, s).
ESI-MS: 354.15 (M+H), 376.16 (M+Na).

[0453] The starting materials were synthesized as follows.

Production example 187-1

4-(1H-Indol-6-yloxy)pyridin-2-ylamine

Sodium hydride (1.04 g, 26.0 mmol, 60% in oil) was suspended in dimethyl sulfoxide (2.5 ml); 6-hydroxyindole (3.46 g, 26.0 mmol) and 2-amino-4-chloropyridine (2.57 g, 20.0 mmol, WO 02/332872) were subsequently added thereto at room temperature under nitrogen stream; and the reaction mixture was stirred at 160 °C for 8.5 hours. After cooled down to room temperature, the reaction mixture was partitioned between ethyl acetate (150 ml) and a solvent mixture of aqueous ammonia: water = 1: 1 (50 ml); the organic layer was washed with a solvent mixture of aqueous ammonia: water = 1: 1, and with brine, dried over anhydrous sodium sulfate, and concentrated. The residue was purified by silica gel column

chromatography (Fuji Silysia BW-300, ethyl acetate, or ethyl acetate: methanol = 93: 7); and the obtained crystals were suspended in diethyl ether, filtered off, washed with diethyl ether, and dried to yield the title compound (477 mg, 2.12 mmol, 10.6%) as pale yellow crystals.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 5.76 (1H, s), 5.82 (2H, brs), 6.13 (1H, d, J=6.0 Hz), 6.44 (1H, s), 6.75 (1H, d, J=8.4 Hz), 7.10 (1H, s), 7.34 (1H, s), 7.56 (1H, d, J=8.4 Hz), 7.75 (1H, d, J=6.0 Hz), 11.12 (1H, brs).

Production example 187-2

6-(2-Aminopyridin-4-yloxy)indole-1-carboxylic acid methylamide

[0454] Similarly to Production example 5-1, the title compound (315 mg, 1.12 mmol, 87.9%) was obtained as colorless crystals from 4-(1H-indol-6-yloxy)pyridin-2-ylamine (285 mg, 1.27 mmol), sodium hydride (63 mg, 1.58 mmol, 60% in oil) and phenyl N-methylcarbamate (239 mg, 1.58 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.80 (3H, d, J=4.4 Hz), 5.77 (1H, d, J=2.0 Hz), 5.85 (2H, m), 6.14 (1H, dd, J=2.0, 5.6 Hz), 6.69 (1H, d, J=3.6 Hz), 6.96 (1H, dd, J=2.0, 8.4 Hz), 7.63 (1H, d, J=8.4 Hz), 7.77 (1H, d, J=5.6 Hz), 7.81 (1H, d, J=3.6 Hz), 7.94 (1H, d, J=2.0 Hz), 8.13 (1H, d, J=4.4 Hz).

Production example 187-3Phenyl (4-(1-methylcarbamoyl-1H-indol-6-yloxy)pyridin-2-yl)-N-(phenoxycarbonyl)carbamate

5 [0455] Similarly to Production example 5-2, the title compound (404 mg, 0.77 mmol, 69%) was obtained as pale pink crystals from 6-(2-aminopyridin-4-yloxy)indole-1-carboxylic acid methylamide (315 mg, 1.12 mmol), triethylamine (0.51 ml, 3.7 mmol), and phenyl
10 chloroformate (0.42 ml, 3.4 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.78 (3H, d, J=4.4 Hz), 6.74 (1H, d, J=3.6 Hz), 7.02 (1H, dd, J=2.4, 5.6 Hz), 7.05 (1H, dd, J=2.4, 8.4 Hz), 7.16 (4H, d, J=7.8 Hz), 7.29 (2H, t, J=7.8 Hz), 7.42 (4H, t, J=7.8 Hz),
15 7.52 (1H, m), 7.69 (1H, d, J=8.4 Hz), 7.86 (1H, d, J=3.6 Hz), 8.04 (1H, d, J=2.4 Hz), 8.15 (1H, m), 8.44 (1H, d, J=5.6 Hz).

Example 188

20 6-(2-(3,3-Diethylureido)pyridin-4-yloxy)indole-1-carboxylic acid methylamide

[0456] Similarly to Example 5, the title compound (55 mg, 0.14 mmol, 76%) was obtained as colorless crystals from phenyl (4-(1-methylcarbamoyl-1H-indol-6-yloxy)pyridin-2-yl)-N-(phenoxycarbonyl)carbamate (100
25 mg, 0.19 mmol) and diethylamine (0.10 ml, 0.96 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.01 (6H, t, J=7.2 Hz), 2.79 (3H, d, J=4.4 Hz), 3.26-3.32 (4H, m), 6.56 (1H, dd, J=2.0, 5.6 Hz), 6.71 (1H, d, J=3.6 Hz), 6.99 (1H, dd, J=2.0, 8.8 Hz), 7.42 (1H, d, J=2.0 Hz), 7.65 (1H, d, J=8.8 Hz), 7.84 (1H, d, J=3.6 Hz), 7.96 (1H, d, J=2.0 Hz), 8.08 (1H, d, J=5.6 Hz), 8.15 (1H, m), 8.63 (1H, s).

ESI-MS: 382.21 (M+H).

10 Example 189

6-(2-(3-(2-Diethylaminoethyl)ureido)pyridin-4-yloxy)indole-1-carboxylic acid methylamide

[0457] Similarly to Example 5, the title compound (51 mg, 0.12 mmol, 63%) was obtained as pale yellow crystals from phenyl (4-(1-methylcarbamoyl-1H-indol-6-yloxy)pyridin-2-yl)-N-(phenoxycarbonyl)carbamate (100 mg, 0.19 mmol) and 2-diethylaminoethylamine (0.14 ml, 0.96 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.93 (6H, t, J=7.6 Hz), 2.41-2.49 (6H, m), 2.79 (3H, d, J=4.0 Hz), 3.14 (2H, m), 6.51 (1H, dd, J=2.4, 6.0 Hz), 6.71 (1H, d, J=3.6 Hz), 6.84 (1H, d, J=2.4 Hz), 6.99 (1H, dd, J=2.4, 8.2 Hz), 7.65 (1H, d, J=8.2 Hz), 7.84 (1H, d, J=3.6 Hz), 7.96 (1H, d, J=2.4 Hz), 8.02 (1H, d, J=6.0 Hz), 8.16 (2H, m), 9.13 (1H, s).

ESI-MS: 425.29 (M+H).

Example 190

6-(2-(((4-(Pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)indole-1-carboxylic acid methylamide

[0458] Similarly to Example 5, the title compound (72 mg, 0.16 mmol, 82%) was obtained as colorless crystals from phenyl (4-(1-methylcarbamoyl-1H-indol-6-yloxy)pyridin-2-yl)-N-(phenoxycarbonyl)carbamate (100 mg, 0.19 mmol) and 4-(pyrrolidin-1-yl)piperidine (148 mg, 0.96 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.19-1.31 (2H, m), 1.63 (4H, m), 1.76 (2H, m), 2.09 (1H, m), 2.44 (4H, m), 2.79 (3H, d, J=4.0 Hz), 2.82 (2H, m), 3.92 (2H, m), 6.55 (1H, dd, J=2.4, 5.6 Hz), 6.71 (1H, d, J=3.8 Hz), 6.98 (1H, dd, J=2.4, 8.8 Hz), 7.32 (1H, d, J=2.4 Hz), 7.65 (1H, d, J=8.8 Hz), 7.84 (1H, d, J=3.8 Hz), 7.96 (1H, d, J=2.4 Hz), 8.08 (1H, d, J=5.6 Hz), 8.15 (1H, m), 9.12 (1H, s).

ESI-MS: 436.32 (M+H).

Example 191

6-(6-(3-Ethylureido)pyrimidin-4-yloxy)indole-1-carboxylic acid methylamide

[0459] Similarly to Production example 5-2, an intermediate, phenyl (4-(1-methylcarbamoyl-1H-indol-6-

yloxy)pyrimidin-6-yl)-N-(phenoxycarbonyl)carbamate, was obtained as pale yellow crystals (597 mg) from 6-(6-aminopyrimidin-4-yloxy)indole-1-carboxylic acid methylamide (245 mg, 0.865 mmol), triethylamine (0.40 ml, 2.9 mmol), and phenyl chloroformate (0.33 ml, 2.6 mmol). Similarly to Example 5, the title compound (43 mg, 0.12 mmol) was obtained as colorless crystals from this intermediate (143 mg), ethylamine hydrochloride (88 mg, 1.1 mmol), and triethylamine (0.5 ml).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.03 (3H, t, J=6.8 Hz), 2.80 (3H, s), 3.10 (2H, m), 6.71 (1H, s), 6.99 (1H, d, J=8.4 Hz), 7.01 (1H, s), 7.04 (1H, m), 7.62 (1H, d, J=8.4 Hz), 7.84 (1H, s), 7.98 (1H, s), 8.16 (1H, m), 8.36 (1H, s), 9.45 (1H, s).

ESI-MS: 355.27 (M+H), 377.26 (M+Na).

[0460]The starting material was synthesized as follows.

Production example 191-1

6-(1H-Indol-6-yloxy)pyrimidin-4-ylamine

Sodium hydride (200 mg, 5.00 mmol) was suspended in dimethyl sulfoxide (8 ml); while stirring at room temperature, 6-hydroxyindole (666 mg, 5.00 mmol) and 6-amino-4-chloropyrimidine (518 mg, 4.00 mmol) were added thereto one by one; and the reaction mixture was stirred at 60 °C for 2 hours, at 80 °C for 1 hour, and at 100 °C for 1.5 hours. After cooled down to room

temperature, the reaction mixture was partitioned between ethyl acetate and water; and the organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated. The residue was purified by silica gel column chromatography (Fuji Silysia BW-300, hexane: ethyl acetate = 1: 1, ethyl acetate, then ethyl acetate: methanol = 98: 2); the obtained crystals were suspended in ethyl acetate (50 ml) and stirred at room temperature overnight, filtered off, washed with diethyl ether, and dried to yield the title compound (322 mg, 1.42 mmol, 35.6%) as pale yellow crystals.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 5.56 (1H, s), 6.44 (1H, s), 6.72 (2H, s), 6.76 (1H, d, J=8.4 Hz), 7.12 (1H, s), 7.34 (1H, s), 7.55 (1H, d, J=8.4 Hz), 8.05 (1H, s), 11.13 (1H, brs).

Production example 191-2

6-(6-Aminopyrimidin-4-yloxy)indole-1-carboxylic acid methylamide

[0461] Similarly to Production example 5-1, the title compound (245 mg, 0.865 mmol, 61.3%) was obtained as colorless crystals from 6-(1H-indol-6-yloxy)pyrimidin-4-ylamine (320 mg, 1.41 mmol), sodium hydride (68 mg, 1.7 mmol, 60% in oil), and phenyl N-methylcarbamate (257 mg, 1.70 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.80 (3H, d, J=4.4

Hz), 5.64 (1H, s), 6.69 (1H, d, J=3.6 Hz), 6.77 (2H, s),
6.96 (1H, dd, J=2.0, 8.4 Hz), 7.61 (1H, d, J=8.4 Hz),
7.81 (1H, d, J=3.6 Hz), 7.94 (1H, d, J=2.0 Hz), 8.05
(1H, s), 8.12 (1H, m).

5

Example 192

6-(6-(3,3-Diethylureido)pyrimidin-4-yloxy)indole-1-
carboxylic acid methylamide

[0462] Similarly to Example 5, the title compound (63
10 mg, 0.16 mmol) was obtained as milky white crystals
from the intermediate obtained in Example 191 (149 mg)
and diethylamine (0.11 ml, 1.1 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.03 (6H, t, J=7.2
Hz), 2.80 (3H, d, J=4.4 Hz), 3.33 (4H, q, J=7.2 Hz),
15 6.71 (1H, d, J=3.8 Hz), 7.00 (1H, dd, J=2.0, 8.4 Hz),
7.31 (1H, s), 7.62 (1H, d, J=8.4 Hz), 7.83 (1H, d,
J=3.8 Hz), 7.98 (1H, d, J=2.0 Hz), 8.15 (1H, m), 8.38
(1H, s), 9.31 (1H, s).

ESI-MS: 383.23 (M+H), 405.26 (M+Na).

20

Example 193

6-(6-(3-(2-Diethylaminoethyl)ureido)pyrimidin-4-
yloxy)indole-1-carboxylic acid methylamide

[0463] Similarly to Example 5, the title compound (63
25 mg, 0.15 mmol) was obtained as grayish white crystals
from the intermediate obtained in Example 191 (164 mg)

and 2-diethylaminoethylamine (0.15 ml, 1.1 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.93 (6H, t, J=7.0 Hz), 2.44 (6H, m), 2.80 (3H, d, J=4.0 Hz), 3.13 (2H, m), 6.70 (1H, d, J=3.6 Hz), 6.90 (2H, m), 7.43 (1H, brs), 7.62 (1H, d, J=8.4 Hz), 7.83 (1H, d, J=3.6 Hz), 7.98 (1H, d, J=1.6 Hz), 8.15 (1H, m), 8.34 (1H, s), 9.63 (1H, s).

ESI-MS: 426.31 (M+H).

10 Example 194

6-(6-(((4-Pyrrolidin-1-yl)piperidin-1-ylcarbonyl)amino)pyrimidin-4-yloxy)indole-1-carboxylic acid methylamide

15 [0464] Similarly to Example 5, the title compound (59 mg, 0.13 mmol) was obtained as colorless crystals from the intermediate obtained in Example 191 (141 mg) and 4-(pyrrolidin-1-yl)piperidine (167 mg, 1.08 mmol).

20 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.22-1.34 (2H, m), 1.64 (4H, m), 1.78 (2H, m), 2.12 (1H, m), 2.44 (4H, m), 2.80 (3H, d, J=4.0 Hz), 2.88 (2H, m), 3.93 (2H, m), 6.70 (1H, d, J=3.6 Hz), 6.99 (1H, dd, J=2.0, 8.4 Hz), 7.20 (1H, s), 7.62 (1H, d, J=8.4 Hz), 7.83 (1H, d, J=3.6 Hz), 7.97 (1H, d, J=2.0 Hz), 8.15 (1H, m), 8.38 (1H, s), 9.73 (1H, s).

25 ESI-MS: 464.36 (M+H).

Example 1954-(6-(3-Ethylureido)pyrimidin-4-yloxy)indole-1-carboxylic acid methylamide

[0465] Similarly to Production example 5-2, an
5 intermediate (a mixture of phenyl (4-(1-methylcarbamoyl-1H-indol-4-yloxy)pyrimidin-6-yl)-N-(phenoxycarbonyl)carbamate and phenyl (4-(1-methylcarbamoyl-1H-indol-4-yloxy)pyrimidin-6-yl)carbamate, 379 mg) was obtained as pale yellow
10 crystals from 4-(6-Aminopyrimidin-4-yloxy)indole-1-carboxylic acid methylamide (245 mg, 0.865 mmol), triethylamine (0.40 ml, 2.9 mmol) and phenyl chloroformate (0.33 ml, 2.6 mmol). Similarly to
15 Example 5, the title compound (41 mg, 0.12 mmol) was obtained as a colorless crystal from this intermediate (94 mg), ethylamine hydrochloride (78 mg, 0.96 mmol), and triethylamine (0.5 ml).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.03 (3H, t, J=7.4 Hz), 2.82 (3H, d, J=4.0 Hz), 3.12 (2H, m), 6.40 (1H, d, J=3.8 Hz), 6.98 (1H, d, J=8.0 Hz), 7.05 (1H, s), 7.28 (1H, t, J=8.0 Hz), 7.31 (1H, m), 7.76 (1H, d, J=3.8 Hz), 8.14 (1H, d, J=8.0 Hz), 8.17 (1H, m), 8.33 (1H, m), 9.48 (1H, s).

ESI-MS: 355.20 (M+H), 377.25 (M+Na).

25

[0466] The starting materials were synthesized as

follows.

Production example 195-1

6-(1H-Indol-4-yloxy)pyrimidin-4-ylamine

The title compound (568 mg, 2.51 mmol, 41.8%)
5 was obtained as grayish white crystals by performing a
reaction similar to that in Production example 191-1
using 6-amino-4-chloropyrimidine (777 mg, 6.00 mmol),
4-hydroxyindole (999 mg, 7.50 mmol) and sodium hydride
(300 mg, 7.50 mmol) at 100 °C for 6 hours.

10 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 5.56 (1H, s), 6.13
(1H, m), 6.70 (2H, brs), 6.74 (1H, d, J=8.0 Hz), 7.09
(1H, t, J=8.0 Hz), 7.29 (2H, m), 8.05 (1H, s), 11.28
(1H, s).

15 Production example 195-2

4-(6-Aminopyrimidin-4-yloxy)indole-1-carboxylic acid
methanamide

[0467] Similarly to Production example 5-1, the title
compound (279 mg, 0.985 mmol, 74.0%) was obtained as
20 colorless crystals from 6-(1H-indol-4-yloxy)pyrimidin-
4-ylamine (300 mg, 1.33 mmol), sodium hydride (83 mg,
2.1 mmol, 60% in oil), and phenyl N-methylcarbamate
(314 mg, 2.07 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.82 (3H, d, J=4.4
25 Hz), 5.64 (1H, s), 6.39 (1H, d, J=3.6 Hz), 6.77 (2H,
brs), 6.94 (1H, d, J=8.0 Hz), 7.27 (1H, t, J=8.0 Hz),

7.75 (1H, d, J=3.6 Hz), 8.04 (1H, s), 8.12 (1H, d, J=8.0 Hz), 8.15 (1H, m).

Example 196

5 4-(6-(3,3-Diethylureido)pyrimidin-4-yloxy)indole-1-carboxylic acid methylamide

[0468] Similarly to Example 5, the title compound (54 mg, 0.14 mmol) was obtained as colorless crystals from the intermediate obtained in Example 195 (94 mg) and
10 diethylamine (0.10 ml, 0.96 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.04 (6H, t, J=6.8 Hz), 2.82 (3H, d, J=4.0 Hz), 3.34 (4H, q, J=6.8 Hz), 6.41 (1H, d, J=3.8 Hz), 6.98 (1H, d, J=8.0 Hz), 7.28 (1H, t, J=8.0 Hz), 7.36 (1H, s), 7.76 (1H, d, J=3.8 Hz),
15 8.14 (1H, d, J=8.0 Hz), 8.17 (1H, m), 8.35 (1H, s), 9.34 (1H, s).

ESI-MS: 383.31 (M+H), 405.22 (M+Na).

Example 197

20 4-(6-(3-(2-Diethylaminoethyl)ureido)pyrimidin-4-yloxy)indole-1-carboxylic acid methylamide

[0469] Similarly to Example 5, the title compound (49 mg, 0.12 mmol) was obtained as colorless crystals from the intermediate obtained in Example 195 (94 mg) and 2-
25 diethylaminoethylamine (0.14 ml, 0.96 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.93 (6H, t, J=7.0

Hz), 2.45 (6H, m), 2.82 (3H, d, J=4.0 Hz), 3.14 (2H, m),
6.40 (1H, d, J=3.4 Hz), 6.98 (1H, d, J=8.0 Hz), 7.04
(1H, s), 7.28 (1H, t, J=8.0 Hz), 7.45 (1H, m), 7.76 (1H,
d, J=3.4 Hz), 8.14 (1H, d, J=8.0 Hz), 8.17 (1H, m),
5 8.32 (1H, s), 8.65 (1H, brs).

ESI-MS: 426.27 (M+H).

Example 198

10 4-(6-(((4-(Pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino)pyrimidin-4-yloxy)indole-1-carboxylic acid methylamide

[0470] Similarly to Example 5, the title compound (57
mg, 0.12 mmol) was obtained as colorless crystals from
the intermediate obtained in Example 195 (94 mg) and 4-
15 (pyrrolidin-1-yl)piperidine (148 mg, 0.96 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.22-1.35 (2H, m),
1.64 (4H, m), 1.78 (2H, m), 2.13 (1H, m), 2.45 (4H, m),
2.82 (3H, d, J=3.2 Hz), 2.89 (2H, m), 3.94 (2H, m),
6.40 (1H, m), 6.98 (1H, d, J=8.0 Hz), 7.26 (1H, s),
20 7.28 (1H, t, J=8.0 Hz), 7.76 (1H, m), 8.13 (1H, d,
J=8.0 Hz), 8.16 (1H, m), 8.35 (1H, s), 9.35 (1H, s).

ESI-MS: 464.35 (M+H).

Example 199

25 5-(2-(3-(3-Diethylaminopropyl)ureido)pyridin-4-ylamino)indole-1-carboxylic acid methylamide

[0471] 1-(4-Chloropyridin-2-yl)-3-(3-diethylaminopropyl)urea (30 mg, 0.11 mmol) was dissolved in ethoxyethanol (1.1 ml); pyridine hydrochloride (24 mg, 0.22 mmol) and 5-aminoindole-1-carboxylic acid methylamide (22 mg, 0.12 mmol, Production example 218-2) was added thereto; and the reaction mixture was stirred at 130 °C for 2 hours. After cooled down to room temperature, the reaction mixture was partitioned between a saturated aqueous solution of sodium hydrogencarbonate and ethyl acetate; and the organic layer was washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Fuji Silysia NH, hexane: ethyl acetate = 1: 3, ethyl acetate, ethyl acetate: methanol = 93: 7 in this order) to yield the title compound (8 mg, 0.018 mmol, 17%) as pale yellow powder.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.94 (6H, t, J=6.8 Hz), 1.54 (2H, m), 2.37-2.46 (6H, m), 2.83 (3H, d, J=3.6 Hz), 3.16 (2H, m), 6.42 (1H, d, J=5.8 Hz), 6.63 (1H, d, J=3.2 Hz), 6.73 (1H, s), 7.07 (1H, d, J=8.8 Hz), 7.37 (1H, s), 7.76 (1H, d, J=5.8 Hz), 7.80 (1H, m), 8.08 (1H, m), 8.19 (1H, d, J=8.8 Hz), 8.66 (1H, s), 8.81 (1H, m), 8.86 (1H, s).

ESI-MS : 438.36 (M+H).

[0472] The starting materials were synthesized as follows.

Production example 199-1

Phenyl (4-chloropyridin-2-yl)-N-(phenoxycarbonyl)carbamate

2-Amino-4-chloropyridine (5.00 g, 38.9 mmol, WO 02/32872) was dissolved in tetrahydrofuran (200 ml); and triethylamine (17.9 ml, 128 mmol) was added thereto. While stirring with a waterbath, phenyl chloroformate (14.6 ml, 117 mmol) was added thereto dropwise; the reaction mixture was stirred at room temperature for 1.5 hours. The reaction mixture was partitioned between water and ethyl acetate; the organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was filtered by silica gel; the crystals obtained after the concentration were suspended in diethyl ether, filtered off, washed with diethyl ether, and dried to yield the title compound (3.77 g, 10.2 mmol, 26.3%) as pale yellow crystals. The mother liquor was concentrated under reduced pressure, which was then treated by the similar methods to yield the title compound (3.98 g, 10.5 mmol, 27.1%) as pale yellow crystals (secondary crystals).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 7.20 (4H, d, J=7.6 Hz), 7.30 (2H, t, J=7.6 Hz), 7.44 (4H, t, J=7.6 Hz),

7.68 (1H, dd, J=1.6, 5.2 Hz), 8.21 (1H, d, J=1.6 Hz),
8.60 (1H, d, J=5.2 Hz).

Production example 199-2

5 1-(4-Chloropyridin-2-yl)-3-(3-diethylaminopropyl)urea
 [0473] Phenyl (4-chloropyridin-2-yl)-N-
 (phenoxy-carbonyl)carbamate (738 mg, 2.00 mmol) was
 dissolved in N,N-dimethylformamide (8.0 ml); N,N-
 diethyl-1,3-diaminopropane (1.57 ml, 10.0 mmol) was
 added thereto; and the reaction mixture was stirred at
 room temperature for 1 hour. The reaction mixture was
 partitioned between water and ethyl acetate; and the
 organic layer was washed with brine, dried over
 anhydrous sodium sulfate, and concentrated under
 reduced pressure. The residue was purified by silica
 gel column chromatography (Fuji Silysia NH, hexane-
 ethyl acetate-methanol system) to yield the title
 compound (309 mg, 1.09 mmol, 54.3%) as a pale brown oil.
 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.92 (6H, t, J=7.0
 Hz), 1.54 (2H, m), 2.35-2.44 (6H, m), 3.16 (2H, m),
 7.02 (1H, d, J= 5.6 Hz), 7.54 (1H, s), 7.73 (1H, brs),
 8.13 (1H, d, J=5.6 Hz), 9.31 (1H, m).

Example 200

25 5-(N-(2-(3-(3-Diethylaminopropyl)ureido)pyridin-4-yl)-
 N-methylamino)indole-1-carboxylic acid methylamide

[0474] Similarly to Example 199, the title compound (6 mg, 0.013 mmol, 12%) was obtained as pale yellow powder from 5-(N-methylamino)indol-1-carboxylic acid methylamide (22 mg, 0.11 mmol), 1-(4-chloropyridin-2-yl)-3-(3-diethylaminopropyl)urea (30 mg, 0.11 mmol, Production example 199-2) and pyridine hydrochloride (25 mg, 0.22 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.91 (6H, t, J=7.0 Hz), 1.51 (2H, m), 2.34-2.43 (6H, m), 2.83 (3H, d, J=4.0 Hz), 3.13 (2H, m), 3.23 (3H, s), 6.11 (1H, d, J=6.0 Hz), 6.40 (1H, s), 6.70 (1H, d, J=3.6 Hz), 7.10 (1H, d, J=8.6 Hz), 7.44 (1H, s), 7.69 (1H, d, J=6.0 Hz), 7.84 (1H, d, J=3.6 Hz), 8.14 (1H, m), 8.27 (1H, d, J=8.6 Hz), 8.76 (1H, s), 8.78 (1H, brs).

ESI-MS: 452.38 (M+H).

[0475]The starting material was synthesized as follows.

Production example 200-1

5-(N-Methylamino)indole-1-carboxylic acid methylamide

5-Aminoindole-1-carboxylic acid methylamide (22 mg, 0.11 mmol, Production example 218-2) was dissolved in methanol (5.5 ml); and benzotriazol-1-ylmethanol (434 mg, 2.91 mmol) was added thereto. Because crystals were precipitated immediately, methanol (5.5 ml) was added to dissolve the precipitation, and the reaction mixture was stirred at room temperature for

1.25 hours. Then, the reaction mixture was heated and stirred at 60 °C for an hour. After cooled to room temperature, precipitated crystals were filtered off, washed by methanol, and dried to yield colorless crystals (421 mg). The crystals were dissolved in a solvent mixture of N,N-dimethylformamide (4.2 ml) and methanol (21 ml); sodium borohydride (99 mg, 2.63 mmol) was added while stirring at room temperature; and the reaction mixture was stirred for 1.5 hours. Sodium borohydride (99 mg, 2.63 mmol) was further added thereto; and the reaction mixture was stirred at room temperature for 12 hours. A similar reaction was performed using the residue obtained by the concentration of the mother liquor the crystals were previously given from under reduced pressure, and sodium borohydride (342 mg, 9.02 mmol). Both reaction mixtures mentioned above were partitioned between a saturated aqueous solution of sodium hydrogencarbonate and ethyl acetate; both organic layers are combined, washed with a saturated aqueous solution of sodium hydrogencarbonate, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Fuji Silysia BW-300, hexane-ethyl acetate-methanol system). The obtained crystals were suspended in ethyl acetate, filtered off, washed with ethyl acetate, and dried to

yield the title compound (255 mg, 1.25 mmol, 43.1%) was obtained as pale pink crystals.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.66 (3H, s), 2.78 (3H, d, J=4.4 Hz), 5.32 (1H, brs), 6.42 (1H, d, J=3.6 Hz), 6.56 (1H, d, 2.4 Hz), 6.57 (1H, dd, J=2.4, 9.0 Hz), 7.61 (1H, d, J=3.6 Hz), 7.84 (1H, d, J=4.4 Hz), 7.93 (1H, d, J=9.0 Hz).

Example 201

10 5-(2-(3,3-Diethylureido)pyridin-4-ylamino)indole-1-carboxylic acid phenylamide

[0476] 5-(2-Aminopyridin-4-ylamino)indole-1-carboxylic acid phenylamide (69 mg, 0.20 mmol) was dissolved in tetrahydrofuran (14 ml); triethylamine (0.055 ml, 0.40 mmol) was added thereto; and phenyl chloroformate (0.038 ml, 0.30 mmol) was added thereto while cooling with ice and stirring. A portion of 7.0 ml of the reaction mixture was transferred to another vessel and concentrated under reduced pressure. After the residue was dissolved in N,N-dimethylformamide (1.0 ml), the similar reaction to Example 27 was performed by use of diethylamine (0.031 ml, 0.30 mmol). The obtained crude product was purified by TLC plate (Fuji Silysia NH, developing solvent: ethyl acetate) to yield the title compound (2.0 mg, 0.005 mmol) as pale yellow crystals.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.07 (6H, t, J=7.0

Hz), 3.33 (4H, m), 6.52 (1H, m), 6.72 (1H, m), 7.14 (2H, m), 7.40 (3H, m), 7.52 (1H, s), 7.66 (2H, d, J=7.6 Hz), 7.83 (1H, d, J=6.4 Hz), 8.04 (1H, d, J=2.8 Hz), 8.18 (2H, m), 8.65 (1H, s), 10.03 (1H, s).

5 ESI-MS: 443.28 (M+H).

[0477] The starting materials were synthesized as follows.

Production example 201-1

10 5-Nitroindole-1-carboxilic acid phenylamide

Sodium hydride (802 mg, 20.0 mmol, 60% in oil) was suspended in N,N-dimethylformamide (40 ml); 5-nitroindole (2.50 g, 15.4 mmol) was added thereto; and the reaction mixture was stirred at room temperature for 30 minutes. Phenyl isocyanate (2.01 ml, 1.23 mmol) was added thereto, and the reaction mixture was stirred at room temperature for 1.5 hours. Water (80 ml) was added to the reaction mixture; the reaction mixture was stirred at room temperature for 30 minutes; and the precipitated crystals were filtered off, washed by water and diethyl ether one by one, and dried by means of suction to yield the title compound (3.53 g, 12.3 mmol, 79.8%) as pale yellow crystals.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 7.00 (1H, d, J=3.6 Hz), 7.16 (1H, t, J=8.0 Hz), 7.40 (2H, t, J=8.0 Hz), 7.65 (2H, d, J=8.0 Hz), 8.17 (1H, dd, J=2.4, 9.2 Hz),

8.25 (1H, d, J=3.6 Hz), 8.36 (1H, d, J=9.2 Hz), 8.62 (1H, d, J=2.4 Hz), 10.30 (1H, s).

Production example 201-2

5 5-Aminoindole-1-carboxylic acid phenylamide

[0478] 5-Nitroindole-1-carboxylic acid phenylamide (3.53 g, 12.3 mmol) was dissolved in ethanol (250 ml); water (50 ml), electrolytic iron powder (2.75 g, 49.2 mmol), ammonium chloride (5.26 g, 98.4 mmol) were added thereto; and the reaction mixture was heated and stirred at 80 °C for 2 hours. After cooling to room temperature, the reaction mixture was filtered off; insoluble portions were washed with ethyl acetate; and the filtrate was concentrated under reduced pressure. The residue was partitioned between water and a solvent mixture of ethyl acetate and tetrahydrofuran; and the organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was solidified by diethyl ether; the crystals were suspended in diethyl ether, filtered off, washed with diethyl ether, and dried to yield the title compound as pale red powder (681 mg, 2.71 mmol, 22.0%). The mother liquor was concentrated under reduced pressure, which was then treated by the similar methods to yield the title compound (590 mg, 2.35 mmol, 19.1%) as pale red powder

(secondary crystals).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 4.80 (2H, s), 6.48 (1H, d, J=3.4 Hz), 6.59 (1H, dd, J=2.4, 8.8 Hz), 6.71 (1H, d, J=2.4 Hz), 7.09 (1H, t, J=7.6 Hz), 7.34 (2H, t, J=7.6 Hz), 7.61 (2H, d, J=7.6 Hz), 7.84 (1H, d, J=3.4 Hz), 7.88 (1H, d, J=8.8 Hz), 9.79 (1H, s).

Production example 201-3

5-(2-Aminopyridin-4-ylamino)indole-1-carboxylic acid phenylamide

[0479] 2-Amino-4-chloropyridine (500 mg, 0.446 mmol) was dissolved in N-methylpyrrolidone (5.0 ml); pyridine hydrochloride (750 mg) and 5-aminoindole-1-carboxylic acid phenylamide (408 mg, 1.62 mmol) was added thereto; the reaction mixture was stirred at 100 °C for 6.5 hours. After cooling to room temperature, the reaction mixture was partitioned between saturated aqueous solution of sodium hydrogencarbonate and ethyl acetate; the organic layer was washed with brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Fuji Silysia NH, hexane-ethyl acetate-methanol system). The obtained pale yellow oil was solidified with diethyl ether; and the crystals were suspended in diethyl ether, filtered off, washed with diethyl ether, and dried to yield the title

compound (188 mg, 0.464 mmol, 35.7%) as pale yellow crystals.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 5.45 (2H, m), 5.99 (1H, d, J=2.0 Hz), 6.10 (1H, dd, J=2.0, 6.0 Hz), 6.69 (1H, d, J=3.6 Hz), 7.07 (1H, dd, J=2.0, 8.6 Hz), 7.12 (1H, t, J=7.6 Hz), 7.37 (3H, m), 7.56 (1H, d, J=6.0 Hz), 7.63 (2H, d, J=7.6 Hz), 8.00 (1H, d, J=3.6 Hz), 8.14 (1H, d, J=8.6 Hz), 8.26 (1H, s), 9.98 (1H, s).

Example 202

5-(2-(3-(3-Diethylaminopropyl)ureido)pyridin-4-ylamino)indole-1-carboxylic acid phenylamide

[0480] 5-(2-Aminopyridin-4-ylamino)indole-1-carboxylic acid phenylamide (69 mg, 0.20 mmol, Production example 201-3) was dissolved in tetrahydrofuran (14 ml); triethylamine (0.055 ml, 0.40 mmol) was added thereto; and phenyl chloroformate (0.038 ml, 0.30 mmol) was added while stirring and cooled by ice. A portion of 7.0 ml of this reaction mixture was transferred to another vessel; and the remaining portion of the reaction mixture was concentrated under reduced pressure. The residue was dissolved in N,N-dimethylformamide (1.0 ml); the similar reaction to Example 201 was performed using N,N-diethyl-1,3-diaminopropane (0.047 ml, 0.30 mmol); the crude product obtained was purified by a TLC plate (Fuji Silysia NH,

developing solvent: ethyl acetate/ethanol = 10/1); and the obtained crystals were suspended in ethyl acetate, filtered off, and dried to yield the title compound (3 mg, 0.006 mmol) as colorless crystals.

5 ¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.93 (6H, t, J=7.0 Hz), 1.53 (2H, m), 2.42 (6H, m), 3.18 (2H, m), 6.43 (1H, d, J=5.6 Hz), 6.70 (1H, s), 6.75 (1H, s), 7.12 (2H, m), 7.38 (3H, m), 7.64 (2H, d, J=8.0 Hz), 7.76 (1H, d, J=5.6 Hz), 8.03 (1H, s), 8.16 (1H, d, J=9.2 Hz), 8.70
10 (1H, s), 8.78 (1H, m), 8.86 (1H, s), 10.01 (1H, s).
ESI-MS: 500.54 (M+H).

Example 203

15 5-(5-Cyano-2-(3-(2-diethylaminoethyl)ureido)pyridin-4-ylamino)indole-1-carboxylic acid phenylamide

[0481] A reaction similar to Production example 5-2 was performed using 5-(2-amino-5-cyanopyridin-4-ylamino)indole-1-carboxylic acid phenylamide (60 mg, 0.16 mmol), triethylamine (0.056 ml, 0.41mmol), and
20 phenyl chloroformate (0.082 ml, 0.66 mmol); the solvent was concentrated under reduced pressure. Similarly to Example 5, the title compound (63 mg, 0.12 mmol, 76%) was obtained as pale yellow crystals from the residue obtained above, and 2-diethylaminoethylamine (0.115 ml,
25 0.81 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.92 (6H, t, J=7.0

Hz), 2.38-2.46 (6H, m), 3.09 (2H, m), 6.75 (1H, d, J=3.8 Hz), 7.03 (1H, brs), 7.13 (1H, dd, J=6.8, 7.6 Hz), 7.18 (1H, dd, J=2.0, 8.8 Hz), 7.38 (2H, t, J=7.6 Hz), 7.48 (1H, d, J=2.0 Hz), 7.65 (3H, m), 8.07 (1H, d, J=3.8 Hz), 8.21 (1H, d, J=8.8 Hz), 8.25 (1H, s), 8.87 (1H, s), 9.21 (1H, brs), 10.06 (1H, s).

ESI-MS: 511.53 (M+H).

[0482]The starting material was synthesized as follows.

Production example 203-1

5-(2-Amino-5-cyanopyridin-4-ylamino)indole-1-carboxylic acid phenylamide

2-Amino-4-chloro-5-cyanopyridine (200 mg, 1.30 mmol, Production example 215-3) was dissolved in ethoxyethanol (13.0 ml); 5-aminoindole-1-carboxylic acid phenylamide (408 mg, 1.62 mmol, Production example 201-2) and pyridine hydrochloride (315 mg, 2.73 mmol) were added thereto; and the reaction mixture was heated and stirred at 130 °C for 4 hours. After cooling to room temperature, the reaction mixture was partitioned between a saturated aqueous solution of sodium hydrogencarbonate and ethyl acetate; the organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Fuji Silysia NH, hexane: ethyl acetate

= 2: 3, ethyl acetate, ethyl acetate: methanol = 95: 5 in this order). The pale yellow oil obtained was solidified with diethyl ether; the crystals were suspended with diethyl ether, filtered off, washed with diethyl ether, and dried to yield the title compound (171 mg, 0.464 mmol, 35.7%) as colorless crystals.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 5.77 (1H, s), 6.40 (2H, brs), 6.74 (1H, d, J=3.6 Hz), 7.13 (1H, t, J=7.6 Hz), 7.17 (1H, dd, J=2.4, 8.8 Hz), 7.38 (2H, t, J=7.6 Hz), 7.46 (1H, d, J=2.4 Hz), 7.64 (2H, d, J=7.6 Hz), 8.04 (1H, s), 8.05 (1H, d, J=3.6 Hz), 8.20 (1H, d, J=8.8 Hz), 8.35 (1H, s), 10.04 (1H, s).

Example 204

5-(5-Cyano-2-(((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-ylamino)indole-1-carboxylic acid phenylamide

[0483] Similarly to Example 203, the title compound (73 mg, 0.13 mmol, 82%) was obtained as colorless crystals from 5-(2-amino-5-cyanopyridin-4-ylamino)indole-1-carboxylic acid phenylamide (60 mg, 0.16 mmol, Production example 203-1), triethylamine (0.056 ml, 0.41 mmol), phenyl chloroformate (0.082 ml, 0.66 mmol), and 4-(pyrrolidin-1-yl)piperidine (126 mg, 0.81 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.17-1.28 (2H, m), 1.62 (4H, m), 1.73 (2H, m), 2.07 (1H, m), 2.42 (4H, m),

2.80 (2H, m), 3.87 (2H, m), 6.75 (1H, d, J=3.6 Hz),
7.13 (1H, t, J=7.6 Hz), 7.18 (1H, d, J=8.8 Hz), 7.38
(3H, m), 7.48 (1H, s), 7.64 (2H, d, J=7.6 Hz), 8.07 (1H,
d, J=3.6 Hz), 8.20 (1H, d, J=8.8 Hz), 8.30 (1H, s),
5 8.87 (1H, s), 9.20 (1H, brs), 10.06 (1H, s).

ESI-MS: 549.48 (M+H).

Example 205

10 5-(N-(2-(3-(3-Diethylaminopropyl)ureido)-5-
cyanopyridin-4-yl)-N-methylamino)indole-1-carboxylic
acid methylamide

[0484] Similarly to Example 203, the title compound (13
mg, 0.027 mmol, 67%) was obtained as colorless crystals
from 5-(N-(2-amino-5-cyanopyridin-4-yl)-N-
15 methylamino)indole-1-carboxylic acid methylamide (13 mg,
0.041 mmol), phenyl chloroformate (0.011 ml, 0.089
mmol), triethylamine (0.014 ml, 0.10 mmol), and N,N-
diethyl-1,3-diaminopropane (0.032 ml, 0.21 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.92 (6H, t, J=7.0
20 Hz), 1.53 (2H, m), 2.41 (6H, m), 2.82 (3H, d, J=4.0 Hz),
3.14 (2H, m), 3.29 (3H, s), 6.65 (1H, d, J=3.6 Hz),
7.09 (1H, s), 7.13 (1H, dd, J=2.0, 8.8 Hz), 7.45 (1H, d,
J=2.0 Hz), 7.74 (1H, brs), 7.84 (1H, d, J=3.6 Hz), 8.10
(1H, s), 8.15 (1H, m), 8.23 (1H, d, J=8.8 Hz), 9.27 (1H,
25 s).

ESI-MS: 477.40 (M+H).

[0485]The starting material was synthesized as follows.

Production example 205-1

5- (N- (2-Amino-5-cyanopyridin-4-yl) -N-

5 methylamino)indole-1-carboxylic acid methylamide

Similarly to Production example 203, the title compound (13 mg, 0.041 mmol, 35.7%) was obtained as colorless crystals from 2-amino-4-chloro-5-cyanopyridine (27 mg, 0.18 mmol, Production example 10 215-3), 5- (N-methylamino)indole-1-carboxylic acid methylamide (30 mg, 0.15 mmol), and pyridine hydrochloride (38 mg, 0.38 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 5.77 (1H, s), 6.40 (2H, brs), 6.74 (1H, d, J=3.6 Hz), 7.13 (1H, t, J=7.6 15 Hz), 7.17 (1H, dd, J=2.4, 8.8 Hz), 7.38 (2H, t, J=7.6 Hz), 7.46 (1H, d, J=2.4 Hz), 7.64 (2H, d, J=7.6 Hz), 8.04 (1H, s), 8.05 (1H, d, J=3.6 Hz), 8.20 (1H, d, J=8.8 Hz), 8.35 (1H, s), 10.04 (1H, s).

20 Example 206

N1-Methyl-5- (2- (azetidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0486] Azetidine hydrochloride (104 mg, 1.11 mmol) and triethylamine (0.155 ml, 1.11 mmol) were added to a 25 dimethylformamide (1 ml) solution of phenyl N- (4- (1- (methylamino) carbonyl-1H-5-indolyloxy) -2-pyridyl) -N-

(phenoxycarbonyl)carbamate (116 mg, 0.222 mmol) synthesized in Production example 5-2; and the reaction mixture was stirred overnight at room temperature. The reaction mixture was partitioned between ethyl acetate and water; the organic layer was dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was washed with a solvent mixture of ether-hexane = 1: 1; and the resultant solid was filtered off to yield the title compound (50 mg) as crystals.

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 2.03-2.13 (2H, m), 2.83 (3H, d, J=6.2 Hz), 3.99 (4H, t, J=7.9 Hz), 6.54 (1H, dd, J=2.2, 6.7 Hz), 6.68 (1H, d, J=3.9 Hz), 7.03 (1H, dd, J=2.2, 8.3 Hz), 7.35 (1H, d, J=2.2 Hz), 7.41 (1H, d, J=2.2 Hz), 7.87 (1H, d, J=3.9 Hz), 8.04 (1H, d, J=6.7 Hz), 8.03-8.20 (1H, m), 8.28 (1H, t, J=8.3 Hz), 8.88 (1H, s).

Example 207

N1-Ethyl-5-(2-(azetidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0487] Similarly to Example 206, the title compound (50 mg) was obtained as white crystals from phenyl N-(4-(1-(ethylamino)carbonyl-1H-5-indolyloxy)-2-pyridyl)-N-(phenoxycarbonyl)carbamate (120 mg, 0.224 mmol) synthesized in Production example 55-1, dimethylformamide (1 ml), azetidine hydrochloride (105

mg, 1.12 mmol), and triethylamine (0.156 ml, 1.12 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 1.19 (3H, t, J=7.9 Hz), 2.04-2.13 (2H, m), 3.27-3.36 (2H, m), 3.90 (4H, t, J=7.0 Hz), 6.52 (1H, dd, J=1.9, 6.5 Hz), 6.67 (1H, d, J=3.9 Hz), 7.02 (1H, dd, J=1.9, 8.4 Hz), 7.34 (1H, d, J=1.9 Hz), 7.42 (1H, d, J=1.9 Hz), 7.90 (1H, d, J=3.9 Hz), 8.05 (1H, d, J=6.5 Hz), 8.21 (1H, t, J=6.5 Hz), 8.28 (1H, d, J=8.4 Hz), 8.88 (1H, s).

10 Example 208

N1-Cyclopropyl-5-(2-(azetidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0488] Similarly to Example 206, the title compound (80 mg) was obtained as white crystals from a mixture (228 mg) of phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)-N-(phenoxycarbonyl)carbamate and phenyl N-(4-(1-cyclopropylaminocarbonyl-1H-5-indolyl)oxy-2-pyridyl)carbamate obtained by a similar method to Example 68, N,N-dimethylformamide (2 ml), azetidine hydrochloride (194 mg, 2.07 mmol), and triethylamine (0.29 ml, 2.08 mmol).

¹H-NMR Spectrum (DMSO-d₆) δ (ppm): 0.59-0.65 (2H, m), 0.70-0.78 (2H, m), 2.03-2.13 (2H, m), 2.73-2.82 (1H, m), 3.89 (4H, t, J=7.1 Hz), 6.52 (1H, dd, J=2.0, 6.6 Hz), 6.64 (1H, d, J=3.9 Hz), 7.02 (1H, dd, J=2.0, 8.5 Hz), 7.34 (1H, d, J=2.0 Hz), 7.41 (1H, d, J=2.0 Hz), 7.87

(1H, d, J=3.9 Hz), 8.05 (1H, d, J=6.6 Hz), 8.23-8.30
(2H, m), 8.87 (1H, s).

Example 209

5 N1-Methyl-5-(2-(((4-(morpholin-4-yl)piperidin-1-
yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-
indolecarboxamide

[0489] Morpholine (228 mg, 1.64 mmol), sodium
triacetoxyborohydride (372 mg, 1.76 mmol), and acetic
10 acid (0.134 ml, 2.34 mmol) were added to a
dichloromethane (3.5 ml) solution of N1-methyl-5-(2-(4-
oxopiperidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-
indolecarboxamide (476 mg) synthesized in Example 40,
and stirred overnight at room temperature. The
15 reaction mixture was partitioned between ethyl acetate
and water; and the organic layer was dried over
anhydrous sodium sulfate. The solution was
concentrated under reduced pressure; and the residue
was purified by silica gel column chromatography (Fuji
20 Silysia NH, ethyl acetate-methanol system). The
resultant was washed with a solvent mixture of ether-
hexane = 1: 1; and the solid was filtered off to yield
the title compound (110 mg) as crystals.
MS Spectrum (ESI): 479 (M+1), 958 (2M+1).

25

Example 210

N1-Methyl-5-(2-(((4-(azetidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide

[0490] Azetidine hydrochloride (179 mg, 2.00 mmol) and
5 sodium triacetoxymethylborohydride (434 mg, 2.05 mmol) were
added to a dichloromethane (3.7 ml) solution of N1-methyl-5-(2-(4-oxopiperidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide (555 mg, 1.36 mmol)
10 synthesized in Example 40, and stirred overnight at
room temperature. The reaction mixture was partitioned
between ethyl acetate and water; and the organic layer
was dried over anhydrous sodium sulfate. The solution
was concentrated under reduced pressure; and the
15 residue was purified by use of silica gel column
chromatography (Fuji Silysia NH, ethyl acetate-methanol
system). The resultant was washed with a solvent
mixture of ether-hexane = 1: 1; and the solid was
20 filtered off to yield crystals of the title compound (5
mg), and a mixture (410 mg) including the title
compound.
MS Spectrum (ESI): 449 (M+1), 897 (2M+1).

Example 211

N1-Methyl-5-(2-(((4-(diethylamino)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide

25

[0491] Similarly to Example 209, the title compound (20 mg) was obtained as crystals from a dichloromethane (4 ml) solution of N1-methyl-5-(2-(4-oxopiperidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarbox-amide (558 mg) synthesized in Example 40, diethylamine (0.199 ml, 1.92 mmol), sodium triacetoxyborohydride (436 mg, 2.06 mmol) and acetic acid (0.157 ml, 2.74 mmol). A mixture (180 mg) including the title compound was also obtained.

MS Spectrum (ESI): 465 (M+1).

Example 212

N1-Methyl-5-(2-(((4-(4-hydroxypiperidin-1-yl)piperidin-1-yl)carbonyl)amino)pyridin-4-yloxy)-1H-1-indolecarboxamide

[0492] Similarly to Example 209, the title compound (100 mg) was obtained as crystals from a dichloromethane (3.5 ml) solution of N1-methyl-5-(2-(4-oxopiperidin-1-ylcarbonyl)-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (500 mg) synthesized in Example 40, 4-hydroxypiperidine (174 mg, 1.72 mmol), sodium triacetoxyborohydride (389 mg, 1.84 mmol) and acetic acid (0.141 mg, 2.46 mmol).

MS Spectrum (ESI): 493 (M+1), 985 (2M+1).

Example 213

N1-Propyl-5-(2-(pyrrolidin-1-ylcarbonyl)amino-4-pyridyl)oxy-1H-1-indolecarboxamide

[0493] N1-Propyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (477 mg, 1.54 mmol) was suspended in tetrahydrofuran (5 ml) at room temperature; triethylamine (0.536 mg, 3.08 mmol) and phenyl chloroformate (0.389 ml, 3.85 mmol) was added thereto while stirring; and the reaction mixture was stirred at room temperature for 2 hours. Water was added to the reaction mixture; this was subjected to extraction with ethyl acetate, washed with brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. N,N-dimethylformamide (3 ml) and pyrrolidine (0.27 ml, 3.23 mmol) was added to the residue; and the reaction mixture was stirred at room temperature overnight. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was dried over anhydrous sodium sulfate. The solution was concentrated under reduced pressure; the residue was purified by silica gel column chromatography (Fuji Silysia NH, ethyl acetate-methanol). The resultant was washed with a solvent mixture of ether: hexane=1:1; and the solid was filtered off to yield crystals (40 mg) of the title compound.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.93 (3H, t, J=7.1 Hz), 1.52-1.65 (2H, m), 1.74-1.82 (4H, m), 3.20-3.40

(6H, m), 6.56 (1H, dd, J=2.7, 6.3 Hz), 6.68 (1H, d, J=3.6 Hz), 7.04 (1H, dd, J=2.7, 7.6 Hz), 7.37 (1H, d, J=2.7 Hz), 7.44 (1H, d, J=2.7 Hz), 7.94 (1H, d, J=3.6 Hz), 8.08 (1H, d, J=6.3 Hz), 8.23 (1H, t, J=7.1 Hz),
5 8.28 (1H, d, J=7.6 Hz), 8.61 (1H, s).

[0494] The starting materials were synthesized as follows.

Production example 213-1

10 N1-Propyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide

Sodium hydride (60% in oil, 104 mg, 2.6 mmol) was gradually added at room temperature under nitrogen atmosphere to a N,N-dimethylformamide (7 ml) solution
15 of 4-(1H-5-indolyloxy)-2-pyridinamine (487 mg, 2.16 mmol, CAS No.417722-11-3) which was described in WO 02/32872. After the reaction mixture was stirred for 2 hours, phenyl N-propylcarbamate (465 mg, 2.6 mmol) was added thereto, and the reaction mixture was stirred for
20 4 hours. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was washed with water and brine, dried over anhydrous sodium sulfate. The solution was concentrated under reduced pressure; and the residue was filtrated by
25 silica gel column chromatography (Fuji Silysia NH, ethyl acetate-methanol) to yield a mixture (500 mg)

including the title compound.

MS Spectrum (ESI): 311 (M+1).

Example 214

5 N1-Isopropyl-5-(2-(pyrrolidin-1-ylcarbonyl)amino-4-
 pyridyl)oxy-1H-1-indolecarboxamide

[0495] N1-Isopropyl-5-(2-amino-4-pyridyl)oxy-1H-1-indolecarboxamide (90 mg, 0.29 mmol) was suspended in tetrahydrofuran (2 ml) at room temperature; triethylamine (0.121 mg, 0.868 mmol) and phenyl chloroformate (0.08 ml, 0.633 mmol) was added thereto while stirring; and the reaction mixture was stirred at room temperature for 2 hours. Water was added to the reaction mixture; this was subjected to extraction with ethyl acetate, washed with brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. N,N-Dimethylformamide (1 ml) and pyrrolidine (0.2 ml, 2.39 mmol) was added to the residue, and stirred at room temperature overnight. The reaction mixture was partitioned between ethyl acetate and water; the organic layer was dried over anhydrous sodium sulfate. This solution was concentrated under reduced pressure; and the residue was purified by silica gel column chromatography (Fuji Silysia NH, ethyl acetate-methanol). The resultant was washed with a solvent mixture (ether: hexane = 1: 1); and the solid was

10
15
20
25

filtered off to yield the title compound as crystals
(65 mg).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.23 (6H, d, J=6.8
Hz), 1.75-1.82 (4H, m), 3.28-3.46 (4H, m), 3.98-4.09
5 (1H, m), 6.56 (1H, dd, J=2.4, 6.0 Hz), 6.68 (1H, d,
J=3.6 Hz), 7.04 (1H, dd, J=2.4, 8.8 Hz), 7.37 (1H, d,
J=2.4 Hz), 7.42 (1H, d, J=2.4 Hz), 7.84-8.00 (2H, m),
8.08 (1H, d, J=6.0 Hz), 8.28 (1H, d, J=8.8 Hz), 8.61
(1H, s).

10

[0496] The starting materials were synthesized as
follows.

Production example 214-1

N1-Isopropyl-5-(2-amino-4-pyridyl)oxy-1H-1-
15 indolecarboxamide

Similarly to Production example 213-1, the title
compound was obtained as crystals (220 mg) from 4-(1H-
5-indolyloxy)-2-pyridinamine (482 mg, 2.16 mmol, CAS
No.417722-11-3), which was described in WO 02/32872,
20 N,N-dimethylformamide (7 ml), sodium hydride (60% in
oil, 94 mg, 2.57 mmol), and phenyl N-isopropylcarbamate
(460 mg, 2.57 mmol) under nitrogen atmosphere.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm) : 1.22 (6H, d, J=6.8
Hz), 3.97-4.08 (1H, m), 5.76 (1H, d, J=2.0 Hz), 5.85
25 (2H, s), 6.14 (1H, dd, J=2.0, 5.6 Hz), 6.67 (1H, d,
J=3.6 Hz), 7.02 (1H, dd, J=2.0, 8.8 Hz), 7.34 (1H, d,

J=2.4 Hz), 7.77 (1H, d, J=6.0 Hz), 7.94-7.96 (2H, m), 8.27 (1H, d, J=8.8 Hz).

Example 215

5 N1-Methyl-5-(2-(methylaminocarbonyl)amino-5-cyano-4-pyridyl)oxy-1H-1-indolecarboxamide

Production example 215-1

N1-Methyl-5-(2-amino-5-cyano-4-pyridyl)oxy-1H-1-indolecarboxamide.

10 [0497] 6-Amino-4-(1H-5-indolyloxy)nicotinonitrile (63 mg, 0.252 mmol) was dissolved in N,N-dimethylformamide (1 ml); and sodium hydride (60% in oil, 11.6 mg, 0.29 mmol) was gradually added thereto while stirring at room temperature. After the reaction mixture was
15 stirred for 30 minutes, phenyl N-propylcarbamate (49.5 mg, 0.277 mmol) was added thereto; and the reaction mixture was stirred for 3 hours. A saturated aqueous solution of ammonium chloride was added thereto; this was subjected to extraction with ethyl acetate, and
20 dried over anhydrous sodium sulfate. This solution was concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Fuji Silysia NH, ethyl acetate-methanol) to yield the title compound (12 mg) and N1-methyl-5-(2-amino-5-cyano-4-pyridyl)oxy-1H-1-indolecarboxamide (17 mg).
25

N1-Methyl-5-(2-(methylaminocarbonyl)amino-5-cyano-4-

pyridyl)oxy-1H-1-indolecarboxamide;

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.58 (3H, d, J=4.6 Hz), 2.86 (3H, d, J=4.6 Hz), 7.15 (1H, dd, J=2.0, 8.3 Hz), 7.20-7.28 (1H, m), 7.51 (1H, d, J=2.0 Hz), 7.93 (1H, d, J=3.0 Hz), 8.22 (1H, q, J=4.6 Hz), 8.34 (3H, d, J=8.3 Hz), 8.59 (1H, s), 9.51 (1H, s).

N1-Methyl-5-(2-amino-5-cyano-4-pyridyl)oxy-1H-1-indolecarboxamide;

[0498] ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.85 (3H, d, J=4.9 Hz), 5.59 (1H, s), 6.72 (1H, d, J=2.6 Hz), 6.87 (2H, brs), 7.13 (1H, dd, J=1.6, 8.5 Hz), 7.49 (1H, d, J=1.6 Hz), 7.92 (1H, d, J=2.6 Hz), 8.21 (1H, q, J=4.9 Hz), 8.28 (1H, s), 8.34 (1H, d, J=8.5 Hz).

Production example 215-2

2-Amino-4-chloro-5-iodopyridine

[0499] N,N-Dimethylformamide (47 ml) and N-iodosuccinimide (10.7 g, 47.6 mmol) were added to 2-amino-4-chloropyridine (4.72 g, 36.7 mmol); and the reaction mixture was stirred overnight. An aqueous solution of sodium thiosulfate and ethyl acetate were added thereto; the organic layer was separated, concentrated, and dried over anhydrous sodium sulfate. This was concentrated under reduced pressure; a solvent mixture (ether: hexane = 1: 1) was added to the

residue; and the solid was filtered off to yield the title compound (7.0g, 27.5mmol).

¹H-NMR Spectrum (CDCl₃) δ(ppm): 4.56 (2H, brs), 6.68 (1H,s), 8.32 (1H, s).

5

Production example 215-3

2-Amino-4-chloro-5-cyanopyridine

[0500] 1-Methyl-2-pyrrolidone (20 ml), zinc cyanide (0.49 g, 4.17 mmol) and
10 tetrakis(triphenylphosphine)palladium (1.3 g, 1.12 mmol) were added to 2-amino-4-chloro-5-iodopyridine (1.93 g, 7.58 mmol) synthesized in Production example 215-2; and the reaction mixture was stirred at 130-135°C for 5 hours. Approximately 0.28% of aqueous
15 ammonium (100 ml) and ethyl acetate was added to the reaction mixture; and the organic layer was separated, washed with brine, and dried over anhydrous sodium sulfate. This was concentrated under reduced pressure and the residue was filtrated by silica gel column
20 chromatography (Fuji Silysia NH, ethyl acetate). After concentration under reduced pressure, a solvent mixture (ether: hexane = 1: 1) was added to the residue, and stirred; and the solid was filtered off to yield the title compound as crystals (680 mg).
25 ¹H-NMR Spectrum (CDCl₃) δ(ppm): 5.03 (2H, brs), 6.58 (1H,s), 8.32 (1H, s).

MS Spectrum (EI): 153 (M).

Production example 215-4

6-Amino-4-(1H-5-indolyloxy)nicotinonitrile

5 [0501] 5-Hydroxyindole (313 mg, 2.35 mmol) was dissolved in dimethyl sulfoxide (3 ml); and sodium hydride (90 mg, 2.25 mmol) was gradually added thereto while stirring at room temperature. After the reaction mixture was stirred for 1 hour, 2-amino-4-chloro-5-
10 cyanopyridine (300 mg, 1.96 mmol) synthesized in Production example 215-3 was added thereto; and the reaction mixture was heated and stirred at 120 °C for 4 hours. After allowing to be cooled to room temperature, the reaction mixture was partitioned between ethyl
15 acetate and water; and the organic layer was washed with water and brine, and dried over anhydrous sodium sulfate. The solvent was distilled off; the residue was subjected to silica gel column chromatography (Fuji Silysia NH, ethyl acetate-methanol); fractions
20 containing the desired compound was concentrated under reduced pressure; ether was added thereto; and the solid was filtered off, and dried under reduced pressure to yield the title compound (95 mg, 0.38 mmol, 59%).
25 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm) : 5.59 (1H, s), 6.48 (1H, s), 6.82 (2H, s), 6.92 (1H, dd, J=2.0, 9.0 Hz),

7.40 (1H, d, J=2.0 Hz), 7.46 (1H, t, J=2.0 Hz), 7.50 (1H, d, J=9.0 Hz), 8.26 (1H, s), 11.30 (1H, s).

Example 216

5 N1-Methyl-5-(2-(pyrrolidin-1-ylcarbonyl)amino-5-cyano-4-pyridyl)oxy-1H-1-indolecarboxamide

[0502] N1-Methyl-5-(2-amino-5-cyano-4-pyridyl)oxy-1H-1-indolecarboxamide (20 mg) synthesized in Example 215 was suspended in tetrahydrofuran (0.5 ml) at room temperature; triethylamine (0.121 ml, 0.868 mmol) and phenyl chloroformate (0.08 ml, 0.633 mmol) was added thereto while stirring; and the reaction mixture was stirred at room temperature for 2 hours. Water was added to the reaction mixture; this was subjected to extraction with ethyl acetate, washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. N,N-Dimethylformamide and pyrrolidine (0.013 ml) was added to a portion of the residue (14 mg); and the reaction mixture was stirred overnight. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Fuji Silysia NH, ethyl acetate-methanol). The resultant was washed with a solvent mixture (ether: hexane = 1: 1) to

10
15
20
25

yield the title compound as crystals (6 mg).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.74 (4H, brs), 2.85 (3H, d, J=4.4 Hz), 3.15-3.40 (4H, m), 6.72 (1H, d, J=4.7 Hz), 7.15 (1H, dd, J=1.9, 8.4 Hz), 7.37 (1H, s), 7.50 (1H, d, J=1.9 Hz), 7.93 (1H, d, J=4.7 Hz), 8.20-8.26 (1H, m), 8.33 (1H, d, J=8.4 Hz), 8.63 (1H, s), 9.28 (1H, brs).

Example 217

N1-Methyl-5-(2-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino-5-cyano-4-pyridyl)oxy-1H-1-indolecarboxamide

[0503] N1-Methyl-5-(2-amino-5-cyano-4-pyridyl)oxy-1H-1-indolecarboxamide (15 mg, 0.049 mmol) synthesized in Example 215 was suspended in tetrahydrofuran (0.5 ml) at room temperature; triethylamine (17 μl, 0.122 mmol) and phenyl chloroformate (14 μl, 0.072 mmol) was added thereto while stirring; and the reaction mixture was stirred at room temperature for 2 hours. Water was added to the reaction mixture, this was subjected to extraction with ethyl acetate, washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. N,N-Dimethylformamide (0.5 ml) and 4-(1-pyrrolidinyl)piperidine (28 mg, 0.18 mmol) was added to a portion of the residue (14 mg); and the reaction mixture was stirred at room temperature

overnight. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was dried over anhydrous sodium sulfate. This was concentrated under reduced pressure; and the residue was purified by silica gel column chromatography (Fuji Silysia NH, ethyl acetate-methanol). The resultant was washed with a solvent mixture (ether: hexane = 1: 1); and the solid was filtered off to yield the title compound as crystals (6 mg).

MS Spectrum (ESI): 488 (M+1), 975 (2M+1).

Example 218

N1-Methyl-5-(2-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino-5-cyano-4-pyridyl)amino-1H-1-indolecarboxamide

[0504] N1-Methyl-5-(2-amino-5-cyano-4-pyridyl)amino-1H-1-indolecarboxamide (50 mg, 0.16 mmol) was suspended in tetrahydrofuran (1 ml) at room temperature; triethylamine (0.057 ml, 0.41mmol) and phenyl chloroformate (0.041 ml, 0.325 mmol) was added thereto while stirring; and the reaction mixture was stirred at room temperature for 2 hours. Water was added to the reaction mixture; this was subjected to extraction with ethyl acetate, washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. N,N-Dimethylformamide (0.5 ml) and 4-(1-

pyrrolidinyl)piperidine (100 mg, 0.648 mmol) was added to the residue; and the reaction mixture was stirred at room temperature overnight. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was dried over anhydrous sodium sulfate. This was concentrated under reduced pressure; and the residue was purified by silica gel column chromatography (Fuji Silysia NH, ethyl acetate-methanol) to yield the title compound (65 mg, 0.134 mmol).

MS Spectrum (ESI): 487 (M+1).

[0505] The starting materials were synthesized as follows.

Production example 218-1

N1-Methyl-5-nitro-1H-1-indolecarboxamide

Sodium hydride (60% in oil, 228 mg, 5.7 mmol) was gradually added to a N,N-dimethylformamide (0.5 ml) solution of 5-nitroindole (0.841 g, 5.19 mmol) while stirring at room temperature; phenyl N-methylcarbamate (1.02 g, 6.74 mmol) was added thereto; and the reaction mixture was stirred overnight. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was washed with water and brine, dried over anhydrous sodium sulfate. This was concentrated under reduced pressure; and the residue

was purified by silica gel column chromatography (hexane-ethyl acetate, sequentially ethyl acetate) to yield the title compound (600 mg).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.88 (3H, d, J=4.4 Hz), 6.94 (1H, d, J=3.6 Hz), 8.03 (1H, d, J=3.6 Hz), 8.15 (1H, dd, J=2.4, 9.2 Hz), 8.35-8.43 (2H, m), 8.59 (1H, d, J=2.4 Hz).

Production example 218-2

10 N1-Methyl-5-amino-1H-1-indolecarboxamide

[0506] Methanol (6 ml), water (2 ml), iron (0.32 g) and ammonium chloride (0.64 g) were added to N1-methyl-5-nitro-1H-1-indolecarboxamide (0.32 g, 1.46 mmol) synthesized in Production example 218-1; and the reaction mixture was heated to reflux for 2 hours. The reaction mixture was partitioned between ethyl acetate and water; the organic layer was washed with water and brine, and dried over anhydrous sodium sulfate. The residue was filtrated with celite, and concentrated under reduced pressure to yield the title compound (210 mg).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.79 (3H, d, J=4.4 Hz), 4.73 (2H, brs), 6.48 (1H, d, J=3.6 Hz), 6.56 (1H, dd, J=2.4, 8.8 Hz), 6.68 (1H, d, J=1.6 Hz), 7.60 (1H, d, J=3.6 Hz), 7.80-7.88 (1H, m), 7.89 (1H, d, J=8.8 Hz).

Production example 218-3N1-Methyl-5-(2-amino-5-cyano-4-pyridyl)amino-1H-1-indolecarboxamide

[0507] 2-Amino-4-chloro-5-cyanopyridine (123 mg, 0.80 mmol) synthesized in Production example 215-3, ethoxyethanol (3 ml) and pyridine hydrochloride (186 mg, 1.60 mmol) were added to N1-methyl-5-amino-1H-1-indolecarboxamide (198 mg, 1.05 mmol) synthesized in Production example 218-2; and the reaction mixture was stirred at 130 °C for 3 hours. After allowing to be cooled to room temperature, the reaction mixture was partitioned between a saturated aqueous solution of sodium hydrogencarbonate and ethyl acetate; and the organic layer was washed with water and brine, dried over anhydrous sodium sulfate. The solvent was distilled off, and the residue was purified by silica gel column chromatography (hexane-ethyl acetate, ethyl acetate in this order) to yield the title compound (110 mg, 0.359 mmol).

MS Spectrum (ESI): 307 (M+1).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.84 (3H, d, J=4.4 Hz), 5.75 (1H, s), 6.39 (2H, s), 6.66 (1H, d, J=3.2 Hz), 7.13 (1H, dd, J=2.0, 8.8 Hz), 7.42 (1H, d, J=2.0 Hz), 7.82 (1H, d, J=3.2 Hz), 8.04 (1H, s), 8.08-8.14 (1H, m), 8.23 (1H, d, J=8.8 Hz), 8.30 (1H, s).

Example 219N1-Methyl-5-(2-(3-(2-diethylaminoethyl)ureido)amino-5-cyano-4-pyridyl)amino-1H-1-indolecarboxamide

[0508] N1-Methyl-5-(2-amino-5-cyano-4-pyridyl)amino-1H-1-indolecarboxamide (36 mg, 0.12 mmol) synthesized in Production example 218-3 was suspended in tetrahydrofuran (2 ml) at room temperature; triethylamine (0.1 ml, 0.72 mmol) and phenyl chloroformate (0.037 ml, 0.29 mmol) were added thereto while stirring; and the reaction mixture was stirred at room temperature for 2 hours. Water was added to the reaction mixture; this was subjected to extraction with ethyl acetate, washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. N,N-Dimethylformamide (0.5 ml) and N,N-diethylaminoethylamine (0.1 ml) were added to the residue; and the reaction mixture was stirred at room temperature overnight. The reaction mixture was partitioned between ethyl acetate and water; the organic layer was washed with brine, dried over anhydrous sodium sulfate. This was concentrated under reduced pressure; and the residue was purified by silica gel column chromatography (Fuji Silysia NH, ethyl acetate-methanol) to yield the title compound (25 mg, 0.056 mmol).

MS Spectrum (ESI): 449 (M+1).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.93 (6H, t, J=6.8 Hz), 2.37-2.50 (6H, m), 2.84 (3H, d, J=4.4 Hz), 3.10 (2H, q, J=6.8 Hz), 6.68 (1H, d, J=3.2 Hz), 7.05 (1H, s), 7.15 (1H, dd, J=2.4, 8.8Hz), 7.44 (1H, d, J=3.2 Hz), 7.70 (1H, brs), 7.84 (1H, d, J=4.0 Hz), 8.14 (1H, d, J=4.4 Hz), 8.24 (1H, d, J=8.8 Hz), 8.25 (1H, s), 8.84 (1H, s), 9.21 (1H, s).

Example 220

10 N1-Diethyl-2-methyl-5-(2-((4-(pyrrolidin-1-yl)piperidin-1-yl)carbonyl)amino-5-cyano-4-pyridyl)amino-1H-1-indolecarboxamide

[0509] N1-Diethyl-2-methyl-5-(2-amino-5-cyano-4-pyridyl)amino-1H-1-indolecarboxamide (84 mg, 0.249 mmol) was suspended in tetrahydrofuran (1 ml) at room temperature; triethylamine (0.2 ml, 1.43 mmol) and phenyl chloroformate (0.079 ml, 0.626 mmol) were added thereto while stirring; and the reaction mixture was stirred at room temperature for 2 hours. Water was added to the reaction mixture; and this was subjected to extraction with ethyl acetate, washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. N,N-Dimethylformamide (0.5 ml) and 4-(1-pyrrolidinyl)piperidine (173 mg, 0.111 mmol) was added to a portion (80 mg) of the obtained residue (120 mg), and the reaction mixture was stirred at room

temperature overnight. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was dried over anhydrous sodium sulfate. This was concentrated under reduced pressure; and the residue was purified by silica gel column chromatography (Fuji Silysia NH, ethyl acetate-methanol system) to yield the title compound (45 mg, 0.083 mmol). MS Spectrum (ESI):543.5 (M+1).

[0510] The starting materials were synthesized as follows.

Production example 220-1

N1-Diethyl-2-methyl-5-nitro-1H-1-indolecarboxamide

Sodium hydride (60% in oil, 94 mg) was gradually added to a N,N-dimethylformamide (0.5 ml) solution of 2-methyl-5-nitroindole (0.841 g, 5.19 mmol) while stirring at room temperature; diethylcarbamoyl chloride (0.341 ml) was added thereto; and the reaction mixture was heated and stirred at 70 °C for 4 hours. After cooling down to room temperature, the reaction mixture was partitioned between ethyl acetate and water; the organic layer was washed with water and brine, and dried over anhydrous sodium sulfate. This was concentrated under reduced pressure; and the residue was purified by silica gel column chromatography (Fuji Silysia NH; hexane-ethyl acetate, ethyl acetate in this

order) to yield the title compound (420 mg).

MS Spectrum (ESI): 330 (M+55).

Production example 220-2

5 N1-Diethyl-2-methyl-5-amino-1H-1-indolecarboxamide

[0511] Methanol (8 ml), water (2 ml), iron powder (0.42 g) and ammonium chloride (0.84 g) were added to N1-methyl-5-nitro-1H-1-indolecarboxamide (415 g, 1.46 mmol) synthesized in Production example 220-1; and the
10 reaction mixture was heated to reflux for 2 hours. The reaction mixture was partitioned between ethyl acetate and water; and the organic layer was washed with water and brine, and dried over anhydrous sodium sulfate. The filtration with celite, and concentration under
15 reduced pressure yield the title compound (322 mg).

MS Spectrum (ESI): 246 (M+1).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.10 (6H, t, J=7.2 Hz), 2.28 (3H, s), 3.25-3.40 (4H, m), 4.92 (2H, brs),
6.10 (1H, t, J=0.8 Hz), 6.51 (1H, dd, J=2.4, 8.4 Hz),
20 6.64 (1H, d, J=2.4 Hz), 6.89 (1H, d, J=8.4 Hz).

Production example 220-3

N1-Diethyl-2-methyl-5-(2-amino-5-cyano-4-pyridyl)amino-1H-1-indolecarboxamide

25 [0512] 2-Amino-4-chloro-5-cyanopyridine (140 mg, 0.92 mmol) synthesized in Production example 215-3,

ethoxyethanol (2.5 ml), and pyridine hydrochloride (223 mg, 1.92 mmol) was added to N1-diethyl-2-methyl-5-amino-1H-1-indolecarboxamide (320 mg, 1.31 mmol) synthesized in Production example 220-2; and the reaction mixture was stirred at 130 °C for 3 hours. After cooling down to room temperature, the reaction mixture was partitioned between a saturated aqueous solution of sodium hydrogencarbonate and ethyl acetate; and the organic layer was washed with water and brine, and dried over anhydrous sodium sulfate. The solvent was distilled off; and the residue was purified by silica gel column chromatography (hexane-ethyl acetate, sequentially ethyl acetate) to yield the title compound (110 mg, 0.359 mmol).

MS Spectrum (ESI): 363 (M+1).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 1.05-1.20 (6H, m), 2.36 (3H, s), 3.25-3.40 (4H, m), 5.68 (1H, s), 6.35-6.37 (3H, m), 7.02 (1H, dd, J=2.0, 8.4 Hz), 7.19 (1H, d, J=8.4 Hz), 7.30 (1H, d, J=2.0 Hz), 8.01 (1H, s), 8.22 (1H, s).

Example 221

5-(5-Iodo-2-(3-methylureido)pyrimidin-4-yloxy)-1H-indole-1-carboxylic acid methylamide

[0513] Phenyl N-(5-iodo-4-(1-methylaminocarbonyl-1H-indol-5-yloxy)pyrimidin-2-yl)-N-

(phenoxycarbonyl)carbamate (597 mg, 0.919 mmol) was dissolved in dimethylformamide (3.0 ml); a 40% methanol solution of methylamine (1.0 ml) was added while stirring at 0 °C; and the reaction mixture was further stirred for 30 minutes keeping the temperature. Water (10 ml) was added to the reaction mixture after the completion of the reaction; the precipitated crystals were filtered off, washed with water, methanol and diethyl ether, and dried under warm aeration to yield the title compound as white crystals (367 mg, 0.787 mmol, 86%).

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.04 (3H, d, J=4.8 Hz), 2.85 (3H, d, J=4.0 Hz), 6.73 (1H, d, J=3.6 Hz), 7.16 (1H, dd, J=2.4, 8.8 Hz), 7.52 (1H, d, J=2.4 Hz), 7.61 (1H, m), 7.92 (1H, d, J=3.6 Hz), 8.20 (1H, m), 8.35 (1H, d, J=8.8 Hz), 8.69 (1H, s), 9.78 (1H, brs).

[0514] The starting materials were synthesized as follows.

Production example 221-1

N1-Methyl-5-(2-amino-4-pyrimidyl)oxy-1H-1-indolecarboxamide

Similarly to Production example 1-3, the title compound was obtained as white powder from 4-(1H-5-indolyloxy)-2-pyrimidinamine (413 mg, 1.83 mmol) synthesized in Production example 1-2 and phenyl N-

methylcarbamate (332 mg, 2.20 mmol) synthesized in
Production example 2-1.

¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.84 (3H, d, J=4.0
Hz), 6.06 (1H, d, J=5.6 Hz), 6.57 (2H, brs), 6.67 (1H,
5 d, J=3.6 Hz), 7.04 (1H, dd, J=2.4, 8.8 Hz), 7.36 (1H, d,
J=2.4 Hz), 7.85 (1H, d, J=3.6 Hz), 8.08 (1H, d, J=5.6
Hz), 8.14 (1H, m), 8.25 (1H, d, J=8.8 Hz).

Production example 221-2

10 N1-Methyl-5-(2-amino-5-iodo-4-pyrimidyl)oxy-1H-1-
indolecarboxamide

[0515] N1-Methyl-5-(2-amino-4-pyrimidyl)oxy-1H-1-
indolecarboxamide (302 mg, 1.07 mmol) and N-
iodosuccinimide (301 mg, 1.34 mmol) were dissolved in
15 N,N-dimethylformamide (3.0 ml); and the reaction
mixture was stirred at room temperature for 1 hour.
The reaction mixture was partitioned between ethyl
acetate and water; and the organic layer was washed
with water and brine, and was dried over anhydrous
20 magnesium sulfate. The solvent was distilled off; and
the residue was purified by silica gel column
chromatography (eluent; ethyl acetate: hexane = 2: 1)
to yield the title compound as yellow crystals (224 mg,
0.547 mmol, 51%).

25 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 2.84 (3H, d, J=4.4
Hz), 6.67 (1H, d, J=3.6 Hz), 6.72 (2H, brs), 7.04 (1H,

dd, J=2.4, 8.8Hz), 7.36 (1H, d, J=2.4 Hz), 7.85 (1H, d, J=3.6 Hz), 8.15 (1H, m), 8.24 (1H, d, J=8.8 Hz), 8.33 (1H, s).

5 Production example 221-3

Phenyl N-(5-iodo-4-(1-methylaminocarbonyl-1H-indol-5-yloxy)pyrimidin-2-yl)-N-(phenoxycarbonyl)carbamate

[0516] N1-Methyl-5-(2-amino-5-iodo-4-pyrimidyl)oxy-1H-1-indolecarboxamide (205 mg, 0.500 mmol) was suspended
10 in tetrahydrofuran (5.0 ml); triethylamine (0.209 ml, 1.50 mmol) was added thereto while stirring. The suspension was cooled with ice; phenyl chloroformate (0.188 ml, 1.50 mmol) was added thereto; and the reaction mixture was stirred at room temperature for 3
15 hours. The reaction mixture was partitioned between ethyl acetate and a saturated aqueous solution of sodium hydrogencarbonate; and the organic layer was washed with a saturated aqueous solution of sodium hydrogencarbonate and brine, and was dried over
20 anhydrous magnesium sulfate. After solvent distillation, the obtained crude product was crystallized from ethyl acetate-hexane; and the crystals were filtered off, and dried under aeration to yield the title compound as white crystals (207 mg,
25 0.319 mmol, 64%).

¹H-NMR Spectrum (CDCl₃) δ(ppm): 3.09 (3H, d, J=4.8 Hz),

5.56 (1H, m), 6.56 (1H, d, J=3.6 Hz), 6.98-7.14 (4H, m),
7.17-7.34 (6H, m), 7.36-7.42 (2H, m), 7.68 (1H, s),
8.12 (1H, d, J=8.8 Hz), 8.74 (1H, s).

5 Example 222

N1-Cyclopropyl-5-((2-(((2-chloroethylamino)carbonyl)amino)-4-pyridyl)oxy)-1H-1-indolecarboxamide









































[0517] N1-cyclopropyl-5-((2-amino-4-pyridyl)oxy)-1H-1-indolecarboxamide (400 mg, CAS No. 417722-12-4) described in W002/32872, 2-chloroethyl isocyanate (150 mg) and tetrahydrofuran (5 ml) were stirred at 80 °C for 1.5 hours. The mixture was cooled to room temperature, silica gel was added, and the solvent was
10 distilled off under reduced pressure. The silica gel was charged into a dry column packed with silica gel, and purification was performed by column chromatography (hexane : ethyl acetate = 1 : 1, followed by ethyl acetate) to yield 280 mg of a colorless powder.

20 ¹H-NMR Spectrum (DMSO-d₆) δ(ppm): 0.57-0.63 (2H, m), 0.70-0.75 (2H, m), 2.73-2.80 (1H, m), 3.42 (2H, q, J= 6.0Hz), 3.61 (2H, t, J= 6.0Hz), 6.52 (1H, dd, J= 5.6Hz, 2.4Hz), 6.65 (1H, d, J= 3.6Hz), 6.85 (1H, d, J= 2.4Hz), 7.04 (1H, dd, J= 8.8Hz, 2.4Hz), 7.35 (1H, d, J= 2.4Hz),
25 7.86 (1H, d, J= 3.6Hz), 8.04 (1H, d, J= 5.6Hz), 8.27 (1H, s), 8.28 (1H, d, J= 8.8Hz), 8.34 (1H, brs), 9.19

(1H, s).


[0518] The structural formulas of the compounds
obtained in Production examples and Examples above are
5 shown in Tables 5 to 17 below.

[0519] [Table 5]

PRODUCTION EXAMPLE1-1 	PRODUCTION EXAMPLE1-2 	PRODUCTION EXAMPLE1-3 	PRODUCTION EXAMPLE2-1 	PRODUCTION EXAMPLE2-2 
PRODUCTION EXAMPLE2-3 	PRODUCTION EXAMPLE2-4 	PRODUCTION EXAMPLE5-1 	PRODUCTION EXAMPLE5-2 	PRODUCTION EXAMPLE5-3 
PRODUCTION EXAMPLE5-4 	PRODUCTION EXAMPLE5-5 	PRODUCTION EXAMPLE8-1 	PRODUCTION EXAMPLE8-2 	PRODUCTION EXAMPLE8-3 
PRODUCTION EXAMPLE26-1 	PRODUCTION EXAMPLE27-1 	PRODUCTION EXAMPLE27-2 	PRODUCTION EXAMPLE28-1 	PRODUCTION EXAMPLE28-2 
PRODUCTION EXAMPLE29-1 	PRODUCTION EXAMPLE32-1 	PRODUCTION EXAMPLE42-1 	PRODUCTION EXAMPLE42-2 	PRODUCTION EXAMPLE42-3 
PRODUCTION EXAMPLE43-1 	PRODUCTION EXAMPLE43-2 	PRODUCTION EXAMPLE43-3 	PRODUCTION EXAMPLE43-4 	PRODUCTION EXAMPLE44-1 
PRODUCTION EXAMPLE44-2 	PRODUCTION EXAMPLE51-1 	PRODUCTION EXAMPLE51-2 	PRODUCTION EXAMPLE54-1 	PRODUCTION EXAMPLE54-2 
PRODUCTION EXAMPLE54-3 	PRODUCTION EXAMPLE55-1 	PRODUCTION EXAMPLE59-1 	PRODUCTION EXAMPLE90-1 	PRODUCTION EXAMPLE90-2 




























[0520]

[Table 6]






























[0521]

[Table 7]

<p>PRODUCTION EXAMPLE131-1</p> 	<p>PRODUCTION EXAMPLE131-2</p> 	<p>PRODUCTION EXAMPLE134-1</p> 	<p>PRODUCTION EXAMPLE135-1</p> 	<p>PRODUCTION EXAMPLE135-2</p> 
<p>PRODUCTION EXAMPLE137-1</p> 	<p>PRODUCTION EXAMPLE137-2</p> 	<p>PRODUCTION EXAMPLE1401</p> 	<p>PRODUCTION EXAMPLE140-2</p> 	<p>PRODUCTION EXAMPLE140-3</p> 
<p>PRODUCTION EXAMPLE146-1</p> 	<p>PRODUCTION EXAMPLE149-1</p> 	<p>PRODUCTION EXAMPLE151-1</p> 	<p>PRODUCTION EXAMPLE156-1</p> 	<p>PRODUCTION EXAMPLE156-2</p> 
<p>PRODUCTION EXAMPLE156-3</p> 	<p>PRODUCTION EXAMPLE159-1</p> 	<p>PRODUCTION EXAMPLE161-1</p> 	<p>PRODUCTION EXAMPLE161-2</p> 	<p>PRODUCTION EXAMPLE161-3</p> 
<p>PRODUCTION EXAMPLE163-1</p> 	<p>PRODUCTION EXAMPLE165-1</p> 	<p>PRODUCTION EXAMPLE167-1</p> 	<p>PRODUCTION EXAMPLE168-1</p> 	<p>PRODUCTION EXAMPLE168-2</p> 
<p>PRODUCTION EXAMPLE169-1</p> 	<p>PRODUCTION EXAMPLE170-1</p> 			
















[0522]

[Table 8]

PRODUCTION EXAMPLE171-1	PRODUCTION EXAMPLE172-1	PRODUCTION EXAMPLE172-2	PRODUCTION EXAMPLE174-1	PRODUCTION EXAMPLE176-1
				
PRODUCTION EXAMPLE178-1	PRODUCTION EXAMPLE178-2	PRODUCTION EXAMPLE179-1	PRODUCTION EXAMPLE179-2	PRODUCTION EXAMPLE181-1
				
PRODUCTION EXAMPLE183-1	PRODUCTION EXAMPLE184-1	PRODUCTION EXAMPLE187-1	PRODUCTION EXAMPLE187-2	PRODUCTION EXAMPLE187-3
				
PRODUCTION EXAMPLE191-1	PRODUCTION EXAMPLE191-2	PRODUCTION EXAMPLE195-1	PRODUCTION EXAMPLE195-2	PRODUCTION EXAMPLE199-1
				
PRODUCTION EXAMPLE199-2	PRODUCTION EXAMPLE200-1	PRODUCTION EXAMPLE201-1	PRODUCTION EXAMPLE202-2	PRODUCTION EXAMPLE201-3
				
PRODUCTION EXAMPLE203-1	PRODUCTION EXAMPLE205-1			
				


[0523]

[Table 9]

<p>PRODUCTION EXAMPLE213-1</p> 	<p>PRODUCTION EXAMPLE214-1</p> 	<p>PRODUCTION EXAMPLE215-1</p> 	<p>PRODUCTION EXAMPLE215-2</p> 	<p>PRODUCTION EXAMPLE215-3</p> 
<p>PRODUCTION EXAMPLE215-4</p> 	<p>PRODUCTION EXAMPLE218-1</p> 	<p>PRODUCTION EXAMPLE218-2</p> 	<p>PRODUCTION EXAMPLE218-3</p> 	<p>PRODUCTION EXAMPLE220-1</p> 
<p>PRODUCTION EXAMPLE220-2</p> 	<p>PRODUCTION EXAMPLE220-3</p> 	<p>PRODUCTION EXAMPLE221-1</p> 	<p>PRODUCTION EXAMPLE221-2</p> 	<p>PRODUCTION EXAMPLE221-3</p> 


[0524]

[Table 10]




[0525]

[Table 11]




[0526]

[Table 12]




[0527]

[Table 13]




[0528]

[Table 14]




[0529]

[Table 15]




[0530]

[Table 16]



[0531]

[Table 17]



[0532] According to the present invention, it is possible to provide novel compounds that exhibit (1) powerful inhibitory action against tube formation by vascular endothelial cells induced by VEGF or FGF and (2) powerful inhibitory action against receptor kinases for VEGF or FGF, and which are highly useful as medicines.

[0533] It should be noted that the tube formation by vascular endothelial cells is an important process in angiogenesis, and compounds having inhibitory action against them therefore has angiogenesis-inhibiting action. In addition, it is known that angiogenesis in the body progresses by the additive/synergistic effect of a multiple angiogenic factors represented by VEGF and FGF (Koolwijk P, van Erck MGM, de Vree WJA, Vermeer MA, Weich HA, Hance maaijer R, van Hinsbergh VWM, Cooperative effect of TNF-alpha, bFGF and VEGF on the formation of tubular structures of human microvascular endothelial cells in a fibrin matrix. Role of urokinase activity. J. Cell Biol., 132 P. 1177-1188, (1996)).

[0534] Therefore, the compounds of the invention which inhibit tube formation induced by VEGF or FGF produced

by cancer cells and the like are expected to exhibit powerful angiogenesis inhibition *in vivo*, and should be highly useful as angiogenesis inhibitors. Moreover, the compounds of the invention are highly useful as angiogenesis inhibitors, and are also useful as prophylactic or therapeutic agents for diseases for which angiogenesis inhibition is effective, angiogenesis inhibitors, antitumor agents, therapeutic agents for angioma, cancer metastasis inhibitors, therapeutic agents for retinal neovascularization, therapeutic agents for diabetic retinopathy, therapeutic agents for inflammatory disease, therapeutic agents for inflammatory disease selected from deformat arthritis, rheumatoid arthritis, psoriasis or delayed hypersensitivity reaction, therapeutic agents for atherosclerosis, and angiogenesis inhibition-based antitumor agents.

[0535] In addition, when using the compounds of the invention as antitumor agents, the tumors include, for example, a pancreatic cancer, a gastric cancer, a colon cancer, a breast cancer, a prostate cancer, a lung cancer, a renal cancer, a brain tumor, a blood cancer and an ovarian cancer; and the tumors to be suitably targeted are a gastric cancer, a colon cancer, a prostate cancer or a renal cancer.